


VITAL STATISTICS  
MISSISSIPPI  
1967



Mississippi State Board of Health  
Jackson, Mississippi

VITAL STATISTICS

MISSISSIPPI

1967

\*\*\*\*\*

MISSISSIPPI STATE BOARD OF HEALTH  
JACKSON, MISSISSIPPI

## TABLE OF CONTENTS

SYNOPSIS OF VITAL STATISTICS, MISSISSIPPI, 1967	Page
Introduction .....	I
Live Births .....	III
Fetal Deaths .....	VII
Deaths .....	X
Marriages .....	XX
Divorces .....	XXII
 POPULATION	
Table 1. Estimated, by County, by Race, Mississippi, July 1, 1967.....	1
 LIVE BIRTHS	
Table 2. By Place of Occurrence, and by Place of Residence of Mother, by Race, with Rates, Mississippi, 1967 .....	3
Table 3. By Place of Residence of Mother, by Race and Attendant, and and by Sex, Mississippi, 1967 .....	6
 FETAL DEATHS	
Table 4. By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex, Mississippi, 1967 .....	10
 DEATHS	
Table 5. By Cause of Death, by Race, Residence Data, Mississippi, 1967.	14
Table 6. By Place of Occurrence, and by Place of Residence, by Race, with Rates, Mississippi, 1967 .....	25
Table 7. By Cause Groups and Certain Selected Causes, by Race, Sex, and Age, with Total Rates, Residence Data, Mississippi, 1967.	28
 DEATHS UNDER 1 YEAR	
Table 8. By Selected Causes, by Race and Age, with Total Rates, Residence Data, Mississippi, 1967 .....	37

## MARRIAGES AND DIVORCES

Table 9. Marriages by Place of Licensure, and by Place of Residence of Bride, by Race of Bride; Divorces by Place of Occurrence, by Race, Mississippi, 1967 .....	41
---	----

Table 10. Divorces by Cause of Divorce, by Race, Mississippi, 1967 ...	43
--	----

Divorces by Number of Minor Children, by Race, Mississippi, 1967 .....	43
---	----

Divorces by Number of Years Married, by Race, Mississippi, 1967 .....	43
--	----

## SUMMARY TABLES

Table 11. <u>Live Births, Mississippi, 1913-1967</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1967 ..	44
--	----

Table 12. <u>Deaths, Mississippi, 1913-1967</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1967 .....	46
--	----

Table 13. <u>Maternal Deaths, Mississippi, 1913-1967</u> , by Year, by Race, with Rates, Occurrence Data 1913-1943, Residence Data 1944-1967 .....	48
--	----

Table 14. <u>Deaths Under 1 Year, Mississippi, 1917-1967</u> , by Year, by Race, with Rates, Occurrence Data 1917-1943, Residence Data 1944-1967 .....	50
--	----

Table 15. <u>Deaths Under 1 Month, Mississippi, 1924-1967</u> , by Year, by Race, with Rates, Occurrence Data 1924-1944, Residence Data 1945-1967 .....	52
---	----

Table 16. <u>Deaths from Heart Diseases, Vascular Lesions, Malignant Neoplasms, Accidents, and Influenza and Pneumonia, Mississippi, 1958-1967</u> , by Year, by Race, with Rates, Residence Data .....	53
---	----

Table 17. <u>Marriages and Divorces, Mississippi, 1926-1967</u> , by Year, by Race, Occurrence Data .....	54
--	----

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH,  
by Place of Occurrence, and by Place of Residence, by Race, with Rates,  
1967 .....

County Tables (in alphabetical order) .....	57
---	----

State Table .....	139
-------------------	-----

SYNOPSIS OF VITAL STATISTICS  
MISSISSIPPI, 1967

INTRODUCTION

The 1967 Vital Statistics Bulletin is the twenty-fifth consecutive issue prepared by Statistical Services from data obtained from records of live births, fetal deaths, deaths, marriages, and divorces collected by the office of Vital Records Registration.

Occurrence and Residence Allocation

Live births, fetal deaths, and deaths were classified by place of occurrence and by place of residence. Some of the tables contain both occurrence and residence figures, but most of the tables are based on residence data only, as these data are considered more suitable for studying public health problems.

Occurrence data were obtained from certificates filed in Mississippi at the place of the event. Residence data include transcripts of records of events that occurred in other states to Mississippi residents and exclude records filed in this state for nonresidents. Residence assignment for live births and fetal deaths was determined by the usual residence of the mother. For deaths it was determined by the usual residence of the decedent, except that deaths of inmates of institutions were reallocated to place of previous residence regardless of length of stay in the institution.

The number of transcripts received for events to Mississippi residents in other states and the number of events to nonresidents in this state are shown in Table A. Transcripts came from 37 states and the District of Columbia, but the four bordering states of Alabama, Arkansas, Louisiana, and Tennessee accounted for 87 percent of the 2,603 records sent. Mississippi filed 2,265 certificates for nonresidents. Although there was no great disparity between the number of children born in other states to Mississippians and the number born in this state to nonresidents, there was a large difference in the counts on deaths; the death transcript total exceeded that for death certificates of nonresidents by 65 percent.

Table A. Transcripts received, by place of occurrence, and certificates filed in Mississippi for nonresidents

Place of Occurrence	Total Records	Live Births	Fetal Deaths	Deaths
<u>Transcripts</u>	<u>2,603</u>	<u>1,313</u>	<u>46</u>	<u>1,244</u>
Tennessee	1,515	797	32	686
Louisiana	387	197	1	189
Alabama	321	180	5	136
Illinois	53	14	1	38
Texas	52	9	2	41
Arkansas	45	17	0	28
Florida	31	8	0	23
Other places	199	91	5	103
<u>Certificates for nonresidents</u>	<u>2,265</u>	<u>1,433</u>	<u>76</u>	<u>756</u>

Vital Statistics  
Mississippi - 1967

Marriages were classified according to place of issuance of marriage license, place of occurrence of marriage, and place of residence of bride. However, the figures on brides who were Mississippi residents are incomplete since there is no exchange among the states of transcripts of marriage records for nonresidents and the out-of-state marriages of Mississippi brides could not be included. Divorces were classified only according to place where the decree was granted, as there is no information regarding place of residence on the record.

Urban-Rural Classification

Incorporated towns with populations of 2,500 or more according to the 1960 census were considered as urban areas. All other areas were classified as rural. These categories were used for live births, fetal deaths, and deaths only. Information on marriage records is inadequate for proper classification, and divorce records do not contain this item.

Population

The estimates as of July 1, 1967, which are used in this bulletin were prepared by the Division of Sociology and Anthropology of Mississippi State University. According to these figures, the population of the state has increased by approximately 183,000 or 8.4 percent since the census of 1960. The white population has grown 13.5 percent and the nonwhite, only 1.5 percent. Although there has been a gain, comparison of the amount of this gain with the natural increase (births minus deaths) reveals a continued loss through out-migration. During the last two years, however, this loss has been limited to the nonwhite population. The new estimates for individual counties show that 50 have grown since the 1960 census, 29 have lost population, and 3 were considered to have had no change as the difference was less than 0.6 percent.

Rates

Rates and ratios were calculated from residence data on live births, fetal deaths, and deaths. Estimated populations for the state and counties by race were used to prepare rates for live births, total deaths, and deaths from specific causes (except maternal and certain diseases of early infancy). Although compilation of rates by age and sex and for cities is desirable, this could not be done because of the lack of population data for these categories. Figures on live births were used to calculate fetal death ratios and rates for maternal deaths, infant deaths, neonatal deaths, and deaths from certain diseases of early infancy, as they approximate the groups at risk more closely than do general population figures.

The base for each type of rate used in this bulletin is given in the first footnote on page 55.

Classification of Causes of Death

Causes of death were assigned according to the seventh revision (1955) of the International Statistical Classification of Diseases, Injuries, and Causes of Death. On certificates with more than one cause entered, only that indicated as the underlying cause was classified.

Vital Statistics  
Mississippi - 1967

The periodic revisions of the classification system which have taken place in the past have resulted in an unavoidable lack of comparability over the years for some causes. Since the seventh revision differs only slightly from the sixth revision used from 1949 through 1957, figures for 1949-1967 are comparable, but comparison of data by cause for these years with those for the years prior to 1949 should be made with caution.

LIVE BIRTHS (See Tables 2, 3, 11)

In 1967 there were 46,594 live births to Mississippians and the birth rate was 19.7 per 1,000 population. Both the number and the rate were 3.4 percent lower than in 1966, and the number was 10,000 or 18 percent less than in 1964, only three years earlier. The year 1967 is the sixth consecutive one in which Mississippi has experienced a decrease. For the country as a whole the percentage decline in the birth rate in 1967 was almost the same as that in Mississippi, but both in this state and in the nation there has been a slowing of the downward trend. However, the number for Mississippi in 1967 is the smallest since 1933 and the rate is the second lowest on record (the rate for 1917 was smaller and the rate for 1918 was the same as that for 1967). Family planning services which are available to all citizens through the county health departments are considered to have played an important part in the reduction which has occurred; about 15,000 patients a year request these services.

Race

For the past 43 years the number of births to nonwhites has been larger than the number to whites, and in many of these years it was considerably larger; but in 1967 the difference was relatively small with the nonwhite majority amounting to only 51 percent. Births to whites numbered 22,675 with a rate of 15.9 per 1,000 population which was the lowest recorded in the 55 years for which data are available. The count for nonwhites was 23,919, and although the rate of 25.6 far exceeded that among whites, it was the lowest since 1935. In the counties, rates for whites ranged from 6.0 in Kemper County to 21.5 in George County. For nonwhites they varied from 15.0 in Itawamba County to 35.4 in Union County.

Urban-Rural Distribution

Tabulation of births by place of residence of mother showed that 18,641 children were born to urban residents and 27,953 to rural residents. However, although only 40 percent were born to urban residents, 70 percent occurred in urban places. It was estimated from comparison of figures on place of residence and place of occurrence that over 8,000 white mothers and over 5,500 nonwhite mothers from rural areas went to cities and towns for delivery.

Attendant

Eighty-two percent of the births to Mississippi mothers in 1967 took place in hospitals or clinics which provide in-patient care. (See Table B.) A few children, 1 percent, were delivered by physicians in places other than hospitals and clinics, and for the remaining 17 percent there was no medical attendant at delivery. The

Vital Statistics  
Mississippi - 1967

majority of the nonmedical attendants were midwives who must obtain permits from the State Board of Health before they can practice. There was a large difference by race in the number and proportion of babies delivered without medical attention; for whites the total was only 75 or 0.3 percent, but for nonwhites the number was 7,878 and the percentage 32.9.

Table B. Live births by attendant and race

Attendant	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>46,594</u>	<u>100.0</u>	<u>22,675</u>	<u>100.0</u>	<u>23,919</u>	<u>100.0</u>
Physician in hospital	38,236	82.1	22,565	99.5	15,671	65.5
Physician in home	405	0.9	35	0.2	370	1.5
Midwife	7,868	16.9	56	0.2	7,812	32.7
Other	85	0.2	19	0.1	66	0.3

Information in Table C shows that much improvement has been made in securing hospitalization at delivery. The proportion of births occurring in hospitals and clinics has approximately doubled in the past twenty years, rising from 41.7 percent in 1947 to 82.1 percent in 1967. Although the rate of increase among nonwhites has been impressive, the 65.5 percent delivered in hospitals in 1967 was substantially below the national figure of 91.6 percent reported for nonwhites in 1966 (latest year available).

Table C. Percentage of live births in hospitals and clinics,  
by race, 1947, 1957, 1963-1967

Year	Total	White	Nonwhite
1947	41.7	73.8	11.0
1957	68.3	98.0	44.8
1963	74.9	99.3	53.4
1964	76.0	99.4	56.1
1965	77.3	99.4	58.8
1966	79.3	99.4	61.4
1967	82.1	99.5	65.5

Month

August was the month in which the largest number of births occurred, but September had the highest daily average. (Refer to Table D.) The smallest number and also the lowest daily average occurred in April.

Vital Statistics  
Mississippi - 1967

Table D. Live births and average per day, by month

Month	Births	Average per Day	Month	Births	Average per Day
January	4,020	130	July	4,187	135
February	3,586	128	August	4,279	138
March	3,870	125	September	4,177	139
April	3,305	110	October	4,127	133
May	3,587	116	November	3,870	129
June	3,659	122	December	3,927	127

Age of Mother

There were more mothers in the age group 20-24 years than in any other, according to the data in Table E; these mothers comprised slightly over one-third of the total. Large figures were also recorded for the age groups 15-19 and 25-29, with a total of 79 percent in the span 15-29 years of age. Teenage and sub-teenage mothers numbered 11,350 or almost one-fourth (24.4 percent) of the total. Five were only 12 years old and 45 were only 13 years old. A larger proportion of nonwhite than of white mothers were under 20 or were 30 and over. Although there was an excess of nonwhites at both ends of the childbearing age range, they apparently counter-balanced each other as the median ages of white and nonwhite mothers were very close, 23.8 and 23.5 years respectively.

Table E. Live births by age of mother and race,  
with percent of total births

Age of Mother	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	46,594	100.0	22,675	100.0	23,919	100.0
10-14	288	0.6	27	0.1	261	1.1
15-19	11,062	23.7	4,431	19.5	6,631	27.7
20-24	16,349	35.1	9,152	40.4	7,197	30.1
25-29	9,541	20.5	5,196	22.9	4,345	18.2
30-34	5,194	11.1	2,392	10.5	2,802	11.7
35-39	3,005	6.4	1,152	5.1	1,853	7.7
40-44	1,026	2.2	304	1.3	722	3.0
45-49	73	0.2	13	*	60	0.2
50-54	0	-	0	-	0	-
Unknown	56	0.1	8	*	48	0.2

\* Less than 0.06 percent.

The decline in births which has occurred in the 1960's appears to have been less pronounced among women under 25 than among those 25 and over. Comparative figures show that the proportion of nonwhite mothers who were 15-19 years of age has increased and that among whites an increase has occurred in the proportion 20-24 as well as those 15-19. However, it is likely that these changes are the result of an increase in the population in these ages rather than of a smaller decline in the fertility rate among younger mothers. Since children born during the "baby boom"

Vital Statistics  
Mississippi - 1967

after World War II are now reaching the age to have children of their own, a rise in the number of young mothers is to be expected even though the birth rate may be lower than in the past.

Sex

Males numbered 23,706 and females 22,888. The sex ratio was 1,036 males per 1,000 females; and, as usual, the ratio for whites was higher than for nonwhites, 1,081 in comparison with 994. However, the fact that there were more females than males born to nonwhites was unusual. In the 23 years for which this information is available, it has never happened among the whites and only once before, in 1959, among the nonwhites. No reason for this is known.

Plurality (See Plurality in Fetal Death section also)

Among the live births to Mississippi mothers in 1967 were 1,123 from plural deliveries. From this figure an incidence rate of 2.4 percent was calculated; for whites the rate was 1.9 and for nonwhites, 2.9. Six sets of triplets with all 18 children born alive were included in these plural deliveries; and each race group had three sets.

Immaturity

Defining immaturity as birth weight of 5 1/2 pounds or less, it was found that 4,447 or 9.5 percent of the children born to Mississippi mothers in 1967 were immature. The rate of immaturity for nonwhites (12.2 percent) was much higher than that for whites (6.8 percent).

Legitimacy

Illegitimate children numbered 8,344 in 1967 and this figure was 17.9 percent of the total. According to the statements on birth certificates, 671 or 3.0 percent of the white and 7,673 or 32.1 percent of the nonwhite babies were born to unmarried mothers. The rate for whites has increased steadily since 1957 and tripled in this ten-year period. Among nonwhites there has been a steady increase since 1949 with the rate almost doubling during the 18-year span.

Over one-third (36.7 percent) of the births to mothers under age 20 were illegitimate, and in the age groups 20 and over the percentage ranged from 9 to 14. The increase in recent years in the proportion of mothers in the age group with the highest illegitimacy rate plus the fact that their already high rate has risen even higher has been partly responsible for the overall rise in illegitimacy. An additional factor in the general increase has been an upward trend in the illegitimacy rate among mothers 20-24 years of age and this group produces more children than any other. One explanation for the rise in illegitimacy in the age groups under 25 is that there are large number of unmarried women in this group and they are less likely to use contraceptive measures (including the pill, which requires a prescription) than the married ones who in recent years apparently have used contraceptives extensively. Therefore, if the married women have reduced their birth rate to a far greater degree than the unmarried, the unmarried automatically form a larger portion of the total than in the past.

Vital Statistics  
Mississippi - 1967

FETAL DEATHS (See Table 4)

According to the regulations of the Mississippi State Board of Health, all fetal deaths, regardless of the length of gestation, are to be registered. Although this all-inclusive registration has been in effect for 16 years (prior to 1952 only the reporting of fetal deaths of 20 or more weeks gestation was required), it is known that registration of both early and late fetal deaths is still incomplete. Since the figures in this bulletin are based on incomplete information concerning fetal deaths, they could present a false picture.

Fetal deaths occurring in this state in 1967 totaled 2,321, and 2,291 records for Mississippi residents were received. Differences between these numbers and those for the previous year were relatively small. However, the fetal death ratio (fetal deaths per 1,000 live births) of 49.2 was the highest recorded thus far in Mississippi. The ratio for white residents declined slightly from 1966, but that for nonwhites increased and also was the highest on record; the figures were 45.5 and 52.7 respectively for the two race groups. Although the difference between the two groups had been comparatively small in 1966, the changes in opposite directions in 1967 caused a considerable widening of the gap.

Variations in completeness of reporting show up most clearly in the wide range of ratios among the various counties and in the differences from year to year for individual counties. In 1967 the ratios for white residents ranged from zero in four counties to 222.2 in Issaquena County and those for nonwhites from zero in four counties to 115.4 in Prentiss County. Even though factors other than variation in reporting contribute to these large differences, this may be the most important one. If the high ratios for certain counties actually represent more complete reporting and are not exaggerated, it can be estimated from them that one out of every ten pregnancies results in fetal death. Figures from previous years indicate that this figure may be as high as one out of every seven or eight.

Attendant

Table F shows that a little over 86 percent of the fetal deaths to Mississippians in 1967 were delivered in hospitals and clinics; about 3 percent were delivered by physicians in homes and other places, and midwives or other nonmedical attendants were present at 10.5 percent or 240 deliveries. Only 5 white mothers lacked medical attention at delivery, but 235 or nearly 19 percent of the nonwhite mothers were delivered by midwives and other nonmedical persons. However, the figure of 19 percent for nonwhite mothers was significantly better than that of 33 percent for live births among this group. The increased likelihood of seeking medical assistance for premature or difficult deliveries which are frequently associated with fetal death and the probably greater underreporting of fetal deaths in the home than of those attended by physicians could account for this difference.

## VIII

Vital Statistics  
Mississippi - 1967

Table F. Fetal deaths by attendant and race

<u>Attendant</u>	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total</u>	<u>2,291</u>	<u>100.0</u>	<u>1,031</u>	<u>100.0</u>	<u>1,260</u>	<u>100.0</u>
Physician in hospital	1,975	86.2	1,007	97.7	968	76.8
Physician in home	76	3.3	19	1.8	57	4.5
Midwife	224	9.8	2	0.2	222	17.6
Other	16	0.7	3	0.3	13	1.0

Evidence of improved care at delivery is shown in Table G by the increases in the proportion of fetal deaths delivered in hospitals. For several years in the first half of the 1960's no substantial progress was recorded, but in 1966 and 1967 a definite upward trend was again established. Not only were gains registered for both races, but the gap between them has narrowed considerably. The disparity is still large (21 percentage points), but is smaller than that for live births (34 percentage points).

Table G. Percentage of fetal deaths delivered in hospitals and clinics, by race, 1957, 1963-1967

Year	Total	White	Nonwhite
1957	69.5	92.2	54.8
1963	79.0	94.6	65.9
1964	79.7	92.0	70.1
1965	79.4	95.8	67.5
1966	83.7	97.5	71.9
1967	86.2	97.7	76.8

Age of Mother

Mothers in the age groups 15-19 and 20-24 experienced the lowest fetal death ratios. The ratio increased at ages 25-29 and then went up sharply with each successive age group thereafter; the ratio for mothers 10-14 years of age was between those for ages 25-29 and 30-34 but closer to that for ages 30-34. Comparisons by race reveal that ratios for nonwhites were higher than those for whites in the groups under 35 years of age, but that the reverse was true at ages 35-39 and 40-44. See Table H.

Sex

A sex ratio of 1,474 males per 1,000 females was calculated from the records with sex stated; however, the sex was undetermined for 1,074 or 47 percent of the fetal deaths. This is the highest ratio on record in this state and, as usual, was markedly larger than the sex ratio calculated for live births (1,036 in 1967). It is not known whether a great many more males than females are conceived and the males then suffer excessive mortality rates before and during delivery or whether the large number of fetal deaths of undetermined sex obscures a more equitable distribution of males and females.

Vital Statistics  
Mississippi - 1967

Table H. Fetal deaths by age of mother and race, with fetal death ratios

Age of Mother	Total		White		Nonwhite	
	No.	Ratio*	No.	Ratio*	No.	Ratio*
All Ages	<u>2,291</u>	<u>49.2</u>	<u>1,031</u>	<u>45.5</u>	<u>1,260</u>	<u>52.7</u>
10-14	17	59.0	1	#	16	61.3
15-19	422	38.1	168	37.9	254	38.3
20-24	658	40.2	329	35.9	329	45.7
25-29	437	45.8	222	42.7	215	49.5
30-34	328	63.1	146	61.0	182	65.0
35-39	274	91.2	113	98.1	161	86.9
40-44	124	120.8	44	144.7	80	110.8
45-49	15	#	5	#	10	#
Unknown	16		3		13	

\* Fetal deaths per 1,000 live births to mothers in specific age-race group.

# Ratio not calculated on less than 100 live births.

#### Plurality

The percentage of fetuses from plural deliveries was greater among fetal deaths than among live births - 5.4 percent in contrast with 2.4 percent. Although there was a difference in the plurality rates among white and nonwhite fetal deaths (4.5 percent and 6.3 percent respectively), the difference was smaller than it has been in recent years because the rate for whites was higher than usual in 1967.

Counts of plural deliveries among live births and fetal deaths showed that there were 621 such events to Mississippians in 1967; this total included 615 sets of twins and 6 sets of triplets. The children were all born alive in 87 percent of the cases, in almost 6 percent there was a combination of live births and fetal deaths, and in 7 percent all were born dead. All of the triplets were born alive. Further study revealed that mortality among fetuses from plural deliveries was about twice that among single deliveries, 10.0 percent born dead in comparison with only 4.5 percent. Differences by race were minor in 1967, but had been sizeable in previous years.

#### Length of Gestation

The stated length of gestation was 20 or more weeks for 1,122 or almost half of the fetal deaths; for 1,077 it was less than 20 weeks, and for 92 it was not given. A gestation period of 20 or more weeks was reported for 62 percent of the nonwhite but only 33 percent of the white fetal deaths. Although part of this difference can be explained by the fact that the late (20 weeks or more) fetal death ratio for nonwhites was over twice that for whites, it is also likely that the presumed better reporting of early fetal deaths among the white population diminishes the relative significance of late fetal deaths among whites.

#### Legitimacy

Comparatively more fetal deaths (22.2 percent) than live births (17.9 percent) were reported as illegitimate. The contrast was especially notable among deliveries

Vital Statistics  
Mississippi - 1967

to white mothers with the proportion illegitimate about 2 1/2 times greater for fetal deaths than for live births. Illegitimacy rates of 7.4 percent for whites and 34.3 percent for nonwhites in 1967 were the highest recorded in the six years for which legitimacy of fetal deaths has been tabulated. During this period the rate for whites has more than doubled and that for nonwhites has increased by 25 percent. These increases follow the same pattern as those among live births.

DEATHS (See Tables 5-8, 12-16)

Deaths tabulated for Mississippi residents in 1967 numbered 22,609. This count was 741 less than the total for 1966 but close to that for 1965. The death rate of 9.6 per 1,000 population was slightly less than in either of the two previous years and also closer to the rate for 1965. Figures by race showed that most of the numerical decline took place among nonwhites. The total for that group in 1967 was the smallest in the 55 years of death registration in this state. In contrast, although the death toll for white persons was less than in the previous year, it was the second highest ever reported in Mississippi. The diametrically opposite records set by the two race groups in 1967 probably resulted from changes in the racial composition of the population, as the rates by race were not unusual.

The white population of the state experienced a death rate of 9.0 with variation by county from 5.1 in DeSoto County to 16.0 in Yalobusha County. Among nonwhites the total rate was 10.5; rates in the counties ranged from 6.4 in Rankin County to 16.1 in Warren County. Sixty percent of the deaths occurring in the state took place in urban areas, but only 40 percent of the Mississippians who died were urban residents.

Natural Increase (Excess of Birth Rate over Death Rate)

There has been a downward though not steady trend in the rate of natural increase since 1956. The rate of 10.1 per 1,000 population in 1967 was only half that calculated for 1956 and the lowest since 1929. Among the white population the excess of the birth rate over the death rate amounted to only 6.9 which is the smallest ever recorded for this group in Mississippi and less than half the rate of 15.1 reported for the nonwhites. Study of natural increase by county revealed that there were more deaths than births among whites in twelve counties (Attala, Choctaw, Copiah, Franklin, Holmes, Issaquena, Jefferson, Kemper, Lawrence, Newton, Webster, and Yalobusha) and the same number of births and deaths to whites in Carroll County. The highest rates of natural increase for both racial groups were recorded in Jackson County.

Month

December and January ranked in first and second place respectively in both number of deaths and average per day. (See Table I) April had the smallest number, but the lowest daily average was in July.

Vital Statistics  
Mississippi - 1967

Table I. Deaths and average per day, by month

Month	Deaths	Average per Day	Month	Deaths	Average per Day
January	2,097	68	July	1,748	56
February	1,771	63	August	1,804	58
March	1,939	62	September	1,762	59
April	1,733	58	October	1,970	64
May	1,870	60	November	1,957	65
June	1,772	59	December	2,186	70

Age at Death

In 1967 the median age at death for the total population was 68.1, exactly two years more than it was in 1957. Medians of 70.3 years for whites and 65.2 years for nonwhites were calculated. Although there was a difference of 5.1 years between the races, the trend in recent years has been toward a narrowing of this gap. Table J shows the distribution of deaths by age and race. Fewer deaths occurred at ages 5-14 than in any other ten-year period and the largest number at ages 75-84. Excessive mortality of nonwhite infants under one year of age was demonstrated by the fact that almost 12 percent of all deaths among nonwhites were infants in comparison with only 4 percent among whites.

Table J. Deaths by age and race

Age	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
All Ages	22,609	100.0	12,816	100.0	9,793	100.0
Under 1	1,645	7.3	512	4.0	1,133	11.6
1-4	295	1.3	83	0.6	212	2.2
5-9	154	0.7	68	0.5	86	0.9
10-14	153	0.7	77	0.6	76	0.8
15-24	618	2.7	342	2.7	276	2.8
25-34	545	2.4	255	2.0	290	3.0
35-44	1,001	4.4	503	3.9	498	5.1
45-54	1,964	8.7	1,121	8.7	843	8.6
55-64	3,383	15.0	1,956	15.3	1,427	14.6
65-74	4,938	21.8	2,826	22.0	2,112	21.6
75-84	5,038	22.3	3,283	25.6	1,755	17.9
85 & over	2,863	12.7	1,789	14.0	1,074	11.0
Unknown	12	*	1	*	11	0.1

\* Less than 0.06 percent.

Vital Statistics  
Mississippi - 1967

Leading Causes of Death

The ten leading causes of death shown in Table K accounted for 80 percent of the deaths of Mississippians in 1967. The first nine are the same as those on the list for 1966, except that malignant neoplasms moved up into second place ahead of vascular lesions of the central nervous system. A group of diseases entitled "other bronchopulmonic" whose principal exclusions are influenza, pneumonia, and bronchitis replaced congenital malformations which ranked tenth in the previous year.

Table K. Ten leading causes of death by race, with percent of total deaths

Cause and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
<u>Total of ten leading causes</u>	<u>18,082</u>	<u>80.0</u>	<u>10,889</u>	<u>85.0</u>	<u>7,193</u>	<u>73.4</u>
1. Diseases of heart (400-443)	7,300	32.3	4,640	36.2	2,660	27.2
2. Malignant neoplasms (140-205)	3,136	13.9	2,061	16.1	1,075	11.0
3. Vascular lesions affecting central nervous system (330-334)	2,981	13.2	1,632	12.7	1,349	13.8
4. Accidents (800-962)	1,890	8.4	1,166	9.1	724	7.4
5. Certain diseases of early infancy (760-776)	875	3.9	308	2.4	567	5.8
6. Influenza and pneumonia (480-493)	657	2.9	310	2.4	347	3.5
7. General arteriosclerosis (450)	348	1.5	221	1.7	127	1.3
8. Diabetes mellitus (260)	332	1.5	146	1.1	186	1.9
9. Other diseases of circulatory system (451-468)	288	1.3	191	1.5	97	1.0
10. Other bronchopulmonic diseases (525-527)	275	1.2	214	1.7	61	0.6

Leading Causes by Age

Leading causes of death for each age group except infants (see infant mortality section) are shown in Table L. Although no two lists were alike and each reflects the problems distinctive to that particular age group, a number of causes appeared repeatedly. Every list contained malignant neoplasms and accidents. All except that for children 1-4 years old included diseases of the heart, and vascular lesions were important in all groups 25 and over. Children 1-4 and 5-14 and old people 65 or over were seriously affected by influenza and pneumonia.

Ill-defined and Unknown Causes

There was no satisfactory statement of cause on 2,149 or 9.7 percent of the certificates for deaths that occurred in Mississippi in 1967. Senility was the only condition mentioned on 54 of these certificates, ill-defined diseases of early infancy appeared on 522, and the remaining 1,573 records had other ill-defined conditions or no cause at all. The percentages by race were 5.0 for the white population and 15.9 for nonwhites. This problem was much more severe in some counties than in others. In

Vital Statistics  
Mississippi - 1967

Table L. Five leading causes of death by age group

Cause of Death	Age Group	No. of Deaths	% of Total
1-4 years			
<u>All Causes</u>		295	100.0
1. Accidents		100	33.9
2. Influenza and pneumonia		48	16.3
3. Congenital malformations		23	7.8
4. Malignant neoplasms		17	5.8
5. Gastritis, duodenitis, enteritis and colitis		11	3.7
5-14 years			
<u>All Causes</u>		307	100.0
1. Accidents		177	57.6
2. Malignant neoplasms		40	13.0
3. Influenza and pneumonia		12	3.9
4. Congenital malformations		9	2.9
5. Diseases of the heart		8	2.6
15-24 years			
<u>All Causes</u>		618	100.0
1. Accidents		385	62.3
2. Homicide		50	8.1
3. Malignant neoplasms		35	5.7
4. Diseases of the heart		16	2.6
4. Maternal causes		16	2.6
25-44 years			
<u>All Causes</u>		1,546	100.0
1. Accidents		418	27.0
2. Diseases of the heart		223	14.4
3. Malignant neoplasms		189	12.2
4. Homicide		148	9.6
5. Vascular lesions affecting central nervous system		97	6.3
45-64 years			
<u>All Causes</u>		5,347	100.0
1. Diseases of the heart		1,900	35.5
2. Malignant neoplasms		1,082	20.2
3. Vascular lesions affecting central nervous system		509	9.5
4. Accidents		372	7.0
5. Diabetes mellitus		96	1.8
65 years and over			
<u>All Causes</u>		12,839	100.0
1. Diseases of the heart		5,136	40.0
2. Vascular lesions affecting central nervous system		2,364	18.4
3. Malignant neoplasms		1,768	13.8
4. Influenza and pneumonia		365	2.8
5. Accidents		359	2.8

Vital Statistics  
Mississippi - 1967

Carroll County 31 percent of the death certificates were deficient in information on cause. Five other counties (Issaquena, Madison, Noxubee, Washington, and Yazoo) had rates of 20 percent or more, and in three others (Holmes, Kemper, and Quitman) rates were 15-19 percent. It is possible that the order of the leading causes of death and even the causes themselves might be changed if information could be obtained for the large number of certificates filed each year without a satisfactory statement of cause of death.

Specific Causes

Heart disease was the leading cause of death in 1967 and also in every year from 1925 to the present except 1929. It was responsible for 7,300 or 32 percent of the total deaths among Mississippians in 1967. However, both the number of deaths and the death rate of 309.2 per 100,000 population were somewhat lower than in the previous year. The rates by race were 325.2 for whites and 284.7 for nonwhites. Data presented in Table M show that an overwhelming majority of heart disease deaths were caused by the arteriosclerotic and degenerative types of disease.

Table M. Deaths from diseases of the heart, by type and race

Cause and International List Number	Total	White	Nonwhite
Diseases of the heart (400-443)	7,300	4,640	2,660
Rheumatic fever (400-402)	7	3	4
Chronic rheumatic heart disease (410-416)	75	49	26
Arteriosclerotic and degenerative heart disease (420-422)	5,619	3,983	1,636
Other diseases of the heart (430-434)	589	302	287
Hypertensive heart disease (440-443)	1,010	303	707

Malignant neoplasms moved up to second place in the list of leading causes of death in 1967 after being in third place for 20 years, 1947-1966. Although it ranked second, the death toll of 3,136 was less than half that of heart disease. Even so, approximately one out of every seven deaths was attributed to this cause. For cancer as was true for heart disease, the death rate for whites was higher than that for nonwhites - 144.4 in comparison with 115.1 per 100,000 population. The two leading sites of fatal cancers were the digestive organs and peritoneum with 922 deaths and the breast and genito-urinary organs with 892 deaths. Despite the fact that cancer is a leading cause of death in every age group except infants, relatively few deaths from this cause occur among young persons. Only nine percent of the total in 1967 were among persons under 45 years of age.

Vascular lesions affecting the central nervous system ranked third among the leading causes with a total of 2,981 deaths. This cause had been in second place in Mississippi for the previous 21 years even though in the nation as a whole it ranked third (after malignant neoplasms) during this time. The death rate of 144.4 for nonwhites was considerably higher than the rate of 114.4 for whites, and this cause still holds second place among the nonwhite population of Mississippi. Cerebral hemorrhage was the most frequent type of vascular lesion resulting in death.

Vital Statistics  
Mississippi - 1967

Chronic respiratory diseases in recent years have become increasingly important as causes of both morbidity and mortality. The most important of these diseases are shown in Table N with mortality figures for 1967 and 5, 10, and 15 years earlier. A significant decline has been recorded for only one of these diseases, tuberculosis of the respiratory system; the number of deaths dropped from 388 to 76 in the past 15 years. Malignant neoplasms of the respiratory system and emphysema which are now the two most important causes of deaths due to chronic respiratory diseases are the two for which the largest increases have been registered. Mortality from cancer of the respiratory system increased by 182 percent with 595 deaths recorded in 1967. Deaths caused by emphysema rose from 10 to 194 in the past 15 years. None of the other diseases listed has shown a consistent trend over the entire period for which comparable data are available.

Table N. Deaths from chronic respiratory diseases, 1952, 1957, 1962, 1967

Cause and International List Number	1952	1957	1962	1967
Tuberculosis of respiratory system (001-008)	388	150	119	76
Malignant neoplasms of respiratory system (160-165)	211	282	435	595
Malignant neoplasms of lung, bronchus, and trachea (162, 163)	186	247	392	556
Asthma (241)	67	59	43	50
Bronchitis, chronic and unqualified (501, 502)	19	25	40	43
Bronchitis with emphysema (502.0)	3	7	19	32
Other chronic interstitial pneumonia (525)	7	21	25	27
Bronchiectasis (526)	39	28	28	21
Emphysema (527.1)	10	43	111	194
Other diseases of lung and pleural cavity (527.0, 527.2)	8	39	36	33

Influenza and pneumonia, acute respiratory diseases which ranked in sixth place among the leading causes of death, accounted for 657 deaths in 1967. This was the lowest count since 1955 and 116 less than in the preceding year. As usual, mortality from this cause was far greater among nonwhites than among whites. Rates by race were 21.7 per 100,000 population for whites and 37.1 for nonwhites. Pneumonia alone caused 635 of these deaths, influenza with pneumonia 11, and influenza without mention of pneumonia 11.

Typhoid fever which in 1966 caused the first death of a Mississippian in four years killed three more residents in 1967. All were nonwhite males but of different age groups (10-14, 15-24, and 85 years and over).

Accidents, the fourth leading cause of death among the total population and the leading cause at ages 1-44, killed 1,890 Mississippians in 1967. This number and the death rate of 80.0 per 100,000 population were the highest ever recorded in this state although the figures for 1966, the previous record year, were nearly as high. December was the month in which the largest number of accidental deaths occurred. The average number of accidental deaths per day in 1967 was 5.2.

Vital Statistics  
Mississippi - 1967

Table 0 lists the five leading causes of accidental deaths. Those on the list for 1967 accounted for 80 percent of the total and are the same causes which have been the top five for many years. Motor vehicle accidents far outnumbered all other types and accounted for almost half of the total. The year 1967 is the fifth successive one in which all previous records have been surpassed in number of motor vehicle accident deaths occurring in the state, number to state residents, and death rate. Not only have new records been set every year for five years, but the amount of the increase has been spectacular. The total for residents in 1967 was 49 percent greater than in 1962 and the rate was 42 percent higher.

Table 0. Five leading causes of accidental deaths by race,  
with percent of total accidental deaths

Type of Accident and International List Number	Total		White		Nonwhite	
	No.	%	No.	%	No.	%
1. Motor vehicle (810-835)	931	49.2	649	55.7	282	39.0
2. Fire and explosion of combustible material (916)	193	10.2	73	6.3	120	16.6
3. Falls (900-904)	178	9.4	138	11.8	40	5.5
4. Firearm (919)	104	5.5	51	4.4	53	7.3
5. Drowning, except in water transport (929)	98	5.2	39	3.3	59	8.1

Nontransport accidents (codes 870-936, international list) caused 830 deaths in Mississippi, and a count by place of accident showed that 426 or 51 percent occurred in the home. (However, for 100 of these deaths the place of accident was not specified.) The principal causes of accidental death in the home were fire and explosion of combustible material (158 deaths), falls (101 deaths), and firearms (55 deaths). Non-transport accident fatalities were also studied by age group and type. It was found that among infants under one year of age the leading type was mechanical suffocation. For children 1-9 years old fires outranked all other types by far. The greatest hazard among older children and young adults (ages 10-24) was drowning, but firearms were also of importance in causing death among persons 15-24 years of age. In the groups 25-64 years of age no one type of accidental death greatly exceeded all others, but among the elderly, 65 years or older, over half were due to falls.

Suicide and homicide were given as the causes of 164 and 271 deaths of Mississippians in 1967. Neither of these was on the list of ten leading causes for the total population, but both ranked high in certain groups. Suicide was in tenth place among whites and in sixth place among persons 15-24 years of age. Homicide was the second leading cause among persons 15-24, fourth among those 25-44, and seventh among males and also among nonwhites. Comparison of the mortality rates for these two causes by race showed a pronounced difference. The suicide rate for white persons was six and a half times that for nonwhites, whereas the homicide rate for nonwhites was almost five times higher than that among whites.

Maternal deaths are considered to be those from deliveries and complications of pregnancy, childbirth, and the puerperium. The 36 deaths in 1967 and the maternal death rate of 7.7 per 10,000 live births were higher than the corresponding figures for 1966 (the lowest on record) but not quite as high as those recorded in 1965. The yearly fluctuations in maternal mortality in the 1960's are difficult to assess, but the

Vital Statistics  
Mississippi - 1967

figures for the three most recent years have been lower than those for the first five years of the decade. Nine maternal deaths were attributed to toxemia and five each to sepsis and hemorrhage. Although the rate of 10.9 among nonwhites in 1967 was more than twice the rate of 4.4 for whites, the gap between the two has narrowed significantly. Death rates by age and race of mother are shown in Table P, but the number of deaths in each group is small and it is not possible to attach great significance to the differences shown. However, the rates increased with age and within each age group the death rate for nonwhite mothers was higher than that for whites.

Table P. Maternal deaths by age and race, with rates

Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
All Ages	36	7.7	10	4.4	26	10.9
10-14	0	-	0	-	0	-
15-24	16	5.8	6	4.4	10	7.2
25-34	13	8.8	3	4.0	10	14.0
35-44	6	14.9	1	6.9	5	19.4
45-54	1	#	0	-	1	#

\* Deaths per 10,000 live births to mothers in specific age-race group.


# Rate not calculated on less than 100 live births.

#### Infant Mortality

There were 1,645 deaths of infants under one year of age in 1967; this total was 211 less than in the previous year and the smallest ever recorded in Mississippi. The infant death rate of 35.3 per 1,000 live births was also the smallest on record and may indicate a downward trend after about 20 years during which there was no improvement. Both race groups experienced declines, but that for the nonwhites was considerably larger. The accompanying graph of infant mortality rates by race clearly illustrates the lack of progress in bettering infant health since 1946. The line for the whites shows that although improvement continued at a slower pace until 1955 there has been very little change since then. The curve for the nonwhites shows an even worse situation, an upward trend from 1946 through 1965; however the unusual drop in 1967 is a hopeful sign.

Even though Mississippi's infant death rate in 1967 was the lowest in its history, it was still the highest in the United States, exceeding the national rate by 58 percent and that for South Carolina which had the next highest rate in the South by 28 percent. Moreover, Mississippi's race specific rates were also higher than the corresponding national figures, that for whites 16 percent higher and that for nonwhites 32 percent higher.

Vital Statistics  
Mississippi - 1967



Figures in Table Q show the variation of infant mortality rates by age and race. The death rate of 47.4 for nonwhites was more than double the rate of 22.6 for whites, and excess mortality among the nonwhites was noted in each age group. However, trends by age for the two groups were in opposite directions; rates for white infants decreased with age and those for nonwhites increased. The difference between the two was greatest for children 28 days to one year of age, yet many of the deaths in this group are due to causes which are more amenable to preventive measures than are the major causes of death in the neonatal period (first 28 days of life).

Table Q. Infant deaths by age and race, with rates

Age	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>Total under one year</u>	<u>1,645</u>	<u>35.3</u>	<u>512</u>	<u>22.6</u>	<u>1,133</u>	<u>47.4</u>
Under one day	532	11.4	214	9.4	318	13.3
One to 28 days	521	11.2	174	7.7	347	14.5
28 days to one year	592	12.7	124	5.5	468	19.6

\* Deaths per 1,000 live births.

Vital Statistics  
Mississippi - 1967

Table R lists the five leading causes of death among infants in 1967. For the second successive year congenital malformations ranked at the head of the list, but among nonwhite infants influenza and pneumonia ranked first and congenital malformations second. The first three causes shown have been on the list every year from 1949 to the present (though not in the same order) and the fourth cause, birth injuries, every year of this period except 1963. That year and 1967 are the only two in which accidents appeared on the list; during the remainder of the time the other leading cause was gastritis, duodenitis, enteritis, and colitis, except diarrhea of the newborn. However, the number of deaths due to the gastritis group of diseases dropped from 123 in 1966 to only 76 in 1967. The five leading causes were responsible for 36 percent of the total infant deaths in 1967 and for 65 percent of those with a definite cause given. Significant differences between the races were noted for two causes. The accidental death rate was twice as high among nonwhite infants as among whites, and the rate for influenza and pneumonia was eight times greater among nonwhites. Immaturity, which is no longer considered a separate classification in ranking causes of infant deaths, is still an important factor. It was mentioned with another condition or as the only cause on 617 death records, 42 percent of those for white children and 36 percent of those for nonwhites.

Table R. Leading causes of infant deaths by race, with rates

Cause of Death and International List Number	Total		White		Nonwhite	
	No.	Rate*	No.	Rate*	No.	Rate*
<u>All Causes</u>	<u>1,645</u>	<u>35.3</u>	<u>512</u>	<u>22.6</u>	<u>1,133</u>	<u>47.4</u>
1. Congenital malformations (750-759)	164	3.5	88	3.9	76	3.2
2. Postnatal asphyxia and atelectasis (762)	138	3.0	67	3.0	71	3.0
3. Influenza and pneumonia, except pneumonia of newborn (480-493)	131	2.8	14	0.6	117	4.9
4. Birth injuries (760, 761)	86	1.8	44	1.9	42	1.8
5. Accidents (800-962)	79	1.7	25	1.1	54	2.2

\* Deaths per 1,000 live births.

Postnatal asphyxia and atelectasis was the leading cause of death among infants who died in the first day of life and birth injuries ranked second. For babies who died between the ages of one and 28 days the chief cause was congenital malformations; postnatal asphyxia and atelectasis tied with pneumonia of the newborn for second place. Influenza and pneumonia far outranked all other causes of death among children 28 days to one year of age, but the gastritis group of diseases and accidents were significant as the second and third leading causes in this group.

Because 720 or 44 percent of the infant deaths records in 1967 either had no cause of death or had ill-defined causes, the data presented here regarding causes must be considered as incomplete or possibly erroneous. If correct information could be obtained for this large number of deaths, the additional knowledge could be used in planning more effective measures to prevent the excessive loss of life among infants in this state.

Vital Statistics  
Mississippi - 1967

MARRIAGES (See Tables 9, 17)

A total of 23,535 marriages occurred in Mississippi in 1967, the eighth consecutive year in which there was an increase. Even though the upward trend has been uninterrupted, there have not been any spectacular rises and in 1967 the amount of the increase was only 59. Whites registered a small decline in 1967, but this was only one of several fluctuations experienced by both races in their rising cycles. It is probable that population growth as well as changes in the age composition have contributed to the higher marriage counts of the 1960's. Approximately twice as many marriages of white as of nonwhite couples were registered in 1967, 15,752 in comparison with 7,783.

Month

June was the most popular month for weddings as evidenced by the figures in Table S. December ranked second with August a close third.

Table S. Marriages and average per day, by month

Month	Marriages	Average per Day	Month	Marriages	Average per Day
<u>Total</u>	<u>23,535</u>	<u>64</u>			
January	1,492	48	July	2,242	72
February	1,552	55	August	2,451	79
March	1,706	55	September	2,031	68
April	1,625	54	October	1,599	52
May	1,757	57	November	1,680	56
June	2,902	97	December	2,498	80

Officiant

The officiant for 72 percent of the marriages was a minister. Most of the rest were performed by justices of the peace, although mayors, judges, members of boards of supervisors, and other persons officiated at a few.

Age of Bride and Groom

The age distribution of brides and grooms presented in Table T shows that there were 9,937 teenage or subteenage brides and 4,255 of these were under the age of 18; these two figures represented 42 and 18 percent of the total brides. Two were only 12 years old, 17 were only 13 years old, and 95 were only 14. Although the number of teenage grooms (4,451) was much smaller than the number of teenage brides, they comprised 19 percent of the total or nearly one out of every five. In contrast with these young persons were 26 brides and 152 grooms who were 75 or more years of age. The oldest bride and the oldest groom were both 98 years old. Median ages at marriage were calculated by race for brides and grooms and for first as well as all marriages, and it was found that the medians for nonwhites were 0.3-0.5 of a year higher than the corresponding figures for whites and that the medians for grooms were 2.4-2.8 years higher than those for brides. The median ages calculated for all marriages were as follows: white brides 20.7, nonwhite brides 21.0, white grooms 23.1, and nonwhite grooms 23.6. Medians determined

Vital Statistics  
Mississippi - 1967

for persons marrying for the first time were found to be 1.2 years less than the figures quoted above the brides and 1.0 years less for grooms. Although in most instances the difference between the ages of the parties to the marriage was not large, several of the couples were 40 or more years apart in age, and the greatest difference was 59 years in the marriage of a woman 25 years old and a man 84 years old.

Table T. Brides and grooms by age

Age	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>23,535</u>	<u>100.0</u>	<u>23,535</u>	<u>100.0</u>
<u>Under 20</u>	<u>9,937</u>	<u>42.2</u>	<u>4,451</u>	<u>18.9</u>
12	2	*	0	-
13	17	0.1	0	-
14	95	0.4	0	-
15	810	3.4	6	*
16	1,302	5.5	83	0.4
17	2,029	8.6	582	2.5
18	2,888	12.3	1,480	6.3
19	2,794	11.9	2,300	9.8
20-24	6,952	29.5	9,795	41.6
25-34	2,986	12.7	4,572	19.4
35-44	1,702	7.2	1,921	8.2
45-54	1,131	4.8	1,259	5.3
55-64	568	2.4	884	3.8
65-74	221	0.9	492	2.1
75 & over	26	0.1	152	0.6
Unknown	12	*	9	*

\* Less than 0.06 percent.

Previous Marriages

Both the bride and the groom were marrying for the first time in 66 percent of the weddings in 1967; and, as the information in Table U shows, 72-73 percent of the individuals in each group were undertaking their first marriage. Among the previously married were 1,504 brides and 1,560 grooms who had already been married at least twice, and 15 of these people (10 brides and 5 grooms) had been married five or more times. There were over three times as many divorced as widowed persons among those being remarried.

Table U. Number of previous marriages of brides and grooms

Previous Marriages	Bride		Groom	
	No.	%	No.	%
<u>Total</u>	<u>23,535</u>	<u>100.0</u>	<u>23,535</u>	<u>100.0</u>
None	17,030	72.4	17,127	72.8
One	4,931	21.0	4,790	20.4
Two	1,224	5.2	1,243	5.3
Three or more	280	1.2	317	1.3
Not stated	70	0.3	58	0.2

Vital Statistics  
Mississippi - 1967

Nonresident Brides

There were 3,522 brides who came from other states to be married in Mississippi in 1967, and 2,858 of them married grooms who also were from other states. The marriages in which both parties were nonresidents comprised a little over 12 percent of the total or about one out of every eight, and the nonresident brides made up 15 percent of the total. These brides came from 45 of the other 49 states and the District of Columbia, and six came from foreign countries. However, 79 percent of them came from the bordering states of Alabama, Arkansas, Louisiana, and Tennessee, and 75 percent of the licenses were issued in Mississippi counties which are adjacent to these states. Nearly 20 percent of the licenses for out-of-state brides were issued in DeSoto County which is adjacent to Memphis, Tennessee.

Out-of-State Use of Mississippi Licenses

The marriage record which was put into use in Mississippi on July 1, 1956 contained more detailed information than the one used prior to that time. From these revised records it has been learned that each year some Mississippi licenses have been used for weddings performed in other states. Such marriages have not been included in the annual tabulations because they did not occur in this state and are of questionable legality, but special counts have been made each year. The count for 1967 was 83 which was the smallest number in the 11 calendar years for which tallies have been made. About two-thirds (56) of these licenses were used in Louisiana, 10 in Tennessee, 9 in Alabama, and 1-3 each in Arkansas, Illinois, Ohio, South Carolina, and Texas. Nearly all of the licenses were issued in border counties and about two-thirds in counties which border on Louisiana with Wilkinson and Adams Counties heading the list. Study of state of residence and state of occurrence of marriage revealed that in 64 cases the bride and/or groom resided in the state where the marriage took place, in 18 cases both resided in Mississippi but went elsewhere for their wedding, and in one case a bride from the state of Washington and a groom from Kentucky used a Mississippi license in Tennessee. Teenagers were involved in 48 (58 percent) of these weddings; both bride and groom were teenagers in 24, the bride only in 22, and the groom only in 2.

DIVORCES (See Tables 9, 10, 17)

The number of divorces granted in Mississippi in 1967 was 6,455. This was the largest total since 1948 and 1,003 more than were granted only four years earlier in 1963. Increases were noted for both races; the figure of 4,704 for white couples was the second highest on record in this state (1946 was the peak year), and the 1,751 divorces among nonwhites was the highest number since 1952.

Cause and Complainant

Cruel and inhuman treatment or desertion caused 91 percent of the divorces in 1967, according to information submitted by Chancery Clerks. The leading cause listed for whites was cruel and inhuman treatment (71 percent) and for nonwhites, desertion (62 percent). In cases involving white couples the wife was the complainant for 71 percent, but among nonwhite couples the husband was the complainant in slightly over half (52 percent) of the cases.

Vital Statistics  
Mississippi - 1967

Minor Children

There were no minor children involved in 42 percent of the divorces granted to white persons and 49 percent of those to nonwhites. However, a total of 8,074 children or an average of 2.3 were listed for the 3,539 couples who did have minor children. Five or more minor children each were involved in 290 divorces; in three of these there were ten children in each family, and another family had twelve children.

Duration of Marriage

The couples who obtained divorces in 1967 had had marriages lasting from less than one year up to 65 years. For 7 percent the duration was less than a year and for 41 percent it was less than 5 years, but at the other end of the range were 19 couples who had been married for 50 or more years. The two racial groups differed greatly in duration of marriage. The median length of marriage for nonwhites (13.3 years) was 2 1/2 times that for whites (5.3 years). Another illustration of the large difference is that 48 percent of the whites but only 22 percent of the nonwhites had been married for less than 5 years.

Additional Information Needed

The statistical record of divorce used in Mississippi is a form which provides only limited information because of the omission of a number of pertinent items and also because of lack of clarity regarding some of the items which do appear on the form. No data at all are available concerning place of marriage, place of residence of the couple at time of separation, their ages at marriage, their ages at divorce, previous marriages, and socio-economic factors. Information which is presently tabulated on minor children, cause of divorce, and complainant probably is erroneous because of lack of definitions and instructions regarding how such facts should be entered on the form and also because of poor arrangement of the form. These deficiencies seriously hamper meaningful study of the divorce problem on a state level as well as on a national level. Since Mississippi's record does not contain even what is considered minimum information, this state is not eligible to become part of the National Divorce Registration Area. Revision of the statistical record of divorce would be of great help in providing material for analysis of this serious social problem.




TABLE 1  
ESTIMATED POPULATION\*  
MARRIAGES IN MISSISSIPPI, 1967

July 1, 1967

The marriages occurred in Mississippi in 1967, the estimated population  
By County, by Race

County	Total	White	Nonwhite
STATE	<u>2,361,140</u>	<u>1,426,945</u>	<u>934,195</u>
Adams	38,900	21,080	17,820
Alcorn	26,810	13,650	3,160
Amite	14,910	6,630	8,280
Attala	24,310	14,830	9,480
Benton	7,650	4,520	3,130
Bolivar	57,500	20,510	36,990
Calhoun	15,530	11,510	4,020
Carroll	10,800	5,340	5,460
Chickasaw	18,830	11,590	7,240
Choctaw	8,600	6,320	2,280
Claiborne	10,040	3,070	6,970
Clarke	15,350	9,530	5,820
Clay	17,740	8,810	8,930
Coahoma	24,550	14,880	32,670
Copiah	27,430	14,020	13,410
Covington	14,670	9,760	4,910
DeSoto	36,900	22,960	13,940
Forrest	62,710	46,570	16,140
Franklin	9,270	5,770	3,500
George	12,260	10,850	1,410
Greene	8,380	6,580	1,800
Grenada	21,230	11,380	9,850
Hancock	20,580	17,700	2,880
Harrison	140,360	119,690	20,670
Hinds	223,910	126,100	97,810
Holmes	28,480	8,400	20,080
Humphreys	18,960	5,550	13,410
Issaquena	2,780	940	1,840
Itawamba	16,930	15,930	1,000
Jackson	75,010	62,800	12,210
Jasper	16,160	8,330	7,830
Jefferson	11,530	2,900	8,630
Jeff Davis	13,850	6,410	7,440
Jones	59,730	44,280	15,450
Kosciusko	11,500	5,310	6,190
Lafayette	24,930	17,290	7,640

Median ages at marriage for the first marriages were 24.9 years for women and 26.7 for men. Median ages at marriage for the second marriages were 26.0 for women and 27.9 for men. Median ages at marriage for third marriages were 27.1 for women and 29.0 for men. Median ages at marriage for fourth marriages were 28.2 for women and 30.3 for men. Median ages at marriage for fifth marriages were 29.4 for women and 31.6 for men. Median ages at marriage for sixth marriages were 30.6 for women and 33.8 for men. Median ages at marriage for seventh marriages were 31.8 for women and 35.0 for men. Median ages at marriage for eighth marriages were 33.0 for women and 37.2 for men. Median ages at marriage for ninth marriages were 34.2 for women and 38.4 for men. Median ages at marriage for tenth marriages were 35.4 for women and 40.6 for men.

Table 1 - Page 2  
 Estimated Population - Mississippi - July 1, 1967

County	Total	White	Nonwhite
Lamar	15,060	12,800	2,260
Lauderdale	76,700	52,210	24,490
Lawrence	11,390	7,840	3,550
Leake	20,500	13,160	7,340
Lee	49,160	38,440	10,720
Leflore	49,730	19,140	30,590
Lincoln	26,120	17,800	8,320
Lowndes	55,380	35,360	20,020
Madison	32,070	10,330	21,740
Marion	23,540	15,670	7,870
Marshall	24,670	8,420	16,250
Monroe	32,730	21,930	10,800
Montgomery	14,920	8,540	6,380
Neshoba	20,100	14,870	5,230
Newton	20,220	14,400	5,820
Noxubee	15,440	4,520	10,920
Oktibbeha	30,635	18,580	12,055
Panola	30,695	13,155	17,540
Pearl River	29,070	24,240	4,830
Perry	9,260	6,460	2,800
Pike	32,280	18,390	13,890
Pontotoc	18,240	15,030	3,210
Prentiss	20,370	17,970	2,400
Quitman	19,790	7,760	12,030
Rankin	39,120	25,250	13,870
Scott	23,630	15,170	8,460
Sharkey	10,140	3,230	6,910
Simpson	19,920	13,600	6,320
Smith	13,390	9,650	3,740
Stone	8,040	6,150	1,890
Sunflower	43,400	13,660	29,740
Tallahatchie	22,945	7,990	14,955
Tate	19,830	9,210	10,620
Tippah	17,750	14,610	3,140
Tishomingo	18,020	17,440	580
Tunica	15,620	4,020	11,600
Union	19,250	16,510	2,740
Walthall	14,190	7,680	6,510
Warren	45,375	26,040	19,335
Washington	78,140	35,530	42,610
Wayne	16,590	10,510	6,080
Webster	10,510	7,980	2,530
Wilkinson	11,560	3,430	8,130
Winston	17,120	9,490	7,630
Yalobusha	12,200	6,690	5,510
Yazoo	34,250	14,300	19,950

\* Prepared by Division of Sociology and Anthropology,  
 Mississippi State University.

TABLE 2  
LIVE BIRTHS  
MISSISSIPPI  
1967

By Place of Occurrence, and by Place of Residence of Mother, by Race with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
STATE	46,714	22,765	23,949	46,594	22,675	23,919	19.7	15.9	25.6
Adams	1,356	654	702	868	414	454	22.3	19.6	25.5
Alcorn	431	346	85	466	386	80	17.4	16.3	25.3
Amite	98	1	97	268	86	182	18.0	13.0	22.0
Attala	404	175	229	383	149	234	15.8	10.0	24.7
Benton	41	1	40	135	65	70	17.6	14.4	22.4
Bolivar	1,468	298	1,170	1,338	291	1,047	23.3	14.2	28.3
Calhoun	252	96	156	322	189	133	20.7	16.4	33.1
Carroll	86	6	80	182	58	124	16.8	10.9	22.7
Chickasaw	363	168	195	392	195	197	20.8	16.8	27.2
Choctaw	136	53	83	135	78	57	15.7	12.3	25.0
Claiborne	105	20	85	208	44	164	20.7	14.3	23.5
Clarke	82	49	33	312	151	161	20.3	15.8	27.7
Clay	268	111	157	334	143	191	18.8	16.2	21.4
Coahoma	1,057	313	744	1,029	246	783	21.6	16.5	24.0
Copiah	636	183	453	491	155	336	17.9	11.0	25.0
Covington	164	118	46	287	156	131	19.6	16.0	26.7
DeSoto	288	17	271	694	335	359	18.8	14.6	25.8
Forrest	1,373	978	395	1,116	693	423	17.8	14.9	26.2
Franklin	90	35	55	117	60	57	12.6	10.4	16.3
George	307	269	38	266	233	33	21.7	21.5	23.4
Greene	59	28	31	174	117	57	20.8	17.8	31.7
Grenada	384	131	253	412	156	256	19.4	13.7	26.0
Hancock	183	140	43	329	274	55	16.0	15.5	19.1
Harrison	3,468	2,720	748	3,084	2,379	705	22.0	19.9	34.1
Hinds	5,359	2,621	2,738	4,347	2,001	2,346	19.4	15.9	24.0
Holmes	539	73	466	552	83	469	19.4	9.9	23.4
Humphreys	430	56	374	409	88	321	21.6	15.8	23.9
Issaquena	13	0	13	51	9	42	18.3	9.6	22.8
Itawamba	38	29	9	207	192	15	12.2	12.0	15.0
Jackson	1,400	1,015	385	1,765	1,346	419	23.5	21.4	34.3
Jasper	198	127	71	330	121	209	20.4	14.5	26.7
Jefferson	69	12	57	194	26	168	16.8	9.0	19.5
Jeff Davis	198	105	93	263	103	160	19.0	16.1	21.5
Jones	1,779	754	1,025	943	609	334	15.8	13.8	21.6
Kemper	109	15	94	191	32	159	16.6	6.0	25.7
Lafayette	533	320	213	467	272	195	18.7	15.7	25.5

Table 2 - Page 2  
Live Births - Mississippi - 1967

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
Lamar	79	56	23	257	196	61	17.1	15.3	27.0
Lauderdale	1,951	999	952	1,415	748	667	18.4	14.3	27.2
Lawrence	102	42	60	184	97	87	16.2	12.4	24.5
Leake	215	108	107	320	138	182	15.6	10.5	24.8
Lee	1,252	988	264	968	709	259	19.7	18.4	24.2
Leflore	1,195	413	782	1,109	348	761	22.3	18.2	24.9
Lincoln	517	292	225	472	256	216	18.1	14.4	26.0
Lowndes	1,175	660	515	1,123	599	524	20.3	16.9	26.2
Madison	599	98	501	792	174	618	24.7	16.8	28.4
Marion	364	212	152	457	252	205	19.4	16.1	26.0
Marshall	342	22	320	516	129	387	20.9	15.3	23.8
Monroe	747	466	281	674	381	293	20.6	17.4	27.1
Montgomery	310	124	186	256	93	163	17.2	10.9	25.5
Neshoba	365	209	156	365	225	140	18.2	15.1	26.8
Newton	278	157	121	304	137	167	15.0	9.5	28.7
Noxubee	367	67	300	360	66	294	23.3	14.6	26.9
Oktibbeha	589	276	313	542	274	268	17.7	14.7	22.2
Panola	710	223	487	604	185	419	19.7	14.1	23.9
Pearl River	601	483	118	540	405	135	18.6	16.7	28.0
Perry	99	59	40	184	115	69	19.9	17.8	24.6
Pike	612	317	295	553	230	323	17.1	12.5	23.2
Pontotoc	318	236	82	278	209	69	15.2	13.9	21.5
Prentiss	383	331	52	336	284	52	16.5	15.8	21.7
Quitman	492	132	360	474	133	341	24.0	17.1	28.3
Rankin	85	43	42	709	476	233	18.1	18.8	16.8
Scott	416	267	149	465	253	212	19.7	16.7	25.0
Sharkey	97	11	86	255	54	201	25.1	16.7	29.1
Simpson	320	196	124	390	218	172	19.6	16.0	27.2
Smith	16	0	16	260	134	126	19.4	13.9	33.7
Stone	120	79	41	130	78	52	16.2	12.7	27.5
Sunflower	801	132	669	951	178	773	21.9	13.0	26.0
Tallahatchie	491	88	403	529	109	420	23.0	13.6	28.1
Tate	184	0	184	421	147	274	21.2	16.0	25.8
Tippah	362	273	89	300	230	70	16.9	15.7	22.3
Tishomingo	170	158	12	233	215	18	12.9	12.3	31.0
Tunica	367	45	322	387	61	326	24.8	15.2	28.1
Union	345	247	98	345	248	97	17.9	15.0	35.4
Walthall	328	160	168	268	99	169	18.9	12.9	26.0
Warren	1,372	571	801	909	453	456	20.0	17.4	23.6
Washington	1,962	658	1,304	1,851	586	1,265	23.7	16.5	29.7
Wayne	371	260	111	327	184	143	19.7	17.5	23.5
Webster	120	75	45	162	100	62	15.4	12.5	24.5
Wilkinson	230	87	143	231	58	173	20.0	16.9	21.3
Winston	332	142	190	378	153	225	22.1	16.1	29.5
Yalobusha	210	74	136	250	102	148	20.5	15.2	26.9
Yazoo	589	192	397	699	221	478	20.4	15.4	24.0
Out-of-State	1,313	1,045	268	1,433	1,135	298	-	-	-

Table 2 - Page 3  
Live Births - Mississippi - 1967

City**	Place of Occurrence			Place of Residence		
	Number			Number		
	Total	White	Non-white	Total	White	Non-white
Biloxi	2,030	1,770	260	1,291	1,082	209
Clarksdale	759	313	446	514	189	325
Cleveland	549	232	317	316	126	190
Columbus	799	443	356	593	271	322
Corinth	21	1	20	210	153	57
Greenville	1,202	533	669	961	361	600
Greenwood	824	412	412	509	206	303
Gulfport	1,394	948	446	878	604	274
Hattiesburg	1,346	978	368	728	414	314
Jackson	4,800	2,260	2,540	3,253	1,492	1,761
Laurel	1,762	739	1,023	466	226	240
McComb	492	296	196	209	105	104
Meridian	1,930	993	937	965	514	451
Natchez	1,352	654	698	536	215	321
Pascagoula	1,349	1,007	342	629	491	138
Tupelo	1,120	905	215	342	263	79
Vicksburg	1,361	569	792	550	230	320
Yazoo City	342	191	151	288	105	183
Total Urban#	32,772	18,243	14,529	18,641	9,867	8,774
Total Rural	13,942	4,522	9,420	27,953	12,808	15,145

\* Births per 1,000 population.

\*\* Cities having 10,000 population and over according to 1960 census.

# Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 3  
LIVE BIRTHS  
MISSISSIPPI  
1967

By Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total	White			Nonwhite			Sex	
		Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
STATE	46,594	22,675	22,565	35	75	23,919	15,671	370	23,706
Adams	868	414	414	0	0	454	444	5	434
Alcorn	466	386	383	0	3	80	50	1	233
Amité	268	86	86	0	0	182	87	1	134
Attala	383	149	149	0	0	234	81	0	202
Benton	135	65	64	1	0	70	30	0	67
Bolivar	1,338	291	284	3	4	1,047	685	5	676
Calhoun	322	189	189	0	0	133	95	2	159
Carroll	182	58	58	0	0	124	52	0	92
Chickasaw	392	195	193	0	2	197	143	1	213
Choctaw	135	78	78	0	0	57	35	0	75
Claiborne	208	44	44	0	0	164	102	0	102
Clarke	312	151	147	4	0	161	143	1	154
Clay	334	143	143	0	0	191	117	2	172
Coahoma	1,029	246	246	0	0	783	423	5	355
Copiah	491	155	155	0	0	336	293	2	253
Covington	287	156	156	0	0	131	114	0	148
DeSoto	694	335	335	0	0	359	251	2	330
Forrest	1,116	693	693	0	0	423	327	0	560
Franklin	117	60	60	0	0	57	44	1	61
George	266	233	233	0	0	33	26	0	138
Greene	174	117	117	0	0	57	42	0	87
Grenada	412	156	155	0	1	256	87	50	210
Hancock	329	274	271	0	3	55	38	1	155
Harrison	3,084	2,379	2,366	2	11	705	474	8	223
									1,570
									1,514

Table 3 - Page 2  
Live Births - Mississippi - 1967

County	Total	White				Nonwhite				Sex	
		Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female	
Hinds	4,347	2,001	2,000	0	1	2,346	2,154	35	157	2,198	2,149
Holmes	552	83	83	0	0	469	86	4	379	275	277
Humphreys	409	88	87	1	0	321	79	0	242	183	226
Issaquena	51	9	9	0	0	42	29	0	13	26	25
Itawamba	207	192	189	3	0	15	15	0	0	114	93
Jackson	1,765	1,346	1,339	3	4	419	343	9	67	905	860
Jasper	330	121	121	0	0	209	191	0	18	172	158
Jefferson	194	26	26	0	0	168	134	0	34	101	93
Jeff Davis	263	103	103	0	0	160	130	0	30	128	135
Jones	943	609	606	0	3	334	329	4	1	489	454
Kemper	191	32	32	0	0	159	95	2	62	108	83
Lafayette	467	272	271	0	1	195	97	0	98	248	219
Lamar	257	196	195	0	1	61	59	2	0	144	113
Lauderdale	1,415	748	748	0	0	667	645	3	19	696	719
Lawrence	184	97	96	0	1	87	66	2	19	101	83
Leake	320	138	137	0	1	182	108	3	71	168	152
Lee	968	709	708	0	1	259	221	3	35	498	470
Leflore	1,109	348	345	1	2	761	238	0	523	547	562
Lincoln	472	256	254	0	2	216	210	2	4	239	233
Lowndes	1,123	599	592	4	3	524	227	2	295	599	524
Madison	792	174	174	0	0	618	284	15	319	374	418
Marion	457	252	252	0	0	205	121	0	84	246	211
Marshall	516	129	127	1	1	387	201	1	185	268	248
Monroe	674	381	379	0	2	293	220	8	65	363	311
Montgomery	256	93	93	0	0	163	83	2	78	130	126
Neshoba	365	225	225	0	0	140	124	2	14	191	174
Newton	304	137	137	0	0	167	127	36	4	162	142
Noxubee	360	66	66	0	0	294	146	1	147	173	187
Oktibbeha	542	274	274	0	0	268	155	4	109	289	253
Panola	604	185	182	1	2	419	298	52	69	313	291

Table 3 - Page 3  
Live Births - Mississippi - 1967

County	Total	White			Nonwhite			Sex			
		Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
Pearl River	540	405	404	1	0	135	114	4	17	264	276
Perry	184	115	114	1	0	69	63	0	6	91	93
Pike	553	230	230	0	0	323	223	3	97	256	297
Pontotoc	278	209	209	0	0	69	64	0	5	146	132
Prentiss	336	284	284	0	0	52	44	0	8	172	164
Quitman	474	133	130	0	3	341	269	6	66	258	216
Rankin	709	476	472	2	2	233	215	3	15	377	332
Scott	465	253	253	0	0	212	166	7	39	231	234
Sharkey	255	54	54	0	0	201	146	2	53	125	130
Simpson	390	218	218	0	0	172	147	2	23	197	193
Smith	260	134	134	0	0	126	110	4	12	133	127
Stone	130	78	77	0	1	52	30	2	20	66	64
Sunflower	951	178	177	0	1	773	239	4	530	484	467
Tallahatchie	529	109	109	0	0	420	91	6	323	266	263
Tate	421	147	147	0	0	274	87	18	169	233	188
Tippah	300	230	227	2	1	70	54	9	7	155	145
Tishomingo	233	215	215	0	0	18	18	0	0	135	98
Tunica	387	61	61	0	0	326	158	2	166	193	194
Union	345	248	242	1	5	97	37	4	56	173	172
Walthall	268	99	92	0	7	169	105	0	64	140	128
Warren	909	453	450	1	2	456	421	1	34	463	446
Washington	1,851	586	583	2	1	1,265	867	6	392	954	897
Wayne	327	184	183	1	0	143	133	0	10	151	176
Webster	162	100	100	0	0	62	39	2	21	87	75
Wilkinson	231	58	58	0	0	173	82	0	91	115	116
Winston	378	153	153	0	0	225	95	1	129	193	185
Yalobusha	250	102	101	0	1	148	118	5	25	137	113
Yazoo	699	221	219	0	2	478	138	0	340	372	327

Table 3 - Page 4  
Live Births - Mississippi - 1967

City*	Total	White			Nonwhite			Sex	
		Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Biloxi	1,291	1,082	1,078	2	209	185	4	20	670
Clarksdale	514	189	189	0	325	263	4	58	273
Cleveland	316	126	125	1	0	190	145	0	45
Columbus	593	271	267	2	322	149	1	172	300
Corinth	210	153	152	0	1	57	37	0	20
Greenville	961	361	358	2	600	435	3	162	492
Greenwood	509	206	205	0	1	303	149	0	154
Gulfport	878	604	599	0	5	274	192	4	78
Hattiesburg	728	414	414	0	0	314	245	0	69
Jackson	3,253	1,492	1,492	0	0	1,761	1,657	31	73
Laurel	466	226	226	0	0	240	236	3	1
McComb	209	105	105	0	0	104	71	1	32
Meridian	965	514	514	0	0	451	445	2	4
Natchez	536	215	215	0	0	321	316	4	1
Pascagoula	629	491	490	0	1	138	101	3	34
Tupelo	342	263	263	0	0	79	72	2	5
Vicksburg	550	230	229	1	0	320	296	1	23
Yazoo City	288	105	104	0	1	183	65	0	118
									144

\* Cities having 10,000 population and over according to 1960 census.

TABLE 4  
FETAL DEATHS  
MISSISSIPPI  
1967

By Place of Occurrence; by Place of Residence of Mother, by Race and Attendant, and by Sex

County	Total by Occurrence	Total by Residence	White			Nonwhite			Sex		
			Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male	Female
STATE	2,321	2,291	1,031	1,007	19	5	1,260	968	57	725	492
Adams	80	54	32	32	0	0	22	22	0	0	10
Alcorn	8	13	8	8	0	0	5	3	0	4	6
Amite	1	8	2	2	0	0	6	5	0	1	3
Attala	17	17	7	7	0	0	10	5	0	4	3
Benton	4	5	1	1	0	0	4	0	2	6	8
Bolivar	58	53	13	13	0	0	40	29	2	4	1
Calhoun	6	11	6	5	1	0	5	3	1	1	3
Carroll	1	6	4	4	0	0	2	1	0	1	2
Chickasaw	22	12	6	5	1	0	6	4	0	2	6
Choctaw	3	3	1	1	0	0	2	1	0	1	0
Claiborne	2	10	0	0	0	0	10	10	0	3	6
Clarke	5	17	6	6	0	0	11	7	0	4	3
Clay	4	9	3	3	0	0	6	4	0	2	2
Coahoma	52	52	6	6	0	0	46	33	0	13	35
Copiah	13	19	3	3	0	0	16	12	1	3	9
Covington	3	10	3	3	0	0	7	5	0	2	2
DeSoto	4	12	7	7	0	0	5	2	0	3	4
Forrest	32	35	11	11	0	0	24	21	0	3	18
Franklin	1	4	2	2	0	0	0	2	0	0	0
George	4	3	3	3	0	0	0	0	0	1	0
Greene	4	8	4	4	0	0	4	2	0	2	5
Grenada	19	20	4	4	0	0	16	10	0	4	5
Hancock	4	6	6	6	0	0	0	0	0	0	2
Harrison	244	207	155	155	0	0	52	43	7	0	145

Table 4 - Page 2  
Fetal Deaths - Mississippi - 1967

County	Total by Occurrence	White						Nonwhite						Sex	
		Total by Residence	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Total	Male	Female	Unde- termined		
Hinds	561	401	209	205	4	0	192	181	5	6	63	37	301	9	9
Holmes	25	28	3	2	1	0	25	10	4	11	11	8	0	5	5
Humphreys	13	23	4	4	0	0	19	13	2	4	13	5	0	2	2
Issaquena	0	4	2	2	0	0	2	2	0	0	1	1	0	3	0
Itawamba	0	5	5	4	0	1	0	0	0	0	2	0	0	3	0
Jackson	29	59	42	41	1	0	17	15	0	2	15	14	30	5	30
Jasper	9	13	5	4	1	0	8	6	2	0	3	5	5	5	5
Jefferson	2	5	1	1	0	0	4	3	0	1	2	2	1	1	1
Jeff Davis	5	8	0	0	0	0	8	6	0	2	4	2	2	2	2
Jones	84	40	16	12	4	0	24	23	0	1	10	9	21	9	21
Kemper	7	11	1	1	0	0	10	6	0	4	7	2	2	2	2
Lafayette	14	10	4	4	0	0	6	1	0	5	7	2	1	1	1
Lamar	1	8	3	2	1	0	5	5	0	0	3	3	2	2	2
Lauderdale	108	81	52	51	1	0	29	28	0	1	24	15	42	3	42
Lawrence	2	6	3	3	0	0	3	3	0	0	3	0	3	3	3
Leake	3	14	5	5	0	0	9	9	0	0	5	2	7	7	7
Lee	25	33	14	14	0	0	19	16	0	3	14	15	4	4	4
Leflore	57	59	18	14	2	2	41	20	7	14	22	15	22	22	22
Lincoln	19	29	13	13	0	0	16	15	0	1	12	4	13	4	13
Lowndes	34	34	14	14	0	0	20	13	0	7	13	9	12	9	12
Madison	37	52	17	16	1	0	35	21	3	11	8	12	32	7	32
Marion	9	14	6	6	0	0	8	6	0	2	3	4	7	7	7
Marshall	18	28	5	5	0	0	23	15	1	7	16	10	2	2	2
Monroe	20	22	5	5	0	0	17	15	0	2	13	5	4	4	4
Montgomery	9	7	2	2	0	0	5	4	1	0	0	3	2	2	2
Neshoba	18	19	5	5	0	0	14	13	0	1	9	7	3	3	3
Newton	27	27	15	15	0	0	12	7	4	1	7	5	17	2	17
Noxubee	11	10	0	0	0	0	10	9	0	1	5	3	1	2	2
Oktibbeha	14	15	4	4	0	0	11	7	0	4	8	6	1	1	1
Panola	21	22	4	4	0	0	14	14	2	2	11	11	0	0	0

Table 4 - Page 3  
Fetal Deaths - Mississippi - 1967

County	Total by Occurrence	Total by Residence	White			Nonwhite			Sex		
			Total Hosp.	Phys. in Hosp.	Phys. in Home	Total Hosp.	Phys. in Hosp.	Phys. in Home	Male	Female	Under- determined
Pearl River	7	10	8	8	0	2	4	0	5	3	2
Perry	3	8	4	4	0	4	4	0	0	1	0
Pike	32	31	8	8	0	23	17	2	4	12	10
Pontotoc	4	2	1	1	0	1	1	0	0	2	0
Prentiss	17	11	5	5	0	6	6	0	0	4	3
Quitman	8	12	1	1	0	0	11	11	0	6	5
Rankin	1	52	35	1	0	17	17	0	0	3	4
Scott	6	12	5	5	0	7	6	0	1	3	6
Sharkey	1	8	4	4	0	4	3	0	1	3	4
Simpson	11	18	9	9	0	9	9	0	0	6	8
Smith	0	8	3	3	0	5	5	0	0	2	1
Stone	2	2	1	1	0	1	1	0	0	1	0
Sunflower	38	44	5	5	0	0	18	3	18	28	12
Tallahatchie	18	19	1	1	0	0	18	10	0	8	1
Tate	4	15	3	3	0	0	12	8	1	3	1
Tippah	36	32	25	25	0	0	7	4	3	5	19
Tishomingo	5	5	5	5	0	0	0	0	0	5	8
Tunica	12	11	0	0	0	0	11	7	0	4	0
Union	13	8	6	6	0	2	1	0	1	3	4
Walhall	16	13	7	5	0	2	6	2	0	4	2
Warren	122	76	45	45	0	0	31	28	2	13	56
Washington	126	125	57	57	0	0	68	60	3	5	72
Wayne	15	14	8	8	0	0	6	6	0	0	3
Webster	2	4	2	2	0	0	2	1	0	1	0
Wilkinson	6	9	2	2	0	0	7	4	0	3	4
Winston	9	11	4	4	0	0	7	6	0	1	2
Yalobusha	6	9	3	3	0	0	6	5	0	1	2
Yazoo	28	41	7	7	0	0	34	16	0	18	14

Table 4 - Page 4  
Fetal Deaths - Mississippi - 1967

City*	Total by Occurrence	Total by Residence	White			Nonwhite			Sex		
			Total	Phys. in Hosp.	Phys. in Home	Other	Total	Phys. in Hosp.	Phys. in Home	Other	Male
Biloxi	193	132	105	0	0	0	27	25	0	2	10
Clarksdale	42	23	3	0	0	0	20	17	0	3	11
Cleveland	18	8	2	0	0	0	6	4	0	2	4
Columbus	23	16	8	0	0	0	8	6	0	2	3
Corinth	2	4	1	0	0	0	3	1	0	2	6
Greenville	108	86	45	0	0	0	41	35	2	4	25
Greenwood	38	25	8	7	1	0	17	12	1	4	14
Gulfport	50	49	26	26	0	0	23	17	2	4	12
Hattiesburg	31	26	9	9	0	0	17	15	0	2	12
Jackson	478	351	191	187	4	0	160	152	4	4	53
Laurel	82	23	7	5	2	0	16	15	0	1	5
McComb	23	5	0	0	0	0	5	1	2	2	2
Meridian	107	64	40	39	1	0	24	24	0	0	19
Natchez	80	40	27	27	0	0	13	13	0	0	9
Pascagoula	26	14	10	10	0	0	4	3	0	1	3
Tupelo	18	9	6	6	0	0	3	3	0	0	5
Vicksburg	121	51	29	29	0	0	22	20	1	1	4
Yazoo City	10	18	4	4	0	0	14	8	0	6	9
											8

\* Cities having 10,000 population and over according to 1960 census.

TABLE 5  
DEATHS  
MISSISSIPPI  
1967

By Cause of Death, by Race, Residence Data

International List Number*	Cause of Death**	Total	White	Non-white
<u>001-999</u>	<u>TOTAL ALL CAUSES</u>	<u>22,609</u>	<u>12,816</u>	<u>9,793</u>
<u>001-019</u>	<u>Tuberculosis, all forms</u>	<u>77</u>	<u>36</u>	<u>41</u>
002	Pulmonary	68	30	38
003	Pleural	5	4	1
004	Primary tuberculosis complex with symptoms	1	0	1
008	Unspecified site	2	2	0
018	Other organs	1	0	1
<u>020-029</u>	<u>Syphilis and its sequelae</u>	<u>22</u>	<u>11</u>	<u>11</u>
022	Aneurysm of aorta	15	10	5
023	Other cardiovascular syphilis	6	1	5
026	Other syphilis of central nervous system	1	0	1
040	Typhoid fever	3	0	3
042	Other Salmonella infections	2	1	1
<u>045-048</u>	<u>Dysentery, all forms</u>	<u>7</u>	<u>1</u>	<u>6</u>
045	Bacillary	1	0	1
046	Amebiasis	5	1	4
048	Unspecified forms	1	0	1
049	Food poisoning (infections and intoxication)	1	0	1
050	Scarlet fever	1	0	1
051	Streptococcal sore throat	1	0	1
053	Septicemia and pyemia	34	16	18
054	Bacterial toxemia	1	0	1
056	Whooping cough	3	1	2
057	Meningococcal infections	6	3	3
061	Tetanus	8	4	4
072	Leptospirosis	1	0	1
080	Acute poliomyelitis	1	0	1
081	Late effects of acute poliomyelitis	1	1	0
082	Acute infectious encephalitis	4	1	3
083	Late effects of acute infectious encephalitis	2	2	0
085	Measles	3	1	2
087	Chickenpox	2	2	0
088	Herpes zoster	1	1	0
089	Mumps	1	1	0
092	Infectious hepatitis	13	8	5
096	Other diseases attributable to viruses	3	1	2
130	Infestation with worms of other, mixed, and unspecified type	1	0	1
132	Actinomycosis	1	0	1
134	Other fungus infections	9	6	3
138	Other infective and parasitic diseases	2	1	1
<u>140-205</u>	<u>Malignant neoplasms</u>	<u>3,136</u>	<u>2,061</u>	<u>1,075</u>
140-148	Buccal cavity and pharynx	59	48	11
140	Lip	1	1	0

Table 5 - Page 2  
Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
141	Tongue	5	5	0
142	Salivary gland	5	3	2
143	Floor of mouth	4	3	1
144	Other parts of mouth and mouth unspecified	14	12	2
145	Oral mesopharynx	3	3	0
146	Nasopharynx	9	7	2
147	Hypopharynx	3	2	1
148	Pharynx, unspecified	15	12	3
<u>150-159</u>	<u>Digestive organs and peritoneum</u>	<u>922</u>	<u>576</u>	<u>346</u>
150	Esophagus	48	31	17
151	Stomach	190	82	108
152	Small intestine, including duodenum	7	5	2
153	Large intestine, except rectum	280	189	91
154	Rectum	72	48	24
155	Biliary passages and liver (stated to be primary site)	42	35	7
156	Liver (secondary and unspecified)	62	33	29
157	Pancreas	199	139	60
158	Peritoneum	8	7	1
159	Unspecified digestive organs	14	7	7
<u>160-165</u>	<u>Respiratory system</u>	<u>595</u>	<u>445</u>	<u>150</u>
160	Nose, nasal cavities, middle ear, and accessory sinuses	7	4	3
161	Larynx	22	17	5
162	Bronchus and trachea, and lung specified as primary	214	162	52
163	Lung, unspecified as to whether primary or secondary	342	257	85
164	Mediastinum	3	1	2
165	Thoracic organs (secondary)	7	4	3
<u>170-181</u>	<u>Breast and genito-urinary organs</u>	<u>892</u>	<u>502</u>	<u>390</u>
170	Breast	244	151	93
171	Cervix uteri	124	57	67
172	Corpus uteri	18	9	9
173	Other parts of uterus, including choriocarcinoma	1	1	0
174	Uterus, unspecified	59	23	36
175	Ovary, Fallopian tube, and broad ligament	75	50	25
176	Other and unspecified female genital organs	9	5	4
177	Prostate	220	117	103
178	Testis	3	3	0
179	Other and unspecified male genital organs	4	1	3
180	Kidney	50	31	19
181	Bladder and other urinary organs	85	54	31
<u>190-199</u>	<u>Other and unspecified sites</u>	<u>379</u>	<u>274</u>	<u>105</u>
190	Melanoma of skin	29	25	4
191	Skin, other	31	27	4
192	Eye	6	3	3
193	Brain and other parts of nervous system	99	78	21
194	Thyroid gland	11	7	4
195	Other endocrine glands	2	1	1
196	Bone	22	14	8

Table 5 - Page 3  
 Deaths - Mississippi - 1967

International List Number*	Cause of Death***	Total	White	Non-white
197	Connective tissue	15	10	5
198	Lymph nodes (secondary and unspecified)	2	2	0
199	Other and unspecified sites	162	107	55
<u>200-205</u>	<u>Lymphatic and hematopoietic tissues</u>	<u>289</u>	<u>216</u>	<u>73</u>
200	Lymphosarcoma and reticulosarcoma	60	45	15
201	Hodgkin's disease	24	19	5
202	Other forms of lymphoma (reticulososis)	10	9	1
203	Multiple myeloma (plasmacytoma)	42	32	10
204	Leukemia and aleukemia	152	111	41
205	Mycosis fungoides	1	0	1
<u>210-229</u>	<u>Benign neoplasms</u>	<u>20</u>	<u>13</u>	<u>7</u>
211	Other parts of digestive system	1	1	0
212	Respiratory system	4	4	0
214	Uterine fibromyoma	3	0	3
215	Uterus, other	1	0	1
216	Ovary	2	1	1
223	Brain and other parts of nervous system	7	5	2
224	Endocrine glands	1	1	0
225	Bone and cartilage	1	1	0
<u>230-239</u>	<u>Neoplasm of unspecified nature</u>	<u>19</u>	<u>7</u>	<u>12</u>
230	Digestive organs	2	0	2
231	Respiratory organs	3	1	2
236	Other genito-urinary organs	2	0	2
237	Brain and other parts of nervous system	12	6	6
241	Asthma	50	29	21
244	Allergic eczema	1	0	1
250	Simple goiter	1	0	1
252	Thyrotoxicosis with or without goiter	9	2	7
253	Myxedema and cretinism	1	1	0
260	Diabetes mellitus	332	146	186
270	Disorders of pancreatic internal secretion other than diabetes mellitus	2	0	2
271	Diseases of parathyroid gland	1	1	0
272	Diseases of pituitary gland	1	0	1
273	Diseases of thymus gland	1	1	0
286	Other avitaminoses and nutritional deficiency states	26	15	11
287	Obesity, not specified as of endocrine origin	11	7	4
288	Gout	1	1	0
289	Other metabolic diseases	12	10	2
<u>290-293</u>	<u>Anemias</u>	<u>48</u>	<u>12</u>	<u>36</u>
290	Pernicious and other hyperchromic anemias	1	1	0
291	Iron deficiency anemias (hypochromic anemias)	4	1	3
292	Other anemias of specified type	20	5	15
293	Unspecified type	23	5	18
294	Polycythemia	3	3	0
295	Hemophilia	2	1	1
296	Purpura and other hemorrhagic conditions	7	6	1
297	Agranulocytosis	1	1	0
298	Diseases of spleen	3	2	1
299	Other diseases of blood and blood-forming organs	1	1	0

Table 5 - Page 4  
Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
304	Senile psychosis	4	2	2
305	Presenile psychosis	1	1	0
307	Alcoholic psychosis	4	3	1
309	Other and unspecified psychoses	10	10	0
322	Alcoholism	24	16	8
323	Other drug addiction	2	2	0
325	Mental deficiency	5	4	1
<u>330-334</u>	<u>Vascular lesions affecting central nervous system</u>	<u>2,981</u>	<u>1,632</u>	<u>1,349</u>
330	Subarachnoid hemorrhage	95	58	37
331	Cerebral hemorrhage	1,748	872	876
332	Cerebral embolism and thrombosis	784	515	269
334	Other and ill-defined	354	187	167
340	Meningitis, except meningococcal and tuberculous	47	14	33
342	Intracranial and intraspinal abscess	2	2	0
343	Encephalitis, myelitis, and encephalomyelitis (except acute infectious)	8	5	3
344	Late effects of intracranial abscess or pyogenic infection	8	2	6
345	Multiple sclerosis	9	5	4
350	Paralysis agitans	28	24	4
351	Cerebral spastic infantile paralysis	7	4	3
352	Other cerebral paralysis	12	3	9
353	Epilepsy	24	8	16
355	Other diseases of brain	18	12	6
356	Motor neurone disease and muscular atrophy	10	8	2
357	Other diseases of spinal cord	6	4	2
361	Trigeminal neuralgia	1	1	0
364	Polyneuritis and polyradiculitis	1	0	1
368	Other diseases of peripheral nerves except autonomic	1	0	1
385	Cataract	2	1	1
393	Mastoiditis without mention of otitis media	1	1	0
<u>400-443</u>	<u>Diseases of the heart</u>	<u>7,300</u>	<u>4,640</u>	<u>2,660</u>
400	Rheumatic fever without mention of heart involvement	1	1	0
401	Rheumatic fever with heart involvement	6	2	4
410	Diseases of mitral valve	29	22	7
411	Diseases of aortic valve specified as rheumatic	10	7	3
414	Other endocarditis specified as rheumatic	5	3	2
415	Other myocarditis specified as rheumatic	2	0	2
416	Other heart disease specified as rheumatic	29	17	12
420	Arteriosclerotic heart disease, including coronary disease	5,034	3,647	1,387
421	Chronic endocarditis not specified as rheumatic	37	23	14
422	Other myocardial degeneration	548	313	235
430	Acute and subacute endocarditis	15	9	6
431	Acute myocarditis not specified as rheumatic	11	5	6
433	Functional disease of heart	112	71	41
434	Other and unspecified diseases of heart	451	217	234

Table 5 - Page 5  
 Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
441	Essential malignant hypertensive heart disease	15	1	14
442	Hypertensive heart disease with arteriolar nephrosclerosis	153	77	76
443	Other and unspecified hypertensive heart disease	842	225	617
<u>444-447</u>	<u>Other hypertensive disease</u>	<u>196</u>	<u>65</u>	<u>131</u>
444	Essential benign hypertension	59	20	39
445	Essential malignant hypertension	45	3	42
446	Hypertension with arteriolar nephrosclerosis	89	42	47
447	Other hypertensive disease without mention of heart	3	0	3
450	General arteriosclerosis	348	221	127
<u>451-468</u>	<u>Other diseases of circulatory system</u>	<u>288</u>	<u>191</u>	<u>97</u>
451	Aortic aneurysm, nonsyphilitic, and dissecting aneurysm	94	69	25
452	Other aneurysm, except of heart and aorta	13	8	5
453	Peripheral vascular disease	4	2	2
454	Arterial embolism and thrombosis	13	11	2
455	Gangrene of unspecified cause	5	1	4
456	Other diseases of arteries	19	15	4
460	Varicose veins of lower extremities	8	6	2
461	Hemorrhoids	1	0	1
462	Varicose veins of other specified sites	3	3	0
463	Phlebitis and thrombophlebitis of lower extremities	17	10	7
464	Phlebitis and thrombophlebitis of other sites	6	5	1
465	Pulmonary embolism and infarction	67	41	26
466	Other venous embolism and thrombosis	20	13	7
467	Other diseases of circulatory system	17	7	10
468	Certain diseases of lymph nodes and lymph channels	1	0	1
470	Acute nasopharyngitis (common cold)	3	1	2
471	Acute sinusitis	2	1	1
473	Acute tonsillitis	4	0	4
475	Acute upper respiratory infection of multiple or unspecified sites	7	1	6
<u>480-483</u>	<u>Influenza</u>	<u>22</u>	<u>7</u>	<u>15</u>
480	With pneumonia	11	3	8
481	With other respiratory manifestations, and influenza unqualified	10	4	6
482	With digestive manifestations, but without respiratory symptoms	1	0	1
<u>490-493</u>	<u>Pneumonia (excluding pneumonia of newborn)</u>	<u>635</u>	<u>303</u>	<u>332</u>
490	Lobar	106	38	68
491	Broncho	257	138	119
492	Primary atypical	60	22	38
493	Other and unspecified	212	105	107
<u>500-502</u>	<u>Bronchitis</u>	<u>50</u>	<u>43</u>	<u>7</u>
500	Acute	7	6	1
501	Unqualified	4	2	2
502	Chronic	39	35	4
510	Hypertrophy of tonsils and adenoids	1	1	0
511	Peritonsillar abscess (quinsy)	1	0	1

Table 5 - Page 6  
Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
519	Pleurisy	4	3	1
521	Abscess of lung	10	5	5
522	Pulmonary congestion and hypostasis	24	14	10
523	Pneumoconiosis due to silica and silicates (occupational)	1	0	1
<u>525-527</u>	<u>Other bronchopulmonic diseases</u>	<u>275</u>	<u>214</u>	<u>61</u>
525	Other chronic interstitial pneumonia	27	19	8
526	Bronchiectasis	21	17	4
527	Other diseases of lung and pleural cavity	227	178	49
538	Other diseases of buccal cavity	1	0	1
539	Diseases of esophagus	5	3	2
<u>540, 541</u>	<u>Ulcer of stomach and duodenum</u>	<u>88</u>	<u>58</u>	<u>30</u>
540	Stomach	41	25	16
541	Duodenum	47	33	14
542	Gastrojejunal ulcer	3	2	1
543	Gastritis and duodenitis	6	6	0
544	Disorders of function of stomach	3	2	1
545	Other diseases of stomach and duodenum	7	5	2
<u>550-553</u>	<u>Appendicitis</u>	<u>17</u>	<u>9</u>	<u>8</u>
550	Acute	17	9	8
<u>560, 561, 570</u>	<u>Hernia and intestinal obstruction</u>	<u>111</u>	<u>64</u>	<u>47</u>
560	Hernia of abdominal cavity without mention of obstruction	9	7	2
561	Hernia of abdominal cavity with obstruction	19	6	13
570	Intestinal obstruction without mention of hernia	83	51	32
571	Gastro-enteritis and colitis, except ulcerative, age 4 weeks and over	110	24	86
572	Chronic enteritis and ulcerative colitis	48	44	4
573	Functional disorders of intestines	1	1	0
575	Abscess of anal and rectal regions	1	0	1
576	Peritonitis	12	5	7
578	Other diseases of intestines and peritoneum	51	32	19
580	Acute and subacute yellow atrophy of liver	5	1	4
581	Cirrhosis of liver	132	90	42
583	Other diseases of liver	15	7	8
<u>584, 585</u>	<u>Cholelithiasis, cholecystitis, and cholangitis</u>	<u>42</u>	<u>28</u>	<u>14</u>
584	Cholelithiasis	17	12	5
585	Cholecystitis and cholangitis, without mention of calculi	25	16	9
586	Other diseases of gallbladder and biliary ducts	7	6	1
587	Diseases of pancreas	25	16	9
<u>590, 591</u>	<u>Acute nephritis and nephritis with edema, including nephrosis</u>	<u>25</u>	<u>7</u>	<u>18</u>
590	Acute	14	3	11
591	With edema, including nephrosis	11	4	7
<u>592-594</u>	<u>Chronic and unspecified nephritis and other renal sclerosis</u>	<u>185</u>	<u>57</u>	<u>128</u>
592	Chronic nephritis	136	42	94
593	Nephritis not specified as acute or chronic	48	14	34
594	Other renal sclerosis	1	1	0
600	Infections of kidney	156	93	63
601	Hydronephrosis	2	1	1

Table 5 - Page 7  
 Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
602	Calculi of kidney and ureter	8	7	1
603	Other diseases of kidney and ureter	16	6	10
604	Calculi of other parts of urinary system	2	2	0
605	Cystitis	2	1	1
606	Other diseases of bladder	3	1	2
608	Stricture of urethra	4	0	4
609	Other diseases of urethra	10	8	2
610	Hyperplasia of prostate	56	19	37
611	Prostatitis	4	4	0
612	Other diseases of prostate	1	0	1
613	Hydrocele	1	1	0
617	Other diseases of male genital organs	4	3	1
624	Salpingitis and oophoritis, unqualified	1	0	1
626	Diseases of parametrium and pelvic peritoneum (female)	4	2	2
630	Infective disease of uterus, vagina, and vulva	1	1	0
631	Uterovaginal prolapse	2	1	1
<u>640-689</u>	<u>Deliveries and complications of pregnancy, child-birth, and the puerperium</u>	<u>36</u>	<u>10</u>	<u>26</u>
640	Pyelitis and pyelonephritis of pregnancy	1	0	1
642	Toxemias of pregnancy	9	1	8
645	Ectopic pregnancy	1	0	1
646	Anemia of pregnancy	1	0	1
648	Other complications arising from pregnancy	4	0	4
650	Abortion without mention of sepsis or toxemia	1	0	1
651	Abortion with sepsis	2	0	2
660	Delivery without mention of complication	1	0	1
671	Delivery complicated by retained placenta	1	0	1
672	Delivery complicated by other postpartum hemorrhage	4	2	2
674	Delivery complicated by disproportion or mal-position of fetus	2	2	0
675	Delivery complicated by prolonged labor of other origin	1	0	1
677	Delivery with other trauma	1	0	1
678	Delivery with other complications of childbirth	2	1	1
681	Sepsis of childbirth and the puerperium	3	2	1
684	Puerperal pulmonary embolism	1	1	0
687	Cerebral hemorrhage in the puerperium	1	1	0
690	Boil and carbuncle	1	1	0
692	Other cellulitis and abscess without mention of lymphangitis	8	3	5
695	Impetigo	2	0	2
698	Other local infections of skin and subcutaneous tissue	2	1	1
701	Eczema	1	0	1
704	Pemphigus	2	1	1
705	Erythematous conditions	2	2	0
710	Other hypertrophic and atrophic conditions of skin	7	3	4
715	Chronic ulcer of skin	10	7	3
722	Rheumatoid arthritis and allied conditions	16	11	5
723	Osteo-arthritis (arthrosis) and allied conditions	5	3	2

Table 5 - Page 8  
Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
725	Arthritis, unspecified	7	6	1
726	Muscular rheumatism	1	0	1
730	Osteomyelitis and periostitis	4	4	0
733	Other diseases of bone	2	1	1
735	Displacement of intervertebral disc	1	1	0
744	Other diseases of muscle, tendon, and fascia	8	5	3
745	Curvature of spine	1	0	1
749	Other deformities	1	0	1
<u>750-759</u>	<u>Congenital malformations</u>	<u>227</u>	<u>123</u>	<u>104</u>
750	Monstrosity	7	5	2
751	Spina bifida and meningocele	16	13	3
752	Congenital hydrocephalus	16	7	9
753	Nervous system and sense organs, other	5	3	2
754	Circulatory system	111	61	50
755	Cleft palate and harelip	1	1	0
756	Digestive system	23	8	15
757	Genito-urinary system	15	11	4
759	Other and unspecified, not elsewhere classified	33	14	19
<u>760-776</u>	<u>Certain diseases of early infancy</u>	<u>875</u>	<u>308</u>	<u>567</u>
760	Intracranial and spinal injury at birth	25	12	13
761	Other birth injury	61	32	29
762	Postnatal asphyxia and atelectasis	138	67	71
763	Pneumonia of newborn	52	13	39
764	Diarrhea of newborn	15	0	15
767	Umbilical sepsis	1	0	1
768	Other sepsis of newborn	13	4	9
769	Neonatal disorders arising from certain diseases of the mother during pregnancy	10	2	8
770	Hemolytic disease of newborn (erythroblastosis)	10	6	4
771	Hemorrhagic disease of newborn	8	2	6
772	Nutritional maladjustment	22	2	20
773	Ill-defined diseases peculiar to early infancy	193	87	106
774	Immaturity with mention of any other subsidiary condition	24	7	17
776	Immaturity, unqualified	303	74	229
<u>780-795</u>	<u>Symptoms, senility, and ill-defined conditions</u>	<u>1,601</u>	<u>436</u>	<u>1,165</u>
780	Certain symptoms referable to nervous system and special senses	5	1	4
781	Other symptoms referable to nervous system and special senses	1	1	0
782	Symptoms referable to cardiovascular and lymphatic system	36	17	19
783	Symptoms referable to respiratory system	1	0	1
784	Symptoms referable to upper gastro-intestinal tract	2	1	1
785	Symptoms referable to abdomen and lower gastro-intestinal tract	4	2	2
786	Symptoms referable to genito-urinary system	1	1	0
787	Symptoms referable to limbs and back	1	0	1
788	Other general symptoms	4	2	2
792	Uremia .	18	9	9

Table 5 - Page 9  
 Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
794	Senility without mention of psychosis	54	4	50
795	Ill-defined and unknown causes	1,474	398	1,076
<u>800-962</u>		<u>1,890</u>	<u>1,166</u>	<u>724</u>
<u>800-802</u>	<u>Accidents</u>	<u>14</u>	<u>7</u>	<u>7</u>
800	Railway	3	2	1
802	Involving railroad employee	11	5	6
<u>810-835</u>	<u>Motor vehicle</u>	<u>931</u>	<u>649</u>	<u>282</u>
810	Traffic, collision with railway train	30	21	9
812	Traffic, to pedestrian	115	53	62
813	Traffic, to pedal cyclist	9	4	5
815	Traffic, to rider or passenger of motorcycle in collision with other motor vehicle	17	13	4
816	Traffic, other involving two or more motor vehicles	387	297	90
818	Traffic, collision with animal or animal-drawn vehicle	2	1	1
819	Traffic, collision with fixed or unspecified object	51	37	14
821	Traffic, to rider of motorcycle without antecedent collision	1	1	0
822	Traffic, overturning in roadway	22	15	7
823	Traffic, running off roadway	195	141	54
824	Traffic, other noncollision	12	5	7
825	Traffic, unspecified nature	69	50	19
830	Nontraffic, to pedestrian	14	7	7
834	Nontraffic while boarding and alighting	1	1	0
835	Nontraffic, other and unspecified nature	6	3	3
<u>840-845</u>	<u>Other road vehicle</u>	<u>5</u>	<u>3</u>	<u>2</u>
845	Other non-motor road vehicle accidents	5	3	2
<u>850-858</u>	<u>Water transport</u>	<u>36</u>	<u>21</u>	<u>15</u>
850	Submersion of occupant of small boat	33	19	14
851	Other injury by submersion	3	2	1
<u>860-866</u>	<u>Aircraft</u>	<u>23</u>	<u>19</u>	<u>4</u>
860	To personnel in military aircraft	7	6	1
861	To occupant by accident to commercial "transport" aircraft	8	8	0
863	To occupant of other specified aircraft	5	3	2
865	Other than at airfield to person not in aircraft	1	0	1
866	Other and unspecified	2	2	0
<u>870-895</u>	<u>Poisoning</u>	<u>27</u>	<u>17</u>	<u>10</u>
871	Barbituric acid and derivatives	2	2	0
872	Aspirin and salicylates	1	1	0
874	Other and analgesic and soporific drugs	1	1	0
878	Other and unspecified drugs	2	2	0
880	Alcohol	3	1	2
882	Industrial solvents	1	1	0
883	Corrosive aromatics, acids, and caustic alkalies	1	0	1
884	Mercury and its compounds	1	1	0
886	Arsenic and antimony, and their compounds	3	2	1
888	Other and unspecified solid and liquid substances	5	1	4
890	Utility (illuminating) gas	3	2	1

Table 5 - Page 10  
Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
891	Motor vehicle exhaust gas	2	1	1
894	Other specified gases and vapours	2	2	0
<u>900-904</u>	<u>Falls</u>	<u>178</u>	<u>138</u>	<u>40</u>
900	On stairs	4	3	1
901	From ladders	1	1	0
902	Other, from one level to another	36	22	14
903	On same level	31	29	2
904	Unspecified	106	83	23
910	Blow from falling or projected object or missile	32	16	16
912	Caused by machinery	47	32	15
913	Caused by cutting and piercing instruments	3	1	2
914	Caused by electric current	21	17	4
915	Caused by explosion of pressure vessel	1	1	0
916	Caused by fire and explosion of combustible material	193	73	120
917	Caused by hot substance, corrosive liquid, and steam	6	2	4
919	Caused by firearm	104	51	53
921	Inhalation and ingestion of food causing obstruction or suffocation	31	19	12
922	Inhalation and ingestion of other object causing obstruction or suffocation	4	1	3
923	Foreign body entering other orifice	24	13	11
924	Mechanical suffocation in bed and cradle	20	7	13
925	Mechanical suffocation in other and unspecified circumstances	11	2	9
926	Lack of care of infants under 1 year of age	1	0	1
927	Caused by bites and stings of venomous animals and insects	2	0	2
929	Drowning and submersion	98	39	59
931	Excessive heat and insolation	2	1	1
932	Excessive cold	3	1	2
933	Hunger, thirst, and exposure	9	1	8
934	Cataclysm	7	2	5
935	Lightning	4	1	3
936	Other and unspecified accidents	32	18	14
943	Post-immunization jaundice and hepatitis	1	1	0
946	Other complications due to non-therapeutic medical and surgical procedures	1	0	1
950	Therapeutic misadventure in surgical treatment	3	1	2
951	Therapeutic misadventure in infusion or transfusion	1	1	0
953	Therapeutic misadventure in administration of drugs or biologicals	1	1	0
954	Therapeutic misadventure in anesthesia	1	1	0
955	Other and unspecified therapeutic misadventure	1	1	0
960	Late effect of motor vehicle accident	2	2	0
962	Late effect of other accidental injury	10	7	3
<u>963, 970-979</u>	<u>Suicide</u>	<u>164</u>	<u>149</u>	<u>15</u>
970	By poisoning with analgesic and soporific substances	7	7	0

Table 5 - Page 11  
 Deaths - Mississippi - 1967

International List Number*	Cause of Death**	Total	White	Non-white
971	By poisoning with other solid and liquid substances	5	4	1
972	By gases in domestic use	1	1	0
973	By other gases	5	5	0
974	By hanging and strangulation	9	8	1
975	By submersion (drowning)	4	3	1
976	By firearms and explosives	129	117	12
977	By cutting and piercing instruments	2	2	0
979	By other and unspecified means	2	2	0
<u>964, 980-985</u>	<u>Homicide</u>	<u>271</u>	<u>66</u>	<u>205</u>
981	Assault by firearm and explosive	205	56	149
982	Assault by cutting and piercing instruments	48	6	42
983	Assault by other means	14	3	11
984	Injury by intervention of police	4	1	3

\* International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

\*\* When cause of death and International List Number are omitted, no deaths were reported.

TABLE 6  
DEATHS  
MISSISSIPPI  
1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number		Non-white	Rate*		Non-white
	Total	White		Total	White		Total	White	
STATE	22,121	12,538	9,583	22,609	12,816	9,793	9.6	9.0	10.5
Adams	496	226	270	401	180	221	10.3	8.5	12.4
Alcorn	242	203	39	255	210	45	9.5	8.9	14.2
Amite	77	35	42	137	69	68	9.2	10.4	8.2
Attala	228	144	84	244	154	90	10.0	10.4	9.5
Benton	26	17	9	63	42	21	8.2	9.3	6.7
Bolivar	551	124	427	560	165	395	9.7	8.0	10.7
Calhoun	158	121	37	198	157	41	12.7	13.6	10.2
Carroll	49	20	29	108	58	50	10.0	10.9	9.2
Chickasaw	233	140	93	217	134	83	11.5	11.6	11.5
Choctaw	77	50	27	106	80	26	12.3	12.6	11.4
Claiborne	102	23	79	141	41	100	14.0	13.4	14.3
Clarke	111	61	50	182	116	66	11.8	12.2	11.3
Clay	164	86	78	176	87	89	9.9	9.9	10.0
Coahoma	551	192	359	518	151	367	10.9	10.1	11.2
Copiah	243	131	112	326	188	138	11.9	13.4	10.3
Covington	82	58	24	130	92	38	8.9	9.4	7.7
DeSoto	144	53	91	261	118	143	7.1	5.1	10.2
Forrest	610	427	183	595	370	225	9.5	7.9	13.9
Franklin	116	65	51	140	85	55	15.1	14.7	15.7
George	90	75	15	110	94	16	9.0	8.7	11.3
Greene	65	50	15	83	65	18	9.9	9.9	10.0
Grenada	237	121	116	224	101	123	10.6	8.9	12.5
Hancock	162	135	27	180	151	29	8.7	8.5	10.1
Harrison	1,226	982	244	1,010	795	215	7.2	6.6	10.4
Hinds	2,696	1,524	1,172	1,722	944	778	7.7	7.5	8.0
Holmes	270	64	206	337	100	237	11.8	11.9	11.8
Humphreys	156	40	116	190	58	132	10.0	10.4	9.8
Issaquena	20	4	16	42	14	28	15.1	14.9	15.2
Itawamba	77	68	9	146	137	9	8.6	8.6	9.0
Jackson	378	293	85	454	355	99	6.0	5.6	8.1
Jasper	165	86	79	202	99	103	12.5	11.9	13.2
Jefferson	82	15	67	131	37	94	11.4	12.8	10.9
Jeff Davis	135	69	66	143	72	71	10.3	11.2	9.5
Jones	716	466	250	601	439	162	10.1	9.9	10.5
Kemper	78	35	43	114	58	56	9.9	10.9	9.0
Lafayette	198	139	59	196	129	67	7.9	7.5	8.8

Table 6 - Page 2  
 Deaths - Mississippi - 1967

County	Place of Occurrence			Place of Residence					
	Number		Non-white	Number			Rate*		
	Total	White		Total	White	Non-white	Total	White	Non-white
Lamar	77	54	23	136	109	27	9.0	8.5	11.9
Lauderdale	986	627	359	770	459	311	10.0	8.8	12.7
Lawrence	89	56	33	142	99	43	12.5	12.6	12.1
Leake	162	109	53	197	135	62	9.6	10.2	8.4
Lee	496	382	114	452	339	113	9.2	8.8	10.5
Leflore	493	197	296	452	160	292	9.1	8.4	9.5
Lincoln	297	200	97	312	211	101	11.9	11.8	12.1
Lowndes	417	202	215	467	242	225	8.4	6.8	11.2
Madison	300	103	197	334	102	232	10.4	9.9	10.7
Marion	225	149	76	248	155	93	10.5	9.9	11.8
Marshall	188	70	118	263	101	162	10.7	12.0	10.0
Monroe	330	230	100	357	239	118	10.9	10.9	10.9
Montgomery	154	90	64	142	86	56	9.5	10.1	8.8
Neshoba	205	144	61	243	181	62	12.1	12.2	11.8
Newton	192	138	54	230	161	69	11.4	11.2	11.8
Noxubee	140	42	98	168	57	111	10.9	12.6	10.2
Oktibbeha	208	81	127	226	97	129	7.4	5.2	10.7
Panola	249	97	152	320	132	188	10.4	10.0	10.7
Pearl River	212	168	44	246	191	55	8.5	7.9	11.4
Perry	78	54	24	94	68	26	10.2	10.5	9.3
Pike	350	218	132	362	216	146	11.2	11.7	10.5
Pontotoc	181	149	32	215	176	39	11.8	11.7	12.1
Prentiss	195	172	23	198	177	21	9.7	9.8	8.8
Quitman	136	34	102	203	72	131	10.2	9.3	10.9
Rankin	360	210	150	247	158	89	6.3	6.2	6.4
Scott	203	132	71	241	150	91	10.2	9.9	10.8
Sharkey	91	26	65	111	32	79	10.9	9.9	11.4
Simpson	230	161	69	236	172	64	11.8	12.6	10.1
Smith	54	40	14	133	103	30	9.9	10.7	8.0
Stone	66	53	13	77	61	16	9.6	9.9	8.5
Sunflower	369	120	249	416	142	274	9.6	10.4	9.2
Tallahatchie	208	69	139	264	107	157	11.5	13.4	10.5
Tate	86	24	62	155	73	82	7.8	7.9	7.7
Tippah	136	99	37	147	114	33	8.3	7.8	10.5
Tishomingo	114	109	5	153	144	9	8.5	8.2	15.5
Tunica	135	36	99	158	37	121	10.1	9.2	10.4
Union	191	159	32	201	170	31	10.4	10.3	11.3
Walton	123	67	56	144	84	60	10.1	10.9	9.2
Warren	650	277	373	532	220	312	11.7	8.4	16.1
Washington	770	292	478	799	276	523	10.2	7.8	12.3
Wayne	156	92	64	185	113	72	11.2	10.8	11.8
Webster	112	95	17	123	104	19	11.7	13.0	7.5
Wilkinson	109	50	59	135	54	81	11.7	15.7	10.0
Winston	156	110	46	187	126	61	10.9	13.3	8.0
Yalobusha	123	76	47	164	107	57	13.4	16.0	10.3
Yazoo	308	133	175	381	159	222	11.1	11.1	11.1
Out-of-State	1,244	866	378	756	588	168	-	-	-

Table 6 - Page 3  
 Deaths - Mississippi - 1967

City**	Place of Occurrence			Place of Residence		
	Number		Non-white	Number		Non-white
	Total	White		Total	White	
Biloxi	434	370	64	373	310	63
Clarksdale	424	184	240	273	105	168
Cleveland	169	85	84	104	52	52
Columbus	308	167	141	267	145	122
Corinth	59	43	16	121	92	29
Greenville	545	243	302	457	189	268
Greenwood	352	152	200	256	111	145
Gulfport	491	374	117	325	242	83
Hattiesburg	521	373	148	399	231	168
Jackson	2,237	1,249	988	1,326	774	552
Laurel	600	374	226	313	185	128
McComb	211	136	75	146	96	50
Meridian	912	586	326	587	342	245
Natchez	464	214	250	304	135	169
Pascagoula	291	229	62	159	127	32
Tupelo	384	301	83	184	132	52
Vicksburg	581	252	329	381	148	233
Yazoo City	178	99	79	191	73	118
Total Urban#	13,179	7,850	5,329	9,053	5,134	3,919
Total Rural	8,942	4,688	4,254	13,556	7,682	5,874

\* Deaths per 1,000 population.

\*\* Cities having 10,000 population and over according to 1960 census.

# Includes all incorporated towns of 2,500 population or more according to 1960 census.

TABLE 7  
DEATHS  
MISSISSIPPI  
1967

**By Cause Groups and Certain Selected Causes, by Race, Sex, and Age,  
With Total Rates. Residence Data**

Table 7 - Page 2  
Deaths - Mississippi - 1967

International List Number#	Cause of Death	Total		Race by Sex	Total by Race and Sex	Age	Unknown					
		Number	Rate*									
045-048	Dysentery (all forms)	7	0.3	M	1	M	0	0	0	0	0	0
				C	6	F	1	0	0	0	0	0
050,051	Scarlet fever and streptococcal sore throat	2	0.1	W	0	M	0	0	0	0	0	0
				C	2	F	0	0	0	0	0	0
055	Diphtheria	0	-	W	0	M	0	0	0	0	0	0
				C	0	F	0	0	0	0	0	0
056	Whooping cough	3	0.1	W	1	M	0	1	1	0	0	0
				C	2	F	1	1	1	0	0	0
057	Meningococcal infections	6	0.2	W	3	M	0	0	2	1	0	0
				C	3	F	3	0	2	0	0	0
080	Acute poliomyelitis	1	**	W	0	M	0	0	0	1	0	0
				C	1	F	1	0	0	1	0	0
085	Measles	3	0.1	W	1	M	0	1	1	0	0	0
				C	2	F	1	1	0	0	0	0

Table 7 - Page 3  
Deaths - Mississippi - 1967

30

International List Number#	Cause of Death	Total	Rate*	Race	Total by Race and Sex	Age	Unknown												
							Number	Rate*	Total by Race	Sex	Under 1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65 Over 64
140-205	Malignant neoplasms	3,136	132.8	W	2,061	M	1,167	2	5	9	54	133	290	649	0	0	0	0	0
				C	1,075	F	894		1	4	47	117	168	527	0	0	0	0	0
210-239	Benign and unspecified neoplasms	39	1.6	W	20	M	14		0	1	0	10	1	5	4	0	0	0	0
				C	19	F	6		1	0	1	10	1	0	3	3	3	1	
240-289	Allergic, endocrine system, metabolic, & nutritional diseases	449	19.0	W	213	M	92	2	1	2	1	3	11	13	57	0	0	0	0
				C	236	F	121		0	1	0	1	0	10	17	82	0	0	0
241	Asthma	50	2.1	W	29	M	16		0	1	0	0	0	1	1	1	3	5	0
				C	21	F	13		1	0	0	0	0	0	0	11	33	0	
260	Diabetes mellitus	332	14.1	W	146	M	61	0	0	0	0	0	0	8	9	42	0	0	
				C	186	F	85		1	0	0	0	0	1	1	7	11	33	0
290-299	Diseases of the blood and blood-forming organs	65	2.8	W	26	M	13		0	0	1	4	1	0	3	0	7	4	0
				C	39	F	13		1	0	2	3	1	0	1	1	1	1	0
300-326	Mental, psycho-neurotic, & personality disorders	50	2.1	W	38	M	25		0	0	0	0	0	2	0	2	2	6	0
				C	12	F	8		1	0	1	0	1	0	1	1	1	1	0

Table 7 - Page 4  
Deaths - Mississippi - 1967

International List Number#	Cause of Death	Total		Total by Race	Total by Sex	Total by Race	Total by Sex	Age						Unknown		
		Total Number	Rate*					1-4	5-9	10-14	15-24	25-34	35-44	45-54		
330-334	Vascular lesions affecting central nervous system	2,981	126.2	W 1,632	M 806	F 826	M 674	0	0	0	0	3	12	40	82	669
				C 1,349	M 675	F 675	C 1,011	1	1	1	1	1	11	28	57	720
340-398	Other diseases of the nervous system and sense organs	185	7.8	W 94	M 57	F 37	M 7	1	1	1	1	1	5	18	45	484
				C 91	M 50	F 41	C 11	1	1	1	1	1	2	3	53	491
340	Nonmeningococcal meningitis	47	2.0	W 14	M 9	F 5	M 2	1	0	1	0	1	0	1	0	21
				C 33	M 18	F 15	C 15	6	5	6	5	1	0	1	0	17
400-468	Diseases of the circulatory system	8,132	344.4	W 5,117	M 3,079	F 2,038	M 1,597	2	0	0	0	0	5	15	44	628
				C 3,015	M 1,597	F 1,418	C 11	3	3	2	1	1	4	19	17	2,043
400-402	Rheumatic fever	7	0.3	W 3	M 1	F 2	M 0	0	0	0	0	0	0	0	0	80
				C 4	M 2	F 2	C 0	0	0	0	0	0	1	0	163	219
410-416	Chronic rheumatic heart disease	75	3.2	W 49	M 20	F 29	M 13	0	0	0	0	0	1	1	1	163
				C 26	M 13	F 13	C 0	0	0	0	1	1	1	1	108	305
420-422	Arteriosclerotic and degenerative heart disease	5,619	238.0	W 3,983	M 2,491	F 1,492	M 885	1	0	0	0	0	3	2	68	543
				C 1,636	M 751	F 751	C 1	0	1	0	1	0	1	0	1	1,605

Table 7 - Page 5  
Deaths - Mississippi - 1967

Inter-national List Number#	Cause of Death	Total		Race*	Total by Race	Sex	Age						Unknown Over 65				
		Total Number	Rate*				1-4	5-9	10-14	15-24	25-34	35-44	45-54				
430-434	Other diseases of the heart	589	24.9	W	302	M	182	0	0	1	1	9	17	32	121	0	
				C	287	F	120	0	1	0	0	3	10	15	91	0	
440-443	Hypertensive heart disease	1,010	42.8	W	303	M	130	0	0	1	1	1	4	16	108	0	
				F	169	M	118	0	1	1	4	2	8	18	42	88	
					3		3	0	0	0	0	3	8	12	27	63	
444-447	Other hypertensive disease	196	8.3	W	65	M	34	0	0	0	0	1	1	2	3	27	0
				C	131	F	31	0	0	0	0	0	0	0	2	26	0
450	General arteriosclerosis	348	14.7	W	221	M	100	0	0	0	0	0	0	0	3	118	0
				C	127	F	121	0	0	0	0	0	0	0	0	60	0
					66		69	0	0	0	0	0	0	0	5	53	0
470-527	Diseases of the respiratory system	1,039	44.0	W	593	M	384	15	8	3	2	2	7	19	59	267	0
				C	446	F	209	3	6	1	1	2	2	5	10	22	157
					243		203	61	22	0	0	2	5	16	32	100	0
					77		77	0	0	0	0	0	0	10	8	75	0
480-483	Influenza	22	0.9	W	7	M	5	0	0	0	0	0	0	0	0	4	0
				C	15	M	2	0	0	0	0	0	0	0	0	2	0
					3		1	0	0	0	0	0	0	0	0	12	0
490-493	Pneumonia (excluding pneumonia of newborn)	635	26.9	W	303	M	166	12	7	2	2	1	0	4	8	115	0
				F	137		180	2	4	1	0	3	2	0	3	111	0
					180		152	52	20	4	1	3	1	1	6	21	0
					64		64	17	0	2	1	4	4	5	7	48	0

Table 7 - Page 6  
Deaths - Mississippi - 1967

Inter-national List Number#	Cause of Death	Total		Race	Total by Race	Sex	Total by Race	Age	Unknown											
		Number	Rate*						1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65+ Over 65			
500-502	Bronchitis	50	2.1	W	43	M	33	2	0	0	0	0	0	1	3	7	20	0		
				C	7	F	10	0	1	0	0	0	0	0	2	1	6	0		
				F	3		4	1	0	0	0	0	0	0	0	1	2	0		
525-527	Other broncho-pulmonic diseases	275	11.6	W	214	M	166	1	0	0	0	0	0	1	2	7	36	118	0	
				C	61	F	48	0	1	0	0	0	0	0	0	5	9	31	0	
				F	18		43	5	1	0	0	0	0	0	0	7	7	23	0	
					61		6	1	0	0	0	0	0	0	0	3	1	6	0	
530-587	<u>Diseases of the digestive system</u>	690	29.2	W	403	M	244	5	1	0	3	3	4	4	16	38	48	126	0	
				C	287	F	159	13	6	1	2	1	2	1	16	12	17	91	0	
				F	145		47	47	6	4	5	5	5	5	5	9	17	46	0	
					142		32	32	4	4	4	4	4	4	4	9	17	45	0	
540,541	Ulcer of stomach and duodenum	88	3.7	W	58	M	43	0	0	0	0	0	0	0	0	0	7	12	22	0
				C	30	F	15	0	0	0	0	0	0	0	0	0	2	0	9	0
				F	19		11	1	0	0	0	0	0	0	0	3	4	8	0	
					11		0	0	0	0	0	0	0	0	0	3	0	5	0	
560,561, 570	Hernia and intestinal obstruction	111	4.7	W	64	M	36	4	0	0	0	0	0	0	1	1	3	5	22	0
				C	47	F	28	3	0	0	0	0	0	0	0	0	1	1	13	0
				F	20		20	2	0	0	0	0	0	0	0	0	1	7	16	0
					27		3	0	0	0	0	0	0	0	0	0	1	7	16	0
543,572	Gastritis, duodenitis, enteritis, & colitis, except diarrhea of the newborn	164	6.9	W	74	M	30	1	1	0	0	0	0	2	1	1	3	4	17	0
				C	90	F	44	7	1	0	0	0	0	1	0	1	2	5	27	0
				F	54		42	5	0	0	0	0	0	0	0	0	1	0	5	0
					36		26	4	0	0	0	0	0	0	0	0	0	1	3	0
581	Cirrhosis of liver	132	5.6	W	90	M	64	0	0	0	0	0	0	0	1	0	8	20	16	0
				C	42	F	26	0	0	0	0	0	0	0	0	0	1	5	10	5
				F	20		22	0	0	0	0	0	0	0	0	0	2	5	6	0
					42		20	0	0	0	0	0	0	0	0	0	3	6	2	0

Table 7 - Page 7  
Deaths - Mississippi - 1967

Table 7 - Page 8  
Deaths - Mississippi - 1967

Inter-national List Number/ <sup>#</sup>	Cause of Death	Total		Race*	Total by Race	Total by Race	Total by Race and Sex	Age						Unknown Over & 65	Unknown		
		Number	Rate*					1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64		
760-776	Certain diseases of early infancy	875	18.8	W	308	M	181	181	127	0	0	0	0	0	0	0	
				C	567	M	327	327	240	0	0	0	0	0	0	0	
780-795	Symptoms, senility, and ill-defined conditions	1,601	67.8	W	436	M	313	16	1	0	3	4	12	72	154	0	
				C	1,165	M	123	10	15	0	1	11	6	22	67	0	
				F	665	M	665	92	82	12	5	4	26	99	342	1	
					500	F	500					13	23	70	249	3	
800-999	Accidents, poison- ings, and violence	2,325	98.5	W	1,381	M	986	16	16	24	31	198	132	147	162	0	
				C	944	M	395	9	15	16	14	76	31	46	112	0	
				F	708	M	708	35	36	16	14	156	45	31	77	0	
					236	F	236	19	33	22	11	17	24	21	51	0	
810-835	Motor vehicle accidents	931	39.4	W	649	M	452	6	8	12	14	131	69	58	63	44	0
				C	282	M	197	1	8	11	8	63	17	22	21	47	0
				F	216	M	216	2	8	13	9	58	24	35	21	28	0
					66	F	66	1	4	6	5	8	6	7	8	11	0
800-802, 840-962	All other accidents	959	40.6	W	517	M	367	10	8	12	15	54	43	39	48	90	0
				C	442	M	150	8	7	5	5	59	4	14	13	76	0
				F	312	M	312	33	28	19	24	37	31	20	18	43	0
					130	F	130	18	29	14	5	3	7	5	8	36	0
963,	Suicide	164	6.9	W	149	M	112	0	0	0	0	1	9	5	21	25	0
970-979				C	15	M	37	0	0	0	0	1	1	3	7	4	0
				F	4	F	11	0	0	0	0	1	0	1	2	2	0
														0	1	1	0

Table 7 - Page 9  
Deaths - Mississippi - 1967

Inter-national List Number#	Cause of Death	Total		Total by Race		Total by Race and Sex		Age									
		Number	Rate*	Total by Race	Race	Under 14	14-17	18-24	25-34	35-44	45-54	55-64	65 over & Unknown				
964, 980-985	Homicide	271	11.5	W	66	M 55	0	0	1	4	15	18	11	4	2	0	
				C	205	F 11	0	0	0	1	3	2	2	0	1	0	
					36	M 169	0	0	0	1	38	54	37	25	10	4	0
						F 36	0	0	2	1	5	11	9	4	1	3	0

# International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

\* Total deaths per 1,000 population. Deaths from specific causes per 100,000 population. Deaths from maternal causes per 10,000 live births. Deaths from diseases of early infancy per 1,000 live births.

\*\* Rate less than 0.06.

TABLE 8  
DEATHS UNDER 1 YEAR  
MISSISSIPPI  
1967

Table 8 - Page 2  
Deaths Under 1 Year - Mississippi - 1967

Table 8 - Page 3  
Deaths Under 1 Year - Mississippi - 1967

Cause of Death and International List Number#	Total Number	Total Rate*	Total by Race	Days of Age											Months of Age								
				Under 1 Day	1-	6	7-	13	14-	20	21-	27	Under 28 Days	1	2	3	4	5	6	7	8	9	10
Congenital malfor- mations (750-759)	164	3.5	W C	88 76	22 16	23 20	7 4	2 1	0 1	0 0	44 41	0 0	0 0	0 0	1 1	2 1	0 1	1 1	0 0	0 0	0 0	0 0	0 0
Birth injuries (760, 761)	86	1.8	W C	44 42	31 31	12 9	1 0	0 1	0 0	0 0	44 41	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Postnatal asphyxia and atelectasis (762)	138	3.0	W C	67 71	41 44	25 21	1 1	0 2	0 0	0 0	67 68	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Pneumonia of newborn (763)	52	1.1	W C	13 39	1 1	8 14	0 12	0 3	2 9	2 3	13 39	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Diarrhea of newborn (764)	15	0.3	W C	0 15	0 0	0 1	0 7	0 3	0 4	0 15	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Other infections of newborn (765-768)	14	0.3	W C	4 10	1 2	1 6	1 1	1 1	0 0	0 0	4 10	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Neonatal disorders arising from certain diseases of mother dur- ing pregnancy (769)	10	0.2	W C	2 8	2 5	0 3	0 0	0 0	0 0	0 0	2 8	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Hemolytic disease of newborn (erythroblas- tosis) (770)	10	0.2	W C	6 4	3 0	1 1	1 0	1 0	0 0	0 0	6 4	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Hemorrhagic disease of newborn (771)	8	0.2	W C	2 6	0 4	0 2	0 0	0 0	1 0	0 0	2 6	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Nutritional maladjust- ment (772)	22	0.5	W C	2 20	0 0	0 0	0 5	0 1	0 3	1 2	1 0	0 1	0 4	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0

Table 8 - Page 4  
Deaths Under 1 Year - Mississippi - 1967

Cause of Death and International List Number#	Total Number	Rate*	Total by Race	Total by Race	Days of Age										Months of Age					
					Under 1 Day	1- 6	7- 13	14- 20	21- 27	Under 28 Days	1	2	3	4	5	6	7	8	9	10
Ill-defined diseases peculiar to early infancy (773)	193	4.1	W C	87 106	43 55	42 45	1 3	1 2	0 0	87 105	0 0	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Immaturity with mention of any other subsidiary condition (774)	24	0.5	W C	7 17	5 4	2 6	0 0	0 0	0 1	7 11	0 3	0 1	0 1	0 0	0 0	0 0	0 0	0 0	0 0	0 0
Immaturity, unqualified (776)	303	6.5	W C	74 229	59 134	13 56	2 12	0 6	0 4	74 212	0 7	0 3	0 3	0 1	0 0	0 0	0 0	0 0	0 0	0 0
Symptoms & ill-defined conditions (780-793, 795)	200	4.3	W C	26 174	2 20	2 22	1 5	1 5	1 5	7 57	3 29	6 15	2 15	3 14	2 13	1 8	2 5	0 5	0 2	0 2
All other diseases	19	0.4	W C	7 12	1 0	2 0	1 0	0 3	0 0	4 4	1 2	0 1	0 3	0 0	0 0	0 0	0 1	0 0	0 0	0 1
Accidents (800-962)	79	1.7	W C	25 54	0 3	2 7	0 1	0 1	0 1	2 13	4 14	5 6	3 5	0 7	2 1	1 1	3 1	0 1	1 0	2 2
Homicide (964, 980-984)	0	-	W C	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0	0 0

# International List Numbers are those of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Seventh Revision, 1955.

\* Deaths per 1,000 live births.

\*\* Rate less than 0.06.

TABLE 9  
MARRIAGES AND DIVORCES  
MISSISSIPPI  
1967

Marriages by Place of Licensure, and by Place of Residence of Bride, by Race of Bride  
Divorces by Place of Occurrence, by Race

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride			Total	White	Non-white
	Total	White	Non-white	Total	White	Non-white			
<u>STATE</u>	<u>23,535</u>	<u>15,752</u>	<u>7,783</u>	<u>20,013</u>	<u>12,822</u>	<u>7,191</u>	<u>6,455</u>	<u>4,704</u>	<u>1,751</u>
Adams	402	262	140	310	187	123	128	84	44
Alcorn	496	453	43	247	222	25	79	72	7
Amite	125	71	54	105	64	41	20	12	8
Attala	176	99	77	191	109	82	43	32	11
Benton	55	31	24	63	40	23	11	10	1
Bolivar	527	238	289	463	196	267	115	66	49
Calhoun	195	147	48	140	91	49	35	30	5
Carroll	39	18	21	46	19	27	12	6	6
Chickasaw	234	156	78	213	144	69	55	35	20
Choctaw	91	59	32	56	35	21	17	16	1
Claiborne	80	28	52	67	16	51	16	3	13
Clarke	144	92	52	127	76	51	38	25	13
Clay	155	84	71	134	74	60	45	23	22
Coahoma	400	186	214	337	144	193	77	40	37
Copiah	214	105	109	228	113	115	68	32	36
Covington	139	99	40	153	115	38	26	21	5
DeSoto	902	553	349	209	112	97	104	80	24
Forrest	628	481	147	592	434	158	265	192	73
Franklin	77	52	25	65	47	18	19	8	11
George	216	202	14	109	101	8	32	30	2
Greene	75	68	7	72	52	20	18	16	2
Grenada	183	112	71	196	127	69	44	28	16
Hancock	399	359	40	138	114	24	79	74	5
Harrison	1,691	1,453	238	1,417	1,188	229	651	566	85
Hinds	2,131	1,341	790	2,112	1,344	768	672	498	174
Holmes	190	68	122	207	69	138	53	20	33
Humphreys	130	57	73	133	54	79	29	19	10
Issaquena	25	13	12	26	9	17	10	5	5
Itawamba	148	136	12	114	103	11	41	41	0
Jackson	880	701	179	710	557	153	270	231	39
Jasper	151	84	67	164	93	71	21	11	10
Jefferson	75	30	45	71	21	50	16	6	10
Jeff Davis	144	67	77	89	44	45	16	8	8
Jones	608	466	142	572	446	126	171	133	38
Kemper	88	28	60	66	18	48	22	10	12
Lafayette	155	104	51	123	72	51	68	37	31

Table 9 - Page 2  
 Marriages and Divorces - Mississippi - 1967

County	Marriages						Divorces		
	Place of Licensure			Residence of Bride					
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white
Lamar	115	101	14	133	117	16	47	41	6
Lauderdale	569	392	177	585	393	192	233	184	49
Lawrence	84	58	26	125	77	48	22	14	8
Leake	160	91	69	177	102	75	65	47	18
Lee	447	343	104	525	399	126	140	129	11
Leflore	426	194	232	418	196	222	88	65	23
Lincoln	251	184	67	228	162	66	43	29	14
Lowndes	397	279	118	363	247	116	133	96	37
Madison	306	113	193	267	100	167	87	20	67
Marion	265	197	68	245	186	59	96	75	21
Marshall	340	161	179	203	78	125	95	39	56
Monroe	291	190	101	264	166	98	84	70	14
Montgomery	129	72	57	125	67	58	30	19	11
Neshoba	193	137	56	161	106	55	93	78	15
Newton	151	104	47	184	130	54	34	18	16
Noxubee	111	42	69	114	35	79	19	7	12
Oktibbeha	227	137	90	221	130	91	39	22	17
Panola	271	160	111	217	115	102	63	39	24
Pearl River	350	304	46	256	212	44	151	137	14
Perry	107	90	17	102	90	12	22	17	5
Pike	352	224	128	336	200	136	100	71	29
Pontotoc	158	139	19	135	126	9	34	27	7
Prentiss	218	200	18	176	167	9	68	67	1
Quitman	171	87	84	174	81	93	30	14	16
Rankin	287	207	80	268	184	84	63	48	15
Scott	191	96	95	207	112	95	37	25	12
Sharkey	94	34	60	88	37	51	21	9	12
Simpson	216	160	56	181	129	52	80	65	15
Smith	99	82	17	93	66	27	31	25	6
Stone	113	94	19	83	67	16	22	16	6
Sunflower	273	101	172	338	142	196	63	39	24
Tallahatchie	190	88	102	152	62	90	32	20	12
Tate	240	136	104	156	83	73	28	22	6
Tippah	192	172	20	186	162	24	40	35	5
Tishomingo	151	148	3	122	120	2	28	27	1
Tunica	124	50	74	97	30	67	25	11	14
Union	187	165	22	159	138	21	37	30	7
Walthall	133	79	54	96	47	49	26	14	12
Warren	467	285	182	407	241	166	202	149	53
Washington	826	485	341	698	370	328	274	178	96
Wayne	199	125	74	148	93	55	44	32	12
Webster	81	68	13	86	73	13	14	13	1
Wilkinson	449	371	78	103	36	67	19	3	16
Winston	176	99	77	162	85	77	46	34	12
Yalobusha	125	73	52	139	86	53	29	23	6
Yazoo	265	132	133	245	127	118	92	41	51
Out-of-State				3,522	2,930	592			

TABLE 10  
DIVORCES  
MISSISSIPPI  
1967

By Cause of Divorce, by Race

Cause	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>ALL CAUSES</u>	<u>6,455</u>	<u>4,704</u>	<u>1,751</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
Cruel and inhuman treatment	3,887	3,352	535	60.2	71.2	30.6
Desertion	1,960	871	1,089	30.4	18.5	62.2
Adultery	292	206	86	4.5	4.4	4.9
Drunkenness	206	194	12	3.2	4.1	0.7
Penitentiary	47	33	14	0.7	0.7	0.8
Insanity	6	6	0	0.1	0.1	-
Annulment	18	16	2	0.3	0.3	0.1
Bigamy	16	7	9	0.2	0.1	0.5
Miscellaneous and other	11	10	1	0.2	0.2	*
Not stated	12	9	3	0.2	0.2	0.2

\* Less than 0.06 percent.

By Number of Minor Children, by Race

Number of Minor Children	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>6,455</u>	<u>4,704</u>	<u>1,751</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
None	2,828	1,967	861	43.8	41.8	49.2
One	1,359	1,076	283	21.0	22.9	16.2
Two	972	762	210	15.0	16.2	12.0
Three or more	1,208	853	355	18.7	18.1	20.3
Not stated	88	46	42	1.4	1.0	2.4

By Number of Years Married, by Race

Number of Years Married	Number			Percent		
	Total	White	Non-white	Total	White	Non-white
<u>TOTAL DIVORCES</u>	<u>6,455</u>	<u>4,704</u>	<u>1,751</u>	<u>100.0</u>	<u>100.0</u>	<u>100.0</u>
<u>Under five</u>	<u>2,651</u>	<u>2,261</u>	<u>390</u>	<u>41.1</u>	<u>48.1</u>	<u>22.3</u>
Under one	448	426	22	6.9	9.0	1.2
One	724	635	89	11.2	13.5	5.1
Two	569	478	91	8.8	10.2	5.2
Three	503	401	102	7.8	8.5	5.8
Four	407	321	86	6.3	6.8	4.9
Five - nine	1,242	943	299	19.2	20.0	17.1
Ten - nineteen	1,393	924	469	21.6	19.6	26.8
Twenty and over	1,116	540	576	17.3	11.5	32.9
Not stated	53	36	17	0.8	0.8	1.0

TABLE 11  
LIVE BIRTHS  
MISSISSIPPI  
1913-1967

By Year, by Race, with Rates, Occurrence Data 1913-1943,  
Residence Data 1944-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	41,103	21,874	19,229	21.9	26.6	18.2
1914	40,623	21,360	19,263	21.4	25.7	18.0
1915	39,484	20,999	18,485	20.5	23.8	17.1
1916	39,033	20,130	18,903	20.0	23.6	17.2
1917	37,141	19,056	18,085	18.8	22.0	16.3
1918	39,391	20,005	19,386	19.7	22.8	17.2
1919	42,491	22,028	20,463	21.0	24.9	18.0
1920	43,743	22,438	21,305	24.4	26.3	22.7
1921	46,482	23,703	22,779	25.6	27.3	24.1
1922	43,723	21,286	22,437	23.4	24.1	23.6
1923	43,266	21,140	22,126	23.3	23.6	23.1
1924	43,960	22,180	21,780	23.4	24.3	22.5
1925	45,563	22,265	23,298	24.0	24.1	23.9
1926	50,449	24,600	25,849	26.2	26.2	26.3
1927	49,204	24,163	25,041	25.3	25.3	25.3
1928	48,126	23,642	24,484	24.5	24.4	24.5
1929	45,570	22,105	23,465	22.9	22.5	23.3
1930	47,982	23,192	24,790	23.8	23.2	24.0
1931	45,123	21,637	23,486	22.2	21.4	22.9
1932	46,481	21,883	24,598	22.6	21.4	23.9
1933	44,184	20,407	23,777	21.3	19.7	22.9
1934	47,905	22,342	25,563	23.0	21.4	24.6
1935	48,291	21,972	26,319	23.0	20.8	25.1
1936	49,261	22,194	27,067	23.3	20.8	25.7
1937	52,170	23,285	28,885	24.4	21.6	27.3
1938	53,729	24,046	29,683	25.0	22.1	27.8
1939	52,288	23,505	28,783	24.1	21.4	26.8
1940	52,526	23,084	29,442	24.0	20.9	27.3
1941	54,561	24,167	30,394	24.7	21.6	28.0
1942	56,521	25,306	31,215	25.4	22.4	28.6
1943	59,408	27,611	31,797	28.9	26.4	31.6
1944#	56,586	25,647	30,939	28.3	25.2	31.6
1945#	53,540	23,426	30,114	26.9	23.1	30.9
1946#	61,430	29,715	31,715	26.8	25.4	28.2
1947#	66,291	32,423	33,868	28.6	27.5	29.9
1948#	65,291	29,666	35,625	28.0	24.9	31.2
1949#	66,128	28,786	37,342	28.1	24.0	32.5

Table 11 - Page 2  
Live Births - Mississippi - 1913-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	64,427	27,188	37,239	29.6	22.9	37.6
1951#	65,512	27,798	37,714	30.1	23.4	38.1
1952#	64,509	28,073	36,436	29.6	23.6	36.8
1953#	64,061	28,312	35,749	29.4	23.8	36.1
1954#	64,748	28,781	35,967	29.7	24.2	36.3
1955#	63,244	27,546	35,698	29.0	23.2	36.0
1956#	62,178	27,278	34,900	29.5	23.8	36.2
1957#	60,712	26,804	33,908	28.8	23.4	35.2
1958#	59,189	26,812	32,377	27.1	20.9	35.9
1959#	60,562	27,979	32,583	27.2	20.9	36.6
1960#	59,210	27,198	32,012	27.2	21.6	34.8
1961#	59,523	27,435	32,088	27.3	21.8	34.8
1962#	58,921	27,512	31,409	26.1	20.8	33.5
1963#	57,316	26,823	30,493	25.0	19.9	32.4
1964#	56,650	25,942	30,708	24.5	18.9	32.5
1965#	51,171	23,369	27,802	22.0	17.4	28.2
1966#	48,230	22,736	25,494	20.4	16.5	25.9
1967#	46,594	22,675	23,919	19.7	15.9	25.6

\* Births per 1,000 population.

1913-1920, population estimates based on U.S. Censuses of 1910 and 1920.

1921-1930, estimates based on U.S. Censuses of 1920 and 1930.

1931-1940, estimates based on U.S. Censuses of 1930 and 1940.

1941 and 1942, projected estimates based on U.S. Censuses of 1930 and 1940.

1943 and 1944, estimates of civilian population obtained from Ration Book Registration.

1945, estimate of civilian population, July 1, 1945, obtained from Bureau of Census Bulletin, Series P-46, No. 3, dated February 12, 1946.

1946-1949, projected estimates as of July 1, each year based on U.S. Censuses of 1930 and 1940.

1950-1955, U.S. Census population of April 1, 1950.

1956 and 1957, estimate as of July 1, 1956, prepared by Dr. Harald A. Pedersen, Sociology Department, Mississippi State University, with adjustments by Statistical Service, Mississippi State Board of Health.

1958, estimate as of January 1, prepared by Dr. George L. Wilber and Mr. Carl T. Robbins, Sociology Department, Mississippi State University.

1959, estimate as of January 1, prepared by Dr. George L. Wilber and Ellen Bryant, Sociology Department, Mississippi State University.

1960 and 1961, U.S. Census population of April 1, 1960.

1962-1967, estimates as of July 1, prepared by the Division of Sociology and Anthropology, Mississippi State University.

# On basis of place of residence.

TABLE 12  
DEATHS  
MISSISSIPPI  
1913-1967

By Year, by Race, with Rates, Occurrence Data 1913-1943,  
Residence Data 1944-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	21,305	7,643	13,662	11.4	9.3	13.0
1914	21,485	7,442	14,043	11.3	8.9	13.1
1915	24,001	8,088	15,913	12.5	9.6	14.7
1916	21,794	7,588	14,206	11.2	8.9	13.0
1917	23,579	8,048	15,531	11.9	9.3	14.0
1918	30,437	10,177	20,260	15.2	11.6	18.0
1919	22,667	8,142	14,525	12.7	9.5	15.5
1920	22,072	7,916	14,156	12.3	9.3	15.1
1921	19,991	7,308	12,683	11.0	8.4	13.4
1922	19,278	7,312	11,966	10.5	8.3	12.6
1923	20,363	7,688	12,675	11.0	8.6	13.2
1924	21,155	7,809	13,346	11.3	8.6	13.8
1925	22,229	8,143	14,086	11.7	8.8	14.4
1926	23,929	9,133	14,796	12.5	9.7	15.1
1927	23,257	8,765	14,492	12.0	9.2	14.6
1928	25,965	10,111	15,854	13.2	10.4	15.9
1929	25,868	10,048	15,820	13.0	10.2	15.7
1930	24,098	9,072	15,026	11.9	9.0	14.8
1931	22,007	8,641	13,366	10.8	8.5	13.0
1932	20,322	8,295	12,027	9.9	8.1	11.6
1933	21,617	8,898	12,719	10.4	8.6	12.2
1934	21,786	9,350	12,436	10.5	9.0	12.0
1935	21,285	9,186	12,099	10.1	8.7	11.6
1936	24,127	10,367	13,760	11.4	9.7	13.1
1937	24,005	10,055	13,950	11.2	9.3	13.2
1938	22,814	9,471	13,343	10.6	8.7	12.5
1939	22,701	9,464	13,237	10.5	8.6	12.3
1940	23,185	9,565	13,620	10.6	8.6	12.6
1941	22,573	9,442	13,131	10.2	8.4	12.1
1942	20,693	8,750	11,943	9.3	7.7	10.9
1943	20,850	9,139	11,711	10.2	8.7	11.6
1944#	20,199	8,830	11,369	10.1	8.7	11.6
1945#	19,024	8,452	10,572	9.6	8.3	10.8
1946#	19,321	8,995	10,326	8.4	7.7	9.2
1947#	20,007	9,238	10,769	8.6	7.8	9.5
1948#	20,183	9,216	10,967	8.6	7.7	9.6
1949#	20,489	9,233	11,256	8.7	7.7	9.8

Table 12 - Page 2  
 Deaths - Mississippi - 1913-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	20,567	9,466	11,101	9.4	8.0	11.2
1951#	21,007	9,593	11,414	9.6	8.1	11.5
1952#	20,657	9,387	11,270	9.5	7.9	11.4
1953#	20,492	9,495	10,997	9.4	8.0	11.1
1954#	19,353	9,453	9,900	8.9	8.0	10.0
1955#	19,403	9,501	9,902	8.9	8.0	10.0
1956#	19,780	9,969	9,811	9.4	8.7	10.2
1957#	20,476	10,339	10,137	9.7	9.0	10.5
1958#	21,541	10,811	10,730	9.9	8.4	11.9
1959#	20,890	10,742	10,148	9.4	8.0	11.4
1960#	21,713	11,204	10,509	10.0	8.9	11.4
1961#	21,232	11,063	10,169	9.7	8.8	11.0
1962#	22,713	11,868	10,845	10.0	9.0	11.6
1963#	23,339	12,118	11,221	10.2	9.0	11.9
1964#	22,559	12,039	10,520	9.7	8.8	11.1
1965#	22,652	12,376	10,276	9.7	9.2	10.4
1966#	23,350	12,976	10,374	9.9	9.4	10.5
1967#	22,609	12,816	9,793	9.6	9.0	10.5

\* Deaths per 1,000 population. See footnote on Table 11 for source of population estimates.

# On basis of place of residence.

TABLE 13  
MATERNAL DEATHS  
MISSISSIPPI  
1913-1967

By Year, by Race, with Rates, Occurrence Data 1913-1943,  
Residence Data 1944-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1913	447	162	285	108.8	74.1	148.2
1914	413	124	289	101.7	58.0	150.0
1915	289	88	201	73.2	41.9	108.7
1916	326	102	224	83.5	50.7	118.5
1917	297	105	192	80.0	55.1	106.2
1918	302	82	220	76.7	41.0	113.5
1919	407	173	234	95.8	78.5	114.4
1920	393	127	266	89.8	56.6	124.8
1921	391	145	246	84.1	61.2	108.0
1922	313	110	203	71.6	51.7	90.5
1923	321	108	213	74.2	51.1	96.3
1924	413	137	276	93.9	61.8	126.7
1925	435	143	292	95.5	64.2	125.3
1926	388	153	235	76.9	62.2	90.9
1927	409	134	275	83.1	55.4	109.8
1928	421	161	260	87.5	68.1	106.2
1929	410	152	258	90.0	68.8	110.0
1930	489	171	318	101.9	73.7	128.3
1931	352	131	221	78.0	60.5	94.1
1932	279	103	176	60.0	47.1	71.6
1933	315	116	199	71.3	56.8	83.7
1934	306	116	190	63.9	51.9	74.3
1935	334	137	197	69.2	62.4	74.8
1936	358	134	224	72.7	60.4	82.8
1937	370	150	220	70.9	64.4	76.2
1938	333	116	217	62.0	48.2	73.1
1939	311	97	214	59.5	41.3	74.3
1940	327	106	221	62.2	45.9	75.1
1941	293	93	200	53.7	38.5	65.8
1942	248	77	171	43.9	30.4	54.8
1943	233	66	167	39.2	23.9	52.5
1944#	225	72	153	39.8	28.1	49.4
1945#	216	64	152	40.3	27.3	50.5
1946#	199	67	132	32.4	22.5	41.6
1947#	175	50	125	26.4	15.4	36.9
1948#	177	38	139	27.1	12.8	39.0
1949#	147	32	115	22.2	11.1	30.8

Table 13 - Page 2  
 Maternal Deaths - Mississippi - 1913-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	165	30	135	25.6	11.0	36.2
1951#	143	28	115	21.8	10.1	30.5
1952#	119	15	104	18.4	5.3	28.5
1953#	119	24	95	18.6	8.5	26.6
1954#	104	11	93	16.1	3.8	25.8
1955#	76	12	64	12.0	4.4	17.9
1956#	74	12	62	11.9	4.4	17.8
1957#	75	14	61	12.4	5.2	18.0
1958#	53	10	43	9.0	3.7	13.3
1959#	55	9	46	9.1	3.2	14.1
1960#	59	8	51	10.0	2.9	15.9
1961#	49	9	40	8.2	3.3	12.5
1962#	55	7	48	9.3	2.5	15.3
1963#	44	9	35	7.7	3.4	11.5
1964#	57	9	48	10.1	3.5	15.6
1965#	40	3	37	7.8	1.3	13.3
1966#	27	6	21	5.6	2.6	8.2
1967#	36	10	26	7.7	4.4	10.9

\* Deaths per 10,000 live births.

# On basis of place of residence.

TABLE 14  
DEATHS UNDER 1 YEAR  
MISSISSIPPI  
1917-1967

By Year, by Race, with Rates, Occurrence Data 1917-1943,  
Residence Data 1944-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1917	3,689	1,296	2,393	99.3	68.0	132.3
1918	**	**	**	**	**	**
1919	3,554	**	**	83.6	**	**
1920	3,499	1,344	2,155	80.0	59.9	101.1
1921	3,157	1,243	1,914	67.9	52.4	84.0
1922	2,826	1,067	1,759	64.6	50.1	78.4
1923	2,908	1,104	1,804	67.2	52.2	81.5
1924	3,137	1,188	1,949	71.4	53.6	89.5
1925	3,176	1,179	1,997	69.7	53.0	85.7
1926	3,578	1,447	2,131	70.9	58.8	82.4
1927	3,291	1,303	1,988	66.9	54.0	79.4
1928	3,567	1,416	2,151	74.1	59.9	87.9
1929	3,275	1,268	2,007	71.9	57.4	85.5
1930	3,276	1,182	2,094	68.3	51.0	84.5
1931	2,547	961	1,586	56.4	44.4	67.5
1932	2,482	965	1,517	53.4	44.1	61.7
1933	2,800	1,035	1,765	63.4	50.7	74.2
1934	3,070	1,199	1,871	64.1	53.7	73.2
1935	2,587	1,044	1,543	53.6	47.5	58.6
1936	2,883	1,105	1,778	58.5	49.8	65.7
1937	3,098	1,174	1,924	59.4	50.4	66.6
1938	3,051	1,205	1,846	56.8	50.1	62.2
1939	2,908	1,135	1,773	55.6	48.3	61.6
1940	2,864	1,070	1,794	54.5	46.4	60.9
1941	2,969	1,113	1,856	54.4	46.1	61.1
1942	2,654	957	1,697	47.0	37.8	54.4
1943	2,774	1,137	1,637	46.7	41.2	51.5
1944#	2,489	974	1,515	44.0	38.0	49.0
1945#	2,192	846	1,346	40.9	36.1	44.7
1946#	2,298	1,005	1,293	37.4	33.8	40.8
1947#	2,428	1,009	1,419	36.6	31.1	41.9
1948#	2,470	988	1,482	37.8	33.3	41.6
1949#	2,612	900	1,712	39.5	31.3	45.8

Table 14 - Page 2  
 Deaths Under 1 Year - Mississippi - 1917-1967

Year	Number			Rate*		
	Total	White	Non-white	Total	White	Non-white
1950#	2,348	763	1,585	36.4	28.1	42.6
1951#	2,589	814	1,775	39.5	29.3	47.1
1952#	2,592	760	1,832	40.2	27.1	50.3
1953#	2,534	807	1,727	39.6	28.5	48.3
1954#	2,360	744	1,616	36.4	25.8	44.9
1955#	2,338	680	1,658	37.0	24.7	46.4
1956#	2,353	734	1,619	37.8	26.9	46.4
1957#	2,257	694	1,563	37.2	25.9	46.1
1958#	2,424	706	1,718	41.0	26.3	53.1
1959#	2,355	677	1,678	38.9	24.2	51.5
1960#	2,457	714	1,743	41.5	26.2	54.4
1961#	2,250	645	1,605	37.8	23.5	50.0
1962#	2,353	705	1,648	39.9	25.6	52.5
1963#	2,372	610	1,762	41.4	22.7	57.8
1964#	2,230	599	1,631	39.4	23.1	53.1
1965#	2,126	593	1,533	41.5	25.4	55.1
1966#	1,856	520	1,336	38.5	22.9	52.4
1967#	1,645	512	1,133	35.3	22.6	47.4

\* Deaths per 1,000 live births.

\*\* Not available.

# On basis of place of residence.

TABLE 15  
DEATHS UNDER 1 MONTH\*  
MISSISSIPPI  
1924-1967

By Year, by Race, with Rates, Occurrence Data 1924-1944,  
Residence Data 1945-1967

Year	Number			Rate**		
	Total	White	Non-white	Total	White	Non-white
1924	1,598	730	868	36.4	32.9	39.8
1925	1,639	701	938	36.0	31.5	40.3
1926	1,731	823	908	34.3	33.4	35.1
1927	1,644	786	858	33.4	32.5	34.3
1928	1,788	804	984	37.2	34.0	40.2
1929	1,662	773	889	36.5	35.0	37.9
1930	1,692	715	977	35.3	30.8	39.4
1931	1,339	590	749	29.7	27.3	31.9
1932	1,361	572	789	29.3	26.1	32.1
1933	1,540	636	904	34.8	31.2	38.0
1934	1,505	644	861	31.4	28.8	33.7
1935	1,446	642	804	29.9	29.2	30.5
1936	1,494	676	818	30.3	30.4	30.2
1937	1,678	755	923	32.2	32.4	32.0
1938	1,688	747	941	31.4	31.1	31.7
1939	1,637	735	902	31.3	31.3	31.3
1940	1,613	680	933	30.7	29.4	31.7
1941	1,663	734	929	30.5	30.4	30.6
1942	1,544	638	906	27.3	25.2	29.0
1943	1,652	791	861	27.8	28.6	27.1
1944	1,436	661	775	25.2	25.4	25.1
1945#	1,351	624	727	25.2	26.6	24.1
1946#	1,561	794	767	25.4	26.7	24.2
1947#	1,595	750	845	24.1	23.1	24.9
1948#	1,674	773	901	25.6	26.0	25.3
1949#	1,594	661	933	24.1	23.0	25.0
1950#	1,446	560	886	22.4	20.6	23.8
1951#	1,679	626	1,053	25.6	22.5	27.9
1952#	1,658	581	1,077	25.7	20.7	29.6
1953#	1,630	635	995	25.4	22.4	27.8
1954#	1,521	601	920	23.5	20.9	25.6
1955#	1,440	527	913	22.8	19.1	25.6
1956#	1,451	581	870	23.3	21.3	24.9
1957#	1,425	540	885	23.5	20.1	26.1
1958#	1,475	540	935	24.9	20.1	28.9
1959#	1,450	515	935	23.9	18.4	28.7
1960#	1,497	550	947	25.3	20.2	29.6
1961#	1,401	509	892	23.5	18.6	27.8
1962#	1,456	550	906	24.7	20.0	28.8
1963#	1,449	470	979	25.3	17.5	32.1
1964#	1,372	460	912	24.2	17.7	29.7
1965#	1,283	470	813	25.1	20.1	29.2
1966#	1,123	386	737	23.3	17.0	28.9
1967#	1,053	388	665	22.6	17.1	27.8

\* Figures for 1949-1967 include only the deaths under 28 days.

\*\* Deaths per 1,000 live births.

# On basis of place of residence.

TABLE 16  
DEATHS FROM HEART DISEASES, VASCULAR LESIONS,  
MALIGNANT NEOPLASMS, ACCIDENTS, AND INFLUENZA AND PNEUMONIA  
MISSISSIPPI  
1958-1967

By Year, by Race, with Rates, Residence Data

Year	Heart Diseases (400-443)				Vascular Lesions (330-334)				Malignant Neoplasms (140-205)				Accidents (800-962)				Influenza & Pneumonia (480-493)					
	Total		White		Non-white		Total		White		Non-white		Total		White		Non-white		Total		White	
	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	Total	White	Non-white	
1958 No. Rate*	6,528	3,929	2,599	2,867	1,426	1,441	2,525	1,524	1,001	1,379	729	650	909	344	565	41.6	26.8	41.6	26.8	41.6	26.7	
1959 No. Rate*	6,233	3,774	2,459	2,683	1,386	1,297	2,600	1,580	1,020	1,501	837	664	768	328	440	34.5	24.6	34.5	24.6	34.5	24.4	
1960 No. Rate*	6,321	3,882	2,439	2,960	1,513	1,447	2,635	1,644	991	1,504	827	658	958	389	569	44.0	30.9	44.0	30.9	44.0	31.8	
1961 No. Rate*	6,561	4,022	2,539	2,891	1,464	1,427	2,748	1,732	1,016	1,427	760	667	721	307	414	72.4	33.1	72.4	33.1	72.4	34.0	
1962 No. Rate*	7,077	4,206	2,871	3,112	1,552	1,560	2,827	1,793	1,034	1,557	846	711	879	362	517	75.9	38.9	75.9	38.9	75.9	38.9	
1963 No. Rate*	7,300	4,396	2,904	3,154	1,650	1,504	2,945	1,859	1,086	1,571	851	720	1,113	420	693	76.5	48.6	76.5	48.6	76.5	48.6	
1964 No. Rate*	6,981	4,313	2,668	3,077	1,588	1,489	2,968	1,861	1,107	1,612	898	714	791	308	483	75.5	34.2	75.5	34.2	75.5	34.2	
1965 No. Rate*	7,038	4,429	2,609	3,064	1,639	1,425	2,931	1,867	1,064	1,610	881	729	764	350	414	74.0	32.8	74.0	32.8	74.0	32.8	
1966 No. Rate*	7,468	4,729	2,739	3,132	1,681	1,451	3,098	1,990	1,108	1,872	1,072	800	773	341	432	79.3	32.8	79.3	32.8	79.3	32.8	
1967 No. Rate*	7,300	4,640	2,660	2,981	1,632	1,349	3,136	2,061	1,075	1,890	1,166	724	657	310	347	81.7	27.8	81.7	27.8	81.7	27.7	

\* Deaths per 100,000 population.

TABLE 17  
MARRIAGES AND DIVORCES  
MISSISSIPPI  
1926-1967

By Year, by Race, Occurrence Data

Year	Marriages			Divorces		
	Total	White	Non-white	Total	White	Non-white
1926*	28,712			2,766		
1927*	29,048#	11,402	17,383	2,895		
1928	30,231#	11,662	18,257	3,009	1,435	1,574
1929	31,495#	12,864	18,388	3,172	1,598	1,574
1930	25,728#	10,815	14,751	2,743	1,487	1,256
1931	21,589#	9,952	11,629	2,015	1,308	707
1932	22,688#	10,147	12,453	1,918	1,308	610
1933	27,567#	12,201	15,300	1,955	1,396	559
1934	30,149#	13,497	16,541	2,446	1,704	742
1935	28,722#	12,955	15,635	2,425	1,681	744
1936	32,480#	14,120	18,196	2,520	1,690	830
1937	32,055#	14,802	17,246	2,977	1,984	993
1938	31,495	14,224	17,271	2,916	2,069	847
1939	31,956	15,370	16,586	3,034	2,166	868
1940	34,088	16,834	17,254	3,251	2,295	956
1941	42,142	21,611	20,531	3,831	2,444	1,387
1942	49,565	27,227	22,338	4,326	2,606	1,720
1943	46,030	23,445	22,585	5,851	3,062	2,789
1944	41,559	21,173	20,386	6,742	3,425	3,317
1945	44,871	24,738	20,133	6,727	3,919	2,808
1946	64,204	37,099	27,105	9,495	5,867	3,628
1947	53,584	30,514	23,070	7,004	4,323	2,681
1948	53,645	31,504	22,141	6,697	3,864	2,833
1949	52,765	32,191	20,574	6,285	3,621	2,664
1950	56,738	37,871	18,867	6,065	3,711	2,354
1951	57,321	37,857	19,464	5,464	3,601	1,863
1952	58,116	39,162	18,954	5,419	3,607	1,812
1953	61,751	42,961	18,790	5,204	3,515	1,689
1954	61,381	43,268	18,113	5,001	3,424	1,577
1955	66,423	47,618	18,805	4,845	3,372	1,473
1956	66,223	48,140	18,083	5,027	3,452	1,575
1957	62,495	46,868	15,627	4,974	3,350	1,624
1958	36,402	26,998	9,404	5,016	3,485	1,531
1959	20,447	13,547	6,900	5,108	3,557	1,551
1960	21,233	14,029	7,204	5,260	3,553	1,707
1961	21,610	14,438	7,172	5,213	3,545	1,668
1962	22,007	14,432	7,575	5,467	3,827	1,640
1963	22,140	14,508	7,632	5,452	3,930	1,522
1964	22,784	15,159	7,625	5,631	4,077	1,554
1965	23,343	15,613	7,730	5,730	4,220	1,510
1966	23,476	15,860	7,616	6,306	4,650	1,656
1967	23,535	15,752	7,783	6,455	4,704	1,751

\* Total number of marriages for 1926 and total number of divorces for 1926 and 1927 furnished by U.S. Bureau of Census; distribution by race not available.

# Includes some records on which the race of the couple was not stated.

## BIRTHS, FETAL DEATHS, DEATHS

THE COUNTIES AND THE STATE OF MISSISSIPPI, 1967

## Footnotes for Tables on Pages 57-139

\* Total births and deaths per 1,000 population. Fetal deaths, deaths under 1 year, deaths under 28 days, and deaths from diseases of early infancy per 1,000 live births. Maternal deaths per 10,000 live births. Deaths from specific causes per 100,000 population.

\*\* Rate less than 0.06.

# Pneumonia and diarrhea of the newborn (under 28 days) included in diseases of early infancy.

↓  
per 100,000



BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
ADAMS COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,356	654	702	868	414	454	22.3	19.6	25.5
Fetal deaths	80	43	37	54	32	22	62.2	77.3	48.4
Total deaths	496	226	270	401	180	221	10.3	8.5	12.4
Deaths under 1 year	53	16	37	35	12	23	40.3	29.0	50.7
Deaths under 28 days	34	13	21	23	11	12	26.5	26.6	26.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	2.6	-	5.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	1	0	1	2.6	-	5.6
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	72	46	26	60	36	24	154.2	170.8	134.7
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	2.6	-	5.6
Asthma (241)	2	0	2	2	0	2	5.1	-	11.2
Diabetes mellitus (260)	16	8	8	10	6	4	25.7	28.5	22.4
Anemias (290-293)	3	0	3	2	0	2	5.1	-	11.2
Intracranial lesions (330-334)	63	16	47	49	12	37	126.0	56.9	207.6
Nonmeningococcal meningitis (340)	1	1	0	0	0	0	-	-	-
Heart diseases (400-443)	154	74	80	125	56	69	321.3	265.6	387.2
Other hypertensive disease (444-447)	6	2	4	4	1	3	10.3	4.7	16.8
General arteriosclerosis (450)	17	7	10	12	7	5	30.8	33.2	28.0
Other circulatory diseases (451-468)	4	3	1	4	2	2	10.3	9.5	11.2
#Influenza & pneumonia (480-493)	17	6	11	9	4	5	23.1	19.0	28.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	7	5	2	6	5	1	15.4	23.7	5.6
Stomach & duodenal ulcers (540, 541)	2	1	1	2	1	1	5.1	4.7	5.6
Hernia & intestinal obstruction (560, 561, 570)	4	2	2	3	2	1	7.7	9.5	5.6
#Gastritis, etc. (543, 571, 572)	4	1	3	3	0	3	7.7	-	16.8
Cirrhosis of liver (581)	2	1	1	3	2	1	7.7	9.5	5.6
Acute nephritis (590, 591)	1	0	1	1	0	1	2.6	-	5.6
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	0	1	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	6	3	3	6	3	3	15.4	14.2	16.8
Diseases of early infancy (760-776)	30	10	20	20	8	12	23.0	19.3	26.4
Accidents (800-962)	34	20	14	38	21	17	97.7	99.6	95.4
Motor vehicle accidents (810-835)	13	11	2	18	14	4	46.3	66.4	22.4
Suicide (963, 970-979)	2	2	0	2	2	0	5.1	9.5	-
Homicide (964, 980-985)	10	0	10	10	0	10	25.7	-	56.1
Ill-defined (780-795)	15	4	11	14	4	10	36.0	19.0	56.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
ALCORN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	431	346	85	466	386	80	17.4	16.3	25.3
Fetal deaths	8	5	3	13	8	5	27.9	20.7	62.5
Total deaths	242	203	39	255	210	45	9.5	8.9	14.2
Deaths under 1 year	12	8	4	12	8	4	25.8	20.7	50.0
Deaths under 28 days	7	6	1	6	5	1	12.9	13.0	12.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	29	29	0	29	28	1	108.2	118.4	31.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	3	2	1	3	2	1	11.2	8.4	31.6
Diabetes mellitus (260)	6	3	3	6	4	2	22.4	16.9	63.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	23	5	32	26	6	119.4	109.9	189.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	93	78	15	90	74	16	335.7	312.9	506.3
Other hypertensive disease (444-447)	4	2	2	4	2	2	14.9	8.4	63.3
General arteriosclerosis (450)	2	2	0	3	3	0	11.2	12.7	-
Other circulatory diseases (451-468)	1	1	0	2	1	1	7.4	4.2	31.6
#Influenza & pneumonia (480-493)	5	4	1	7	6	1	26.1	25.4	31.6
Bronchitis (500-502)	3	3	0	2	2	0	7.4	8.4	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	2	2	0	2	2	0	7.4	8.4	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	3.7	4.2	-
#Gastritis, etc. (543, 571, 572)	1	1	0	1	1	0	3.7	4.2	-
Cirrhosis of liver (581)	0	0	0	2	2	0	7.4	8.4	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	3.7	4.2	-
Infections of kidney (600)	2	2	0	1	1	0	3.7	4.2	-
Hyperplasia of prostate (610)	0	0	0	1	1	0	3.7	4.2	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	3	3	0	11.2	12.7	-
Diseases of early infancy (760-776)	6	5	1	5	4	1	10.7	10.4	12.5
Accidents (800-962)	23	20	3	25	20	5	93.2	84.6	158.2
Motor vehicle accidents (810-835)	16	15	1	18	16	2	67.1	67.6	63.3
Suicide (963, 970-979)	4	4	0	5	5	0	18.6	21.1	-
Homicide (964, 980-985)	3	2	1	4	3	1	14.9	12.7	31.6
Ill-defined (780-795)	14	10	4	13	9	4	48.5	38.0	126.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
AMITE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	98	1	97	268	86	182	18.0	13.0	22.0
Fetal deaths	1	0	1	8	2	6	29.8	23.2	33.0
Total deaths	77	35	42	137	69	68	9.2	10.4	8.2
Deaths under 1 year	7	1	6	9	0	9	33.6	-	49.4
Deaths under 28 days	2	0	2	3	0	3	11.2	-	16.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	1	1	0	6.7	15.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	12	7	5	25	13	12	167.7	196.1	144.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	6.7	15.1	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	3	1	2	17	10	7	114.0	150.8	84.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	31	17	14	47	28	19	315.2	422.3	229.5
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	4	3	1	26.8	45.2	12.1
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	5	2	3	7	2	5	46.9	30.2	60.4
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	0	1	2	1	1	13.4	15.1	12.1
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	6.7	15.1	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	6.7	-	12.1
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	4	0	4	5	0	5	18.6	-	27.5
Accidents (800-962)	11	5	6	13	7	6	87.2	105.6	72.5
Motor vehicle accidents (810-835)	6	3	3	7	4	3	46.9	60.3	36.2
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	2	0	2	13.4	-	24.2
Ill-defined (780-795)	5	0	5	9	2	7	60.4	30.2	84.5

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
ATTALA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	404	175	229	383	149	234	15.8	10.0	24.7
Fetal deaths	17	7	10	17	7	10	44.4	47.0	42.7
Total deaths	228	144	84	244	154	90	10.0	10.4	9.5
Deaths under 1 year	14	4	10	15	4	11	39.2	26.8	47.0
Deaths under 28 days	5	2	3	6	2	4	15.7	13.4	17.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	4.1	6.7	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	32	20	12	36	23	13	148.1	155.1	137.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	4.1	6.7	-
Diabetes mellitus (260)	6	2	4	8	3	5	32.9	20.2	52.7
Anemias (290-293)	1	1	0	1	1	0	4.1	6.7	-
Intracranial lesions (330-334)	39	24	15	36	22	14	148.1	148.3	147.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	77	54	23	84	59	25	345.5	397.8	263.7
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	4	3	1	3	3	0	12.3	20.2	-
Other circulatory diseases (451-468)	1	1	0	2	2	0	8.2	13.5	-
#Influenza & pneumonia (480-493)	7	1	6	8	2	6	32.9	13.5	63.3
Bronchitis (500-502)	1	1	0	1	1	0	4.1	6.7	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	8.2	13.5	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	2	1	1	8.2	6.7	10.5
#Gastritis, etc. (543, 571, 572)	4	3	1	3	2	1	12.3	13.5	10.5
Cirrhosis of liver (581)	2	2	0	3	3	0	12.3	20.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	12.3	13.5	10.5
Infections of kidney (600)	2	2	0	1	1	0	4.1	6.7	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	0	3	3	0	3	12.3	-	31.6
Diseases of early infancy (760-776)	6	2	4	7	2	5	18.3	13.4	21.4
Accidents (800-962)	15	12	3	16	11	5	65.8	74.2	52.7
Motor vehicle accidents (810-835)	10	9	1	6	6	0	24.7	40.4	-
Suicide (963, 970-979)	1	1	0	1	1	0	4.1	6.7	-
Homicide (964, 980-985)	2	0	2	2	0	2	8.2	-	21.1
Ill-defined (780-795)	12	6	6	11	5	6	45.2	33.7	63.3

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
BENTON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	41	1	40	135	65	70	17.6	14.4	22.4
Fetal deaths	4	0	4	5	1	4	37.0	15.4	57.1
Total deaths	26	17	9	63	42	21	8.2	9.3	6.7
Deaths under 1 year	1	0	1	5	2	3	37.0	30.8	42.8
Deaths under 28 days	0	0	0	3	2	1	22.2	30.8	14.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	13.1	22.1	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	3	2	1	12	8	4	156.9	177.0	127.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	2	2	11	6	5	143.8	132.7	159.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	8	5	3	21	16	5	274.5	354.0	159.7
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	1	0	1	1	0	1	13.1	-	31.9
#Influenza & pneumonia (480-493)	0	0	0	0	0	0	-	-	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	13.1	-	31.9
Cirrhosis of liver (581)	0	0	0	1	1	0	13.1	22.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	0	1	74.1	-	142.8
Congenital malformations (750-759)	0	0	0	1	0	1	13.1	-	31.9
Diseases of early infancy (760-776)	0	0	0	3	2	1	22.2	30.8	14.3
Accidents (800-962)	4	4	0	6	5	1	78.4	110.6	31.9
Motor vehicle accidents (810-835)	3	3	0	4	3	1	52.3	66.4	31.9
Suicide (963, 970-979)	1	1	0	1	1	0	13.1	22.1	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	3	2	1	2	1	1	26.1	22.1	31.9

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
BOLIVAR COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,468	298	1,170	1,338	291	1,047	23.3	14.2	28.3
Fetal deaths	58	10	48	53	13	40	39.6	44.7	38.2
Total deaths	551	124	427	560	165	395	9.7	8.0	10.7
Deaths under 1 year	62	9	53	54	7	47	40.4	24.0	44.9
Deaths under 28 days	36	9	27	29	6	23	21.7	20.6	22.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	3	2	1	5.2	9.8	2.7
Syphilis (020-029)	0	0	0	1	0	1	1.7	-	2.7
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	56	16	40	63	25	38	109.6	121.9	102.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	0	4	4	0	4	7.0	-	10.8
Anemias (290-293)	2	0	2	3	0	3	5.2	-	8.1
Intracranial lesions (330-334)	107	24	83	104	31	73	180.9	151.1	197.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	175	48	127	168	57	111	292.2	277.9	300.1
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	9	3	6	9	4	5	15.6	19.5	13.5
Other circulatory diseases (451-468)	5	1	4	6	2	4	10.4	9.8	10.8
#Influenza & pneumonia (480-493)	19	1	18	19	3	16	33.0	14.6	43.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	5	1	4	5	1	4	8.7	4.9	10.8
Stomach & duodenal ulcers (540, 541)	2	1	1	3	1	2	5.2	4.9	5.4
Hernia & intestinal obstruction (560, 561, 570)	2	0	2	3	0	3	5.2	-	8.1
#Gastritis, etc. (543, 571, 572)	9	0	9	9	0	9	15.6	-	24.3
Cirrhosis of liver (581)	2	1	1	5	2	3	8.7	9.8	8.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	17	0	17	11	0	11	19.1	-	29.7
Infections of kidney (600)	4	2	2	4	1	3	7.0	4.9	8.1
Hyperplasia of prostate (610)	0	0	0	1	0	1	1.7	-	2.7
Maternal causes (640-689)	1	0	1	1	0	1	7.5	-	9.6
Congenital malformations (750-759)	4	0	4	4	1	3	7.0	4.9	8.1
Diseases of early infancy (760-776)	31	9	22	25	6	19	18.7	20.6	18.1
Accidents (800-962)	24	6	18	36	18	18	62.6	87.8	48.7
Motor vehicle accidents (810-835)	6	2	4	16	11	5	27.8	53.6	13.5
Suicide (963, 970-979)	2	2	0	3	2	1	5.2	9.8	2.7
Homicide (964, 980-985)	1	0	1	2	0	2	3.5	-	5.4
Ill-defined (780-795)	50	5	45	49	5	44	85.2	24.4	119.0

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
CALHOUN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	252	96	156	322	189	133	20.7	16.4	33.1
Fetal deaths	6	2	4	11	6	5	34.2	31.7	37.6
Total deaths	158	121	37	198	157	41	12.7	13.6	10.2
Deaths under 1 year	8	4	4	10	6	4	31.0	31.7	30.1
Deaths under 28 days	6	4	2	7	5	2	21.7	26.4	15.0
<b>SELECTED CAUSES OF DEATH</b>									
AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	6.4	8.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	14	8	6	31	24	7	199.6	208.5	174.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	2	1	4	3	1	25.8	26.1	24.9
Anemias (290-293)	1	0	1	1	0	1	6.4	-	24.9
Intracranial lesions (330-334)	16	14	2	22	19	3	141.7	165.1	74.6
Nonmeningococcal meningitis (340)	0	0	0	1	1	0	6.4	8.7	-
Heart diseases (400-443)	63	54	9	69	61	8	444.3	530.0	199.0
Other hypertensive disease (444-447)	4	3	1	4	2	2	25.8	17.4	49.8
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	2	1	1	12.9	8.7	24.9
#Influenza & pneumonia (480-493)	7	7	0	4	4	0	25.8	34.8	-
Bronchitis (500-502)	0	0	0	1	1	0	6.4	8.7	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	12.9	17.4	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	6.4	8.7	-
#Gastritis, etc. (543, 571, 572)	2	1	1	1	1	0	6.4	8.7	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	19.3	17.4	24.9
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	0	1	2	0	2	12.9	-	49.8
Diseases of early infancy (760-776)	5	3	2	7	5	2	21.7	26.4	15.0
Accidents (800-962)	22	14	8	26	18	8	167.4	156.4	199.0
Motor vehicle accidents (810-835)	8	8	0	11	11	0	70.8	95.6	-
Suicide (963, 970-979)	1	1	0	1	1	0	6.4	8.7	-
Homicide (964, 980-985)	0	0	0	2	0	2	12.9	-	49.8
Ill-defined (780-795)	4	3	1	3	2	1	19.3	17.4	24.9

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 CARROLL COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	86	6	80	182	58	124	16.8	10.9	22.7
Fetal deaths	1	0	1	6	4	2	33.0	69.0	16.1
Total deaths	49	20	29	108	58	50	10.0	10.9	9.2
Deaths under 1 year	8	0	8	10	0	10	54.9	-	80.6
Deaths under 28 days	4	0	4	6	0	6	33.0	-	48.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	5	2	3	16	11	5	148.1	206.0	91.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	9.2	18.7	-
Diabetes mellitus (260)	0	0	0	1	1	0	9.2	18.7	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	2	2	19	8	11	175.9	149.8	201.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	9	6	3	24	16	8	222.2	299.6	146.5
Other hypertensive disease (444-447)	2	0	2	2	0	2	18.5	-	36.6
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	2	2	0	18.5	37.4	-
#Influenza & pneumonia (480-493)	2	0	2	4	1	3	37.0	18.7	54.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	1	1	0	9.2	18.7	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	9.2	-	18.3
#Gastritis, etc. (543, 571, 572)	0	0	0	4	4	0	37.0	74.9	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	1	1	0	9.2	18.7	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	3	0	3	5	0	5	27.5	-	40.3
Accidents (800-962)	7	4	3	8	5	3	74.1	93.6	54.9
Motor vehicle accidents (810-835)	4	3	1	6	4	2	55.6	74.9	36.6
Suicide (963, 970-979)	1	1	0	0	0	1	18.5	18.7	18.3
Homicide (964, 980-985)	1	1	0	2	1	1	111.1	37.4	183.2
Ill-defined (780-795)	12	3	9	12	2	10	-	-	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
CHICKASAW COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	363	168	195	392	195	197	20.8	16.8	27.2
Fetal deaths	22	8	14	12	6	6	30.6	30.8	30.4
Total deaths	233	140	93	217	134	83	11.5	11.6	11.5
Deaths under 1 year	9	6	3	7	5	2	17.8	25.6	10.2
Deaths under 28 days	4	4	0	3	3	0	7.6	15.4	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	5.3	-	13.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	5.3	-	13.8
Malignant neoplasms (140-205)	41	32	9	39	31	8	207.1	267.5	110.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	5.3	-	13.8
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	39	16	23	33	15	18	175.2	129.4	248.6
Nonmeningococcal meningitis (340)	1	0	1	0	0	0	-	-	-
Heart diseases (400-443)	84	56	28	77	50	27	408.9	431.4	372.9
Other hypertensive disease (444-447)	12	5	7	11	4	7	58.4	34.5	96.7
General arteriosclerosis (450)	3	2	1	4	3	1	21.2	25.9	13.8
Other circulatory diseases (451-468)	1	1	0	3	2	1	15.9	17.2	13.8
#Influenza & pneumonia (480-493)	3	2	1	0	0	0	-	-	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	1	0	2	2	0	10.6	17.2	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	0	1	5.3	-	13.8
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	5.3	8.6	-
Cirrhosis of liver (581)	0	0	0	1	1	0	5.3	8.6	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	4	2	2	3	1	2	15.9	8.6	27.6
Infections of kidney (600)	2	1	1	2	1	1	10.6	8.6	13.8
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	25.5	-	50.8
Congenital malformations (750-759)	1	0	1	0	0	0	-	-	-
Diseases of early infancy (760-776)	4	4	0	3	3	0	7.6	15.4	-
Accidents (800-962)	19	12	7	17	13	4	90.3	112.2	55.2
Motor vehicle accidents (810-835)	14	7	7	11	7	4	58.4	60.4	55.2
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	4	1	3	4	1	3	21.2	8.6	41.4
Ill-defined (780-795)	6	1	5	5	1	4	26.6	8.6	55.2

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 CHOCTAW COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	136	53	83	135	78	57	15.7	12.3	25.0
Fetal deaths	3	0	3	3	1	2	22.2	12.8	35.1
Total deaths	77	50	27	106	80	26	12.3	12.6	11.4
Deaths under 1 year	3	1	2	5	3	2	37.0	38.5	35.1
Deaths under 28 days	0	0	0	1	1	0	7.4	12.8	-
<b>SELECTED CAUSES OF DEATH</b>									
AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	1	0	1	11.6	-	43.8
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	1	0	1	1	0	11.6	15.8	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	2	2	0	8	7	1	93.0	110.8	43.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	2	1	4	3	1	46.5	47.5	43.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	14	6	8	18	11	7	209.3	174.0	307.0
Nonmeningococcal meningitis (340)	0	0	0	2	1	1	23.2	15.8	43.8
Heart diseases (400-443)	31	23	8	42	36	6	488.4	569.6	263.2
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	4	1	3	3	2	1	34.9	31.6	43.8
Other circulatory diseases (451-468)	5	4	1	4	3	1	46.5	47.5	43.8
#Influenza & pneumonia (480-493)	2	0	2	3	1	2	34.9	15.8	87.7
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	3	0	3	3	0	34.9	47.5	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	11.6	15.8	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	11.6	-	43.8
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	1	1	0	11.6	15.8	-
Infections of kidney (600)	0	0	0	2	2	0	23.2	31.6	-
Hyperplasia of prostate (610)	0	0	0	1	0	1	11.6	-	43.8
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	1	1	0	7.4	12.8	-
Accidents (800-962)	3	3	0	4	3	1	46.5	47.5	43.8
Motor vehicle accidents (810-835)	1	1	0	2	1	1	23.2	15.8	43.8
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	1	0	1	1	0	11.6	15.8	-
Ill-defined (780-795)	2	1	1	3	2	1	34.9	31.6	43.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
CLAIBORNE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	105	20	85	208	44	164	20.7	14.3	23.5
Fetal deaths	2	1	1	10	0	10	48.1	-	61.0
Total deaths	102	23	79	141	41	100	14.0	13.4	14.3
Deaths under 1 year	5	0	5	9	0	9	43.3	-	54.9
Deaths under 28 days	2	0	2	6	0	6	28.8	-	36.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	5	3	2	11	5	6	109.6	162.9	86.1
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	10.0	-	14.3
Asthma (241)	1	1	0	1	1	0	10.0	32.6	-
Diabetes mellitus (260)	1	0	1	1	0	1	10.0	-	14.3
Anemias (290-293)	1	0	1	2	0	2	19.9	-	28.7
Intracranial lesions (330-334)	5	1	4	12	4	8	119.5	130.3	114.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	54	14	40	64	21	43	637.4	684.0	616.9
Other hypertensive disease (444-447)	3	1	2	3	1	2	29.9	32.6	28.7
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	1	0	1	10.0	-	14.3
#Influenza & pneumonia (480-493)	3	0	3	3	0	3	29.9	-	43.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	1	1	0	10.0	32.6	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	10.0	32.6	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	10.0	32.6	-
Cirrhosis of liver (581)	2	1	1	2	1	1	19.9	32.6	14.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	1	4	5	1	4	49.8	32.6	57.4
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	2	1	1	96.2	227.3	61.0
Congenital malformations (750-759)	1	0	1	1	0	1	10.0	-	14.3
Diseases of early infancy (760-776)	1	0	1	5	0	5	24.0	-	30.5
Accidents (800-962)	7	0	7	9	0	9	89.6	-	129.1
Motor vehicle accidents (810-835)	1	0	1	3	0	3	29.9	-	43.0
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	2	0	2	2	0	2	19.9	-	28.7
Ill-defined (780-795)	8	0	8	10	1	9	99.6	32.6	129.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 CLARKE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	82	49	33	312	151	161	20.3	15.8	27.7
Fetal deaths	5	1	4	17	6	11	54.5	39.7	68.3
Total deaths	111	61	50	182	116	66	11.8	12.2	11.3
Deaths under 1 year	7	1	6	15	4	11	48.1	26.5	68.3
Deaths under 28 days	5	1	4	14	4	10	44.9	26.5	62.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	6.5	-	17.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	7	5	2	15	11	4	97.7	115.4	68.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	1	0	1	6.5	-	17.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	14	10	34	22	12	221.5	230.8	206.2
Nommeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	28	17	11	56	44	12	364.8	461.7	206.2
Other hypertensive disease (444-447)	0	0	0	1	1	0	6.5	10.5	-
General arteriosclerosis (450)	6	2	4	6	2	4	39.1	21.0	68.7
Other circulatory diseases (451-468)	1	1	0	4	3	1	26.0	31.5	17.2
#Influenza & pneumonia (480-493)	4	2	2	5	2	3	32.6	21.0	51.5
Bronchitis (500-502)	1	1	0	1	1	0	6.5	10.5	-
Other bronchopul. diseases (525-527)	0	0	0	2	1	1	13.0	10.5	17.2
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	6.5	10.5	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	6.5	10.5	-
Cirrhosis of liver (581)	0	0	0	1	1	0	6.5	10.5	-
Acute nephritis (590, 591)	0	0	0	1	1	0	6.5	10.5	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	13.0	-	34.4
Infections of kidney (600)	0	0	0	2	2	0	13.0	21.0	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	0	1	32.0	-	62.1
Congenital malformations (750-759)	1	0	1	2	1	1	13.0	10.5	17.2
Diseases of early infancy (760-776)	3	1	2	9	2	7	28.8	13.2	43.5
Accidents (800-962)	18	14	4	17	14	3	110.7	146.9	51.5
Motor vehicle accidents (810-835)	9	7	2	10	8	2	65.1	83.9	34.4
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	1	0	1	6.5	-	17.2
Ill-defined (780-795)	13	3	10	13	3	10	84.7	31.5	171.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
CLAY COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	268	111	157	334	143	191	18.8	16.2	21.4
Fetal deaths	4	1	3	9	3	6	26.9	21.0	31.4
Total deaths	164	86	78	176	87	89	9.9	9.9	10.0
Deaths under 1 year	8	1	7	9	3	6	26.9	21.0	31.4
Deaths under 28 days	3	0	3	5	2	3	15.0	14.0	15.7
<b>SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	29	18	11	31	16	15	174.7	181.6	168.0
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	5.6	-	11.2
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	3	1	3	2	1	16.9	22.7	11.2
Anemias (290-293)	1	0	1	1	0	1	5.6	-	11.2
Intracranial lesions (330-334)	20	8	12	19	6	13	107.1	68.1	145.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	60	36	24	64	40	24	360.8	454.0	268.8
Other hypertensive disease (444-447)	1	1	0	2	1	1	11.3	11.4	11.2
General arteriosclerosis (450)	4	2	2	3	1	2	16.9	11.4	22.4
Other circulatory diseases (451-468)	1	0	1	2	1	1	11.3	11.4	11.2
#Influenza & pneumonia (480-493)	3	1	2	4	2	2	22.5	22.7	22.4
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	0	1	1	0	1	5.6	-	11.2
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	0	1	1	0	1	5.6	-	11.2
Acute nephritis (590, 591)	0	0	0	1	0	1	5.6	-	11.2
Chronic nephritis, etc. (592-594)	2	2	0	1	1	0	5.6	11.4	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	5.6	-	11.2
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	5.6	11.4	-
Diseases of early infancy (760-776)	1	0	1	3	2	1	9.0	14.0	5.2
Accidents (800-962)	14	6	8	14	5	9	78.9	56.8	100.8
Motor vehicle accidents (810-835)	9	4	5	9	3	6	50.7	34.0	67.2
Suicide (963, 970-979)	4	3	1	4	3	1	22.5	34.0	11.2
Homicide (964, 980-985)	2	1	1	1	1	0	5.6	11.4	-
Ill-defined (780-795)	9	2	7	10	3	7	56.4	34.0	78.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 COAHOMA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,057	313	744	1,029	246	783	21.6	16.5	24.0
Fetal deaths	52	4	48	52	6	46	50.5	24.4	58.7
Total deaths	551	192	359	518	151	367	10.9	10.1	11.2
Deaths under 1 year	44	7	37	47	8	39	45.7	32.5	49.8
Deaths under 28 days	27	6	21	31	7	24	30.1	28.4	30.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	0	2	1	0	1	2.1	-	3.1
Syphilis (020-029)	0	0	0	1	0	1	2.1	-	3.1
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	1	0	1	2.1	-	3.1
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	65	30	35	59	23	36	124.1	154.6	110.2
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	2.1	-	3.1
Asthma (241)	2	1	1	2	1	1	4.2	6.7	3.1
Diabetes mellitus (260)	8	4	4	6	3	3	12.6	20.2	9.2
Anemias (290-293)	2	0	2	3	0	3	6.3	-	9.2
Intracranial lesions (330-334)	86	34	52	75	15	60	157.7	100.8	183.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	179	77	102	177	69	108	372.2	463.7	330.6
Other hypertensive disease (444-447)	6	0	6	6	0	6	12.6	-	18.4
General arteriosclerosis (450)	3	1	2	4	2	2	8.4	13.4	6.1
Other circulatory diseases (451-468)	8	2	6	9	3	6	18.9	20.2	18.4
#Influenza & pneumonia (480-493)	15	9	6	10	4	6	21.0	26.9	18.4
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	12	5	7	10	3	7	21.0	20.2	21.4
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	2.1	6.7	-
Hernia & intestinal obstruction (560, 561, 570)	3	2	1	3	2	1	6.3	13.4	3.1
#Gastritis, etc. (543, 571, 572)	3	0	3	3	0	3	6.3	-	9.2
Cirrhosis of liver (581)	3	1	2	3	1	2	6.3	6.7	6.1
Acute nephritis (590, 591)	2	1	1	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	1	5	7	1	6	14.7	6.7	18.4
Infections of kidney (600)	3	1	2	4	1	3	8.4	6.7	9.2
Hyperplasia of prostate (610)	4	0	4	1	0	1	2.1	-	3.1
Maternal causes (640-689)	1	0	1	1	0	1	9.7	-	12.8
Congenital malformations (750-759)	6	1	5	8	1	7	16.8	6.7	21.4
Diseases of early infancy (760-776)	22	4	18	25	6	19	24.3	24.4	24.3
Accidents (800-962)	40	10	30	32	9	23	67.3	60.5	70.4
Motor vehicle accidents (810-835)	25	4	21	21	4	17	44.2	26.9	52.0
Suicide (963, 970-979)	1	1	0	1	1	0	2.1	6.7	-
Homicide (964, 980-985)	6	0	6	6	0	6	12.6	-	18.4
Ill-defined (780-795)	42	1	41	43	2	41	90.4	13.4	125.5

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
COPIAH COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	636	183	453	491	155	336	17.9	11.0	25.0
Fetal deaths	13	4	9	19	3	16	38.7	19.4	47.6
Total deaths	243	131	112	326	188	138	11.9	13.4	10.3
Deaths under 1 year	13	1	12	24	5	19	48.9	32.2	56.5
Deaths under 28 days	5	1	4	14	5	9	28.5	32.2	26.8
<b>SELECTED CAUSES OF DEATH</b>									
AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	3.6	-	7.4
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	40	25	15	53	33	20	193.2	235.4	149.1
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	3.6	7.1	-
Asthma (241)	0	0	0	1	0	1	3.6	-	7.4
Diabetes mellitus (260)	1	0	1	1	0	1	3.6	-	7.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	12	23	42	18	24	153.1	128.4	179.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	97	67	30	119	85	34	433.8	606.3	253.5
Other hypertensive disease (444-447)	1	0	1	2	1	1	7.3	7.1	7.4
General arteriosclerosis (450)	8	4	4	9	6	3	32.8	42.8	22.4
Other circulatory diseases (451-468)	1	1	0	2	1	1	7.3	7.1	7.4
#Influenza & pneumonia (480-493)	10	2	8	13	3	10	47.4	21.4	74.6
Bronchitis (500-502)	0	0	0	1	1	0	3.6	7.1	-
Other bronchopul. diseases (525-527)	1	1	0	3	2	1	10.9	14.3	7.4
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	3.6	7.1	-
#Gastritis, etc. (543, 571, 572)	3	1	2	5	2	3	18.2	14.3	22.4
Cirrhosis of liver (581)	0	0	0	1	1	0	3.6	7.1	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	2	0	2	7.3	-	14.9
Infections of kidney (600)	4	0	4	4	0	4	14.6	-	29.8
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	3.6	7.1	-
Diseases of early infancy (760-776)	5	1	4	12	5	7	24.4	32.2	20.8
Accidents (800-962)	14	9	5	23	16	7	83.8	114.1	52.2
Motor vehicle accidents (810-835)	5	4	1	9	8	1	32.8	57.1	7.4
Suicide (963, 970-979)	1	1	0	1	1	0	3.6	7.1	-
Homicide (964, 980-985)	1	0	1	3	0	3	10.9	-	22.4
Ill-defined (780-795)	14	4	10	15	5	10	54.7	35.7	74.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
COVINGTON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	164	118	46	287	156	131	19.6	16.0	26.7
Fetal deaths	3	1	2	10	3	7	34.8	19.2	53.4
Total deaths	82	58	24	130	92	38	8.9	9.4	7.7
Deaths under 1 year	4	2	2	11	6	5	38.3	38.5	38.2
Deaths under 28 days	2	2	0	8	6	2	27.9	38.5	15.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	6	3	15	9	6	102.2	92.2	122.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	6.8	10.2	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	5	4	11	5	6	75.0	51.2	122.2
Normeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	28	24	4	42	36	6	286.3	368.8	122.2
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	4	4	0	5	4	1	34.1	41.0	20.4
Other circulatory diseases (451-468)	0	0	0	2	2	0	13.6	20.5	-
#Influenza & pneumonia (480-493)	3	2	1	2	1	1	13.6	10.2	20.4
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	1	0	1	1	0	6.8	10.2	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	6.8	10.2	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	2	0	3	3	0	20.4	30.7	-
Cirrhosis of liver (581)	0	0	0	1	1	0	6.8	10.2	-
Acute nephritis (590, 591)	2	1	1	2	1	1	13.6	10.2	20.4
Chronic nephritis, etc. (592-594)	1	1	0	2	1	1	13.6	10.2	20.4
Infections of kidney (600)	0	0	0	3	3	0	20.4	30.7	-
Hyperplasia of prostate (610)	0	0	0	1	1	0	6.8	10.2	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	6.8	10.2	-
Diseases of early infancy (760-776)	2	2	0	8	6	2	27.9	38.5	15.3
Accidents (800-962)	9	3	6	14	6	8	95.4	61.5	162.9
Motor vehicle accidents (810-835)	7	2	5	11	4	7	75.0	41.0	142.6
Suicide (963, 970-979)	2	2	0	2	2	0	13.6	20.5	-
Homicide (964, 980-985)	0	0	0	2	2	0	13.6	20.5	-
Ill-defined (780-795)	9	5	4	9	4	5	61.3	41.0	101.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
DESOTO COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	288	17	271	694	335	359	18.8	14.6	25.8
Fetal deaths	4	0	4	12	7	5	17.3	20.9	13.9
Total deaths	144	53	91	261	118	143	7.1	5.1	10.2
Deaths under 1 year	16	0	16	28	7	21	40.3	20.9	58.5
Deaths under 28 days	7	0	7	17	5	12	24.5	14.9	33.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	13	5	8	30	15	15	81.3	65.3	107.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	5	2	3	13.6	8.7	21.5
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	5	12	37	18	19	100.3	78.4	136.3
Nonmeningococcal meningitis (340)	0	0	0	1	1	0	2.7	4.4	-
Heart diseases (400-443)	44	23	21	75	47	28	203.3	204.7	200.9
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	3	2	1	8.1	8.7	7.2
Other circulatory diseases (451-468)	1	1	0	4	2	2	10.8	8.7	14.3
#Influenza & pneumonia (480-493)	6	1	5	8	1	7	21.7	4.4	50.2
Bronchitis (500-502)	1	1	0	1	1	0	2.7	4.4	-
Other bronchopul. diseases (525-527)	1	1	0	3	3	0	8.1	13.1	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	2.7	4.4	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	4	0	4	3	0	3	8.1	-	21.5
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	2	1	1	5.4	4.4	7.2
Infections of kidney (600)	0	0	0	2	0	2	5.4	-	14.3
Hyperplasia of prostate (610)	0	0	0	1	0	1	2.7	-	7.2
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	0	3	14	4	10	37.9	17.4	71.7
Diseases of early infancy (760-776)	6	0	6	9	3	6	13.0	9.0	16.7
Accidents (800-962)	18	7	11	26	10	16	70.5	43.6	114.8
Motor vehicle accidents (810-835)	7	6	1	8	4	4	21.7	17.4	28.7
Suicide (963, 970-979)	3	3	0	3	3	0	8.1	13.1	-
Homicide (964, 980-985)	4	1	3	5	0	5	13.6	-	35.9
Ill-defined (780-795)	13	1	12	13	0	13	35.2	-	93.2

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
FORREST COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,373	978	395	1,116	693	423	17.8	14.9	26.2
Fetal deaths	32	17	15	35	11	24	31.4	15.9	56.7
Total deaths	610	427	183	595	370	225	9.5	7.9	13.9
Deaths under 1 year	31	25	6	26	14	12	23.3	20.2	28.4
Deaths under 28 days	27	21	6	22	11	11	19.7	15.9	26.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	4	1	3	4	1	3	6.4	2.1	18.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	1	0	1	1	0	1	1.6	-	6.2
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	81	60	21	86	56	30	137.1	120.2	185.9
Benign & unspec. neoplasms (210-239)	2	0	2	3	1	2	4.8	2.1	12.4
Asthma (241)	2	1	1	2	1	1	3.9	2.1	6.2
Diabetes mellitus (260)	13	7	6	15	7	8	23.9	15.0	49.6
Anemias (290-293)	1	1	0	0	0	0	-	-	-
Intracranial lesions (330-334)	58	45	13	64	45	19	102.0	96.6	117.7
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	1.6	-	6.2
Heart diseases (400-443)	210	152	58	193	131	62	307.8	281.3	384.1
Other hypertensive disease (444-447)	8	2	6	8	2	6	12.8	4.3	37.2
General arteriosclerosis (450)	7	6	1	7	6	1	11.2	12.9	6.2
Other circulatory diseases (451-468)	14	14	0	9	9	0	14.4	19.3	-
#Influenza & pneumonia (480-493)	18	11	7	16	7	9	25.5	15.0	55.8
Bronchitis (500-502)	2	2	0	3	3	0	4.8	6.4	-
Other bronchopul. diseases (525-527)	16	14	2	16	13	3	25.5	27.9	18.6
Stomach & duodenal ulcers (540, 541)	4	4	0	2	2	0	3.2	4.3	-
Hernia & intestinal obstruction (560, 561, 570)	2	2	0	1	1	0	1.6	2.1	-
#Gastritis, etc. (543, 571, 572)	4	3	1	3	2	1	4.8	4.3	6.2
Cirrhosis of liver (581)	2	2	0	2	2	0	3.2	4.3	-
Acute nephritis (590, 591)	1	1	0	1	1	0	1.6	2.1	-
Chronic nephritis, etc. (592-594)	6	2	4	6	1	5	9.6	2.1	31.0
Infections of kidney (600)	5	2	3	7	3	4	11.2	6.4	24.8
Hyperplasia of prostate (610)	4	1	3	3	0	3	4.8	-	18.6
Maternal causes (640-689)	1	0	1	1	0	1	9.0	-	23.6
Congenital malformations (750-759)	3	3	0	3	3	0	4.8	6.4	-
Diseases of early infancy (760-776)	24	18	6	19	9	10	17.0	13.0	23.6
Accidents (800-962)	51	42	9	54	35	19	86.1	75.2	117.7
Motor vehicle accidents (810-835)	23	22	1	24	17	7	38.3	36.5	43.4
Suicide (963, 970-979)	0	0	0	1	1	0	1.6	2.1	-
Homicide (964, 980-985)	12	5	7	11	3	8	17.5	6.4	49.6
Ill-defined (780-795)	36	12	24	35	11	24	55.8	23.6	148.7

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
FRANKLIN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	90	35	55	117	60	57	12.6	10.4	16.3
Fetal deaths	1	1	0	4	2	2	34.2	33.3	35.1
Total deaths	116	65	51	140	85	55	15.1	14.7	15.7
Deaths under 1 year	4	0	4	8	1	7	68.4	16.7	122.8
Deaths under 28 days	1	0	1	3	1	2	25.6	16.7	35.1
<b>SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	17	11	6	21	16	5	226.5	277.3	142.8
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	0	4	6	2	4	64.7	34.7	114.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	9	6	19	13	6	205.0	225.3	171.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	51	31	20	53	34	19	571.7	589.2	542.8
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	4	3	1	3	2	1	32.4	34.7	28.6
Other circulatory diseases (451-468)	2	2	0	2	2	0	21.6	34.7	-
#Influenza & pneumonia (480-493)	2	0	2	5	1	4	53.9	17.3	114.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	1	1	2	1	1	21.6	17.3	28.6
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	3	1	2	3	1	2	32.4	17.3	57.1
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	1	0	2	2	0	21.6	34.7	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	10.8	17.3	-
Diseases of early infancy (760-776)	1	0	1	3	1	2	25.6	16.7	35.1
Accidents (800-962)	10	4	6	13	6	7	140.2	104.0	200.0
Motor vehicle accidents (810-835)	4	2	2	5	3	2	53.9	52.0	57.1
Suicide (963, 970-979)	0	0	0	1	1	0	10.8	17.3	-
Homicide (964, 980-985)	1	0	1	1	0	1	10.8	-	28.6
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
GEORGE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	307	269	38	266	233	33	21.7	21.5	23.4
Fetal deaths	4	4	0	3	3	0	11.3	12.9	-
Total deaths	90	75	15	110	94	16	9.0	8.7	11.3
Deaths under 1 year	8	5	3	10	8	2	37.6	34.3	60.6
Deaths under 28 days	6	5	1	8	7	1	30.1	30.0	30.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	1	1	0	8.2	9.2	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	9	0	13	13	0	106.0	119.8	-
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	8.2	9.2	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	8.2	9.2	-
Anemias (290-293)	0	0	0	2	1	1	16.3	9.2	70.9
Intracranial lesions (330-334)	4	2	2	4	3	1	32.6	27.6	70.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	42	38	4	46	41	5	375.2	377.9	354.6
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	2	2	0	16.3	18.4	-
Other circulatory diseases (451-468)	4	4	0	4	4	0	32.6	36.9	-
#Influenza & pneumonia (480-493)	1	1	0	3	2	1	24.5	18.4	70.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	4	3	1	4	3	1	32.6	27.6	70.9
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	8.2	9.2	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	8.2	9.2	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	1	1	0	8.2	9.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	8.2	9.2	-
Infections of kidney (600)	1	1	0	1	1	0	8.2	9.2	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	2	2	0	16.3	18.4	-
Diseases of early infancy (760-776)	4	4	0	5	5	0	18.8	21.4	-
Accidents (800-962)	4	3	1	6	4	2	48.9	36.9	141.8
Motor vehicle accidents (810-835)	1	1	0	3	2	1	24.5	18.4	70.9
Suicide (963, 970-979)	0	0	0	1	1	0	8.2	9.2	-
Homicide (964, 980-985)	4	2	2	2	1	1	16.3	9.2	70.9
Ill-defined (780-795)	1	0	1	2	1	1	16.3	9.2	70.9

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
GREENE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	59	28	31	174	117	57	20.8	17.8	31.7
Fetal deaths	4	1	3	8	4	4	46.0	34.2	70.2
Total deaths	65	50	15	83	65	18	9.9	9.9	10.0
Deaths under 1 year	4	0	4	7	0	7	40.2	-	122.8
Deaths under 28 days	3	0	3	3	0	3	17.2	-	52.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	5	5	0	7	7	0	83.5	106.4	-
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	11.9	15.2	-
Diabetes mellitus (260)	2	2	0	2	2	0	23.9	30.4	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	5	5	0	8	7	1	95.5	6.4	55.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	25	23	2	37	33	4	441.5	501.5	222.2
Other hypertensive disease (444-447)	2	1	1	2	1	1	23.9	15.2	55.6
General arteriosclerosis (450)	2	2	0	2	2	0	23.9	30.4	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	5	4	1	8	6	2	95.5	91.2	111.1
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	11.9	15.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	35.8	-	166.7
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	4	1	3	4	1	3	47.7	15.2	166.7
Diseases of early infancy (760-776)	1	0	1	1	0	1	5.7	-	17.5
Accidents (800-962)	7	3	4	4	2	2	47.7	30.4	111.1
Motor vehicle accidents (810-835)	5	2	3	2	1	1	23.9	15.2	55.6
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	1	0	0	0	0	-	-	-
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
GRENADA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	384	131	253	412	156	256	19.4	13.7	26.0
Fetal deaths	19	3	16	20	4	16	48.5	25.6	62.5
Total deaths	237	121	116	224	101	123	10.6	8.9	12.5
Deaths under 1 year	16	1	15	17	2	15	41.3	12.8	58.6
Deaths under 28 days	9	0	9	11	1	10	26.7	6.4	39.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	4.7	-	10.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	30	20	10	27	16	11	127.2	140.6	111.7
Benign & unspec. neoplasms (210-239)	2	1	1	2	1	1	9.4	8.8	10.2
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	2	0	1	1	0	4.7	8.8	-
Anemias (290-293)	1	1	0	1	1	0	4.7	8.8	-
Intracranial lesions (330-334)	62	27	35	57	23	34	268.5	202.1	345.2
Nonmeningococcal meningitis (340)	1	1	0	0	0	0	-	-	-
Heart diseases (400-443)	54	33	21	54	28	26	254.4	246.0	264.0
Other hypertensive disease (444-447)	6	1	5	7	1	6	33.0	8.8	60.9
General arteriosclerosis (450)	2	2	0	2	1	1	9.4	8.8	10.2
Other circulatory diseases (451-468)	1	0	1	2	1	1	9.4	8.8	10.2
#Influenza & pneumonia (480-493)	7	5	2	4	2	2	18.8	17.6	20.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	3	0	2	2	0	9.4	17.6	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	4.7	-	10.2
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	4.7	8.8	-
Cirrhosis of liver (581)	1	1	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	1	0	4.7	8.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	24.3	-	39.1
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	10	0	10	12	1	11	29.1	6.4	43.0
Accidents (800-962)	25	15	10	23	13	10	108.3	114.2	101.5
Motor vehicle accidents (810-835)	9	9	0	10	9	1	47.1	79.1	10.2
Suicide (963, 970-979)	3	1	2	2	1	1	9.4	8.8	10.2
Homicide (964, 980-985)	4	1	3	4	1	3	18.8	8.8	30.4
Ill-defined (780-795)	16	5	11	13	4	9	61.2	35.1	91.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
HANCOCK COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	183	140	43	329	274	55	16.0	15.5	19.1
Fetal deaths	4	4	0	6	6	0	18.2	21.9	-
Total deaths	162	135	27	180	151	29	8.7	8.5	10.1
Deaths under 1 year	8	6	2	9	6	3	27.4	21.9	54.5
Deaths under 28 days	4	3	1	5	4	1	15.2	14.6	18.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	15	3	31	26	5	150.6	146.9	173.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	2	2	0	2	2	0	9.7	11.3	-
Diabetes mellitus (260)	4	4	0	4	4	0	19.4	22.6	-
Anemias (290-293)	1	0	1	1	0	1	4.8	-	34.7
Intracranial lesions (330-334)	12	9	3	13	10	3	63.2	56.5	104.2
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	4.8	-	34.7
Heart diseases (400-443)	50	44	6	61	55	6	296.4	310.7	208.3
Other hypertensive disease (444-447)	0	0	0	1	1	0	4.8	5.6	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	2	2	0	1	1	0	4.8	5.6	-
#Influenza & pneumonia (480-493)	3	3	0	4	3	1	19.4	16.9	34.7
Bronchitis (500-502)	1	1	0	1	1	0	4.8	5.6	-
Other bronchopul. diseases (525-527)	1	1	0	2	2	0	9.7	11.3	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	4.8	5.6	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	4.8	5.6	-
#Castritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	6	5	1	6	5	1	29.2	28.2	34.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	2	2	0	2	2	0	9.7	11.3	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	3	3	0	14.6	16.9	-
Diseases of early infancy (760-776)	3	2	1	3	2	1	9.1	7.3	18.2
Accidents (800-962)	30	21	9	22	14	8	106.9	79.1	277.8
Motor vehicle accidents (810-835)	19	13	6	13	7	6	63.2	39.5	208.3
Suicide (963, 970-979)	2	2	0	3	3	0	14.6	16.9	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	16	14	2	12	10	2	58.3	56.5	69.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
HARRISON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	3,468	2,720	748	3,084	2,379	705	22.0	19.9	34.1
Fetal deaths	244	189	55	207	155	52	67.1	65.2	73.8
Total deaths	1,226	982	244	1,010	795	215	7.2	6.6	10.4
Deaths under 1 year	91	57	34	82	49	33	26.6	20.6	46.8
Deaths under 28 days	62	37	25	55	30	25	17.8	12.6	35.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	0.7	-	4.8
Syphilis (020-029)	1	0	1	1	1	0	0.7	0.8	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	1	1	0	0.7	0.8	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	213	180	33	161	137	24	114.7	114.5	116.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	3	2	1	2	1	1	1.4	0.8	4.8
Diabetes mellitus (260)	11	9	2	9	6	3	6.4	5.0	14.5
Anemias (290-293)	1	1	0	1	1	0	0.7	0.8	-
Intracranial lesions (330-334)	103	77	26	83	60	23	59.1	50.1	111.3
Nonmeningococcal meningitis (340)	3	0	3	4	1	3	2.8	0.8	14.5
Heart diseases (400-443)	409	339	70	341	277	64	242.9	231.4	309.6
Other hypertensive disease (444-447)	10	5	5	7	4	3	5.0	3.3	14.5
General arteriosclerosis (450)	10	7	3	9	7	2	6.4	5.8	9.7
Other circulatory diseases (451-468)	17	15	2	17	15	2	12.1	12.5	9.7
#Influenza & pneumonia (480-493)	40	35	5	24	20	4	17.1	16.7	19.4
Bronchitis (500-502)	7	7	0	3	3	0	2.1	2.5	-
Other bronchopul. diseases (525-527)	24	23	1	17	16	1	12.1	13.4	4.8
Stomach & duodenal ulcers (540, 541)	3	2	1	3	2	1	2.1	1.7	4.8
Hernia & intestinal obstruction (560, 561, 570)	5	5	0	3	3	0	2.1	2.5	-
#Gastritis, etc. (543, 571, 572)	4	4	0	5	5	0	3.6	4.2	-
Cirrhosis of liver (581)	16	13	3	11	9	2	7.8	7.5	9.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	0.7	0.8	-
Infections of kidney (600)	7	7	0	7	7	0	5.0	5.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	24	15	9	21	12	9	15.0	10.0	43.5
Diseases of early infancy (760-776)	45	29	16	40	25	15	13.0	10.5	21.3
Accidents (800-962)	114	93	21	101	82	19	72.0	68.5	91.9
Motor vehicle accidents (810-835)	65	58	7	50	46	4	35.6	38.4	19.4
Suicide (963, 970-979)	13	13	0	11	11	0	7.8	9.2	-
Homicide (964, 980-985)	10	2	8	10	4	6	7.1	3.3	29.0
Ill-defined (780-795)	98	71	27	86	59	27	61.3	49.3	130.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
HINDS COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	5,359	2,621	2,738	4,347	2,001	2,346	19.4	15.9	24.0
Fetal deaths	561	292	269	401	209	192	92.2	104.4	81.8
Total deaths	2,696	1,524	1,172	1,722	944	778	7.7	7.5	8.0
Deaths under 1 year	191	72	119	115	36	79	26.4	18.0	33.7
Deaths under 28 days	137	57	80	84	30	54	19.3	15.0	23.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	18	7	11	14	5	9	6.3	4.0	9.2
Syphilis (020-029)	4	3	1	2	2	0	1.0	1.6	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	1	0	1	0.4	-	1.0
Meningococcal infections (057)	2	1	1	1	0	1	0.4	-	1.0
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	495	322	173	282	196	86	125.9	155.4	87.9
Benign & unspec. neoplasms (210-239)	8	4	4	2	1	1	1.0	0.8	1.0
Asthma (241)	8	3	5	2	1	1	1.0	0.8	1.0
Diabetes mellitus (260)	50	22	28	28	10	18	12.5	7.9	18.4
Anemias (290-293)	7	1	6	3	0	3	1.3	-	3.1
Intracranial lesions (330-334)	312	161	151	201	85	116	89.8	67.4	118.6
Nonmeningococcal meningitis (340)	14	3	11	5	0	5	2.2	-	5.1
Heart diseases (400-443)	724	475	249	516	335	181	230.4	265.7	185.0
Other hypertensive disease (444-447)	27	9	18	17	6	11	7.6	4.8	11.2
General arteriosclerosis (450)	36	22	14	26	12	14	11.6	9.5	14.3
Other circulatory diseases (451-468)	59	34	25	33	19	14	14.7	15.1	14.3
Influenza & pneumonia (480-493)	60	35	25	46	28	18	20.5	22.2	18.4
Bronchitis (500-502)	16	14	2	9	8	1	4.0	6.3	1.0
Other bronchopul. diseases (525-527)	34	19	15	14	10	4	6.2	7.9	4.1
Stomach & duodenal ulcers (540, 541)	23	11	12	12	4	8	5.4	3.2	8.2
Hernia & intestinal obstruction (560, 561, 570)	22	13	9	11	7	4	4.9	5.6	4.1
#Gastritis, etc. (543, 571, 572)	17	11	6	12	7	5	5.4	5.6	5.1
Cirrhosis of liver (581)	29	18	11	17	9	8	7.6	7.1	8.2
Acute nephritis (590, 591)	4	0	4	3	0	3	1.3	-	3.1
Chronic nephritis, etc. (592-594)	18	3	15	4	1	3	1.8	0.8	3.1
Infections of kidney (600)	17	11	6	9	6	3	4.0	4.8	3.1
Hyperplasia of prostate (610)	4	2	2	0	0	0	-	-	-
Maternal causes (640-689)	6	1	5	4	0	4	9.2	-	17.0
Congenital malformations (750-759)	47	29	18	16	13	3	7.1	10.3	3.1
Diseases of early infancy (760-776)	106	40	66	69	20	49	15.9	10.0	20.9
Accidents (800-902)	214	123	91	133	83	50	59.4	65.8	51.1
Motor vehicle accidents (810-835)	105	69	36	65	47	18	29.0	37.3	18.4
Suicide (963, 970-979)	18	17	1	16	14	2	7.1	11.1	2.0
Homicide (964, 980-985)	46	9	37	35	5	30	15.6	4.0	30.7
Ill-defined (780-795)	112	29	83	103	20	83	46.0	15.9	84.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
HOLMES COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	539	73	466	552	83	469	19.4	9.9	23.4
Fetal deaths	25	3	22	28	3	25	50.7	36.1	53.3
Total deaths	270	64	206	337	100	237	11.8	11.9	11.8
Deaths under 1 year	22	3	19	25	3	22	45.3	36.1	46.9
Deaths under 28 days	11	2	9	14	2	12	25.4	24.1	25.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	27	9	18	37	18	19	129.9	214.3	94.6
Benign & unspec. neoplasms (210-239)	2	1	1	2	1	1	7.0	11.9	5.0
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	5	1	4	8	1	7	28.1	11.9	34.9
Anemias (290-293)	3	1	2	3	1	2	10.5	11.9	10.0
Intracranial lesions (330-334)	38	10	28	50	15	35	175.6	178.6	174.3
Nonmeningococcal meningitis (340)	2	1	1	1	0	1	3.5	-	5.0
Heart diseases (400-443)	75	22	53	92	31	61	323.0	369.0	303.8
Other hypertensive disease (444-447)	6	1	5	6	1	5	21.1	11.9	24.9
General arteriosclerosis (450)	3	0	3	4	1	3	14.0	11.9	14.9
Other circulatory diseases (451-468)	0	0	0	2	1	1	7.0	11.9	5.0
#Influenza & pneumonia (480-493)	18	4	14	16	3	13	56.2	35.7	64.7
Bronchitis (500-502)	1	0	1	1	0	1	3.5	-	5.0
Other bronchopul. diseases (525-527)	0	0	0	1	0	1	3.5	-	5.0
Stomach & duodenal ulcers (540, 541)	0	0	0	2	0	2	7.0	-	10.0
Hernia & intestinal obstruction (560, 561, 570)	3	1	2	6	4	2	21.1	47.6	10.0
#Gastritis, etc. (543, 571, 572)	1	0	1	2	1	1	7.0	11.9	5.0
Cirrhosis of liver (581)	0	0	0	1	0	1	3.5	-	5.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	1	4	5	1	4	17.6	11.9	19.9
Infections of kidney (600)	1	1	0	3	3	0	10.5	35.7	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.5	-	5.0
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	3	1	2	10.5	11.9	10.0
Diseases of early infancy (760-776)	12	2	10	14	2	12	25.4	24.1	25.6
Accidents (800-962)	14	3	11	18	5	13	63.2	59.5	64.7
Motor vehicle accidents (810-835)	8	2	6	6	2	4	21.1	23.8	19.9
Suicide (963, 970-979)	2	1	1	3	1	2	10.5	11.9	10.0
Homicide (964, 980-985)	3	0	3	4	1	3	14.0	11.9	14.9
Ill-defined (780-795)	39	3	36	39	4	35	136.9	47.6	174.3

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
HUMPHREYS COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	430	56	374	409	88	321	21.6	15.8	23.9
Fetal deaths	13	3	10	23	4	19	56.2	45.4	59.2
Total deaths	156	40	116	190	58	132	10.0	10.4	9.8
Deaths under 1 year	19	2	17	17	2	15	41.6	22.7	46.7
Deaths under 28 days	14	2	12	13	2	11	31.8	22.7	34.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	14	5	9	20	9	11	105.5	162.2	82.0
Benign & unspec. neoplasms (210-239)	0	0	0	1	0	1	5.3	-	7.4
Asthma (241)	1	0	1	1	0	1	5.3	-	7.4
Diabetes mellitus (260)	3	0	3	3	0	3	15.8	-	22.4
Anemias (290-293)	1	0	1	1	0	1	5.3	-	7.4
Intracranial lesions (330-334)	23	5	18	29	7	22	153.0	126.1	164.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	57	18	39	71	25	46	374.5	450.4	343.0
Other hypertensive disease (444-447)	1	0	1	2	0	2	10.5	-	14.9
General arteriosclerosis (450)	1	0	1	2	0	2	10.5	-	14.9
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	3	0	3	3	0	3	15.8	-	22.4
Bronchitis (500-502)	1	0	1	1	0	1	5.3	-	7.4
Other bronchopul. diseases (525-527)	1	0	1	2	1	1	10.5	18.0	7.4
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	5.3	18.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	0	6	8	0	8	42.2	-	59.6
Infections of kidney (600)	2	1	1	2	1	1	10.5	18.0	7.4
Hyperplasia of prostate (610)	1	0	1	1	0	1	5.3	-	7.4
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	1	1	0	5.3	18.0	-
Diseases of early infancy (760-776)	10	1	9	10	1	9	24.4	11.4	28.0
Accidents (800-962)	12	6	6	15	8	7	79.1	144.1	52.2
Motor vehicle accidents (810-835)	6	5	1	7	6	1	36.9	108.1	7.4
Suicide (963, 970-979)	2	1	1	2	1	1	10.5	18.0	7.4
Homicide (964, 980-985)	2	0	2	2	0	2	10.5	-	14.9
Ill-defined (780-795)	13	1	12	10	1	9	52.7	18.0	67.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
ISSAQENA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	13	0	13	51	9	42	18.3	9.6	22.8
Fetal deaths	0	0	0	4	2	2	78.4	222.2	47.6
Total deaths	20	4	16	42	14	28	15.1	14.9	15.2
Deaths under 1 year	2	0	2	6	1	5	117.6	111.1	119.0
Deaths under 28 days	0	0	0	3	1	2	58.8	111.1	47.6
<b>SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	1	0	1	1	0	1	36.0	-	54.3
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	0	0	0	3	0	3	107.9	-	163.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	0	1	36.0	-	54.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	4	0	4	6	2	4	215.8	212.8	217.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	7	3	4	10	5	5	359.7	531.9	271.7
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	1	1	0	1	0	1	36.0	-	54.3
#Influenza & pneumonia (480-493)	0	0	0	3	1	2	107.9	106.4	108.7
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	2	1	1	71.9	106.4	54.3
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	36.0	-	54.3
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	0	1	36.0	-	54.3
Diseases of early infancy (760-776)	1	0	1	2	1	1	39.2	111.1	23.8
Accidents (800-962)	1	0	1	5	4	1	179.8	425.5	54.3
Motor vehicle accidents (810-835)	1	0	1	3	3	0	107.9	319.1	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	1	0	1	36.0	-	54.3
Ill-defined (780-795)	4	0	4	5	0	5	179.8	-	271.7

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
ITAWAMBA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Total	White	Non-white
				Total	White	Non-white			
Live births	38	29	9	207	192	15	12.2	12.0	15.0
Fetal deaths	0	0	0	5	5	0	24.2	26.0	-
Total deaths	77	68	9	146	137	9	8.6	8.6	9.0
Deaths under 1 year	1	1	0	3	3	0	14.5	15.6	-
Deaths under 28 days	0	0	0	1	1	0	4.8	5.2	-
<b>SELECTED CAUSES OF DEATH</b>									
<b>AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	5.9	6.3	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	16	14	2	25	24	1	147.7	150.6	100.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	2	1	1	11.8	6.3	100.0
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	18	2	27	25	2	159.5	156.9	200.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	26	22	4	52	47	5	307.1	295.0	500.0
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	2	2	0	4	4	0	23.6	25.1	-
Other circulatory diseases (451-468)	1	1	0	2	2	0	11.8	12.6	-
#Influenza & pneumonia (480-493)	1	1	0	4	4	0	23.6	25.1	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	1	0	2	2	0	11.8	12.6	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	5.9	6.3	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	5.9	6.3	-
Cirrhosis of liver (581)	0	0	0	2	2	0	11.8	12.6	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	5.9	6.3	-
Infections of kidney (600)	1	1	0	1	1	0	5.9	6.3	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	1	0	48.3	52.1	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	1	1	0	4.8	5.2	-
Accidents (800-962)	4	4	0	14	14	0	82.7	87.9	-
Motor vehicle accidents (810-835)	2	2	0	6	6	0	35.4	37.7	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
JACKSON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,400	1,015	385	1,765	1,346	419	23.5	21.4	34.3
Fetal deaths	29	14	15	59	42	17	33.4	31.2	40.6
Total deaths	378	293	85	454	355	99	6.0	5.6	8.1
Deaths under 1 year	31	18	13	43	28	15	24.4	20.8	35.8
Deaths under 28 days	27	17	10	35	24	11	19.8	17.8	26.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	2	2	0	2.7	3.2	-
Syphilis (020-029)	1	0	1	1	0	1	1.3	-	8.2
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	57	47	10	75	63	12	100.0	100.3	98.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	1.3	1.6	-
Diabetes mellitus (260)	3	3	0	5	5	0	6.7	8.0	-
Anemias (290-293)	2	1	1	0	0	0	-	-	-
Intracranial lesions (330-334)	37	23	14	41	26	15	54.6	41.4	122.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	108	93	15	129	109	20	172.0	173.6	163.8
Other hypertensive disease (444-447)	2	1	1	2	1	1	2.7	1.6	8.2
General arteriosclerosis (450)	6	6	0	4	4	0	5.3	6.4	-
Other circulatory diseases (451-468)	6	6	0	7	7	0	9.3	11.1	-
#Influenza & pneumonia (480-493)	8	4	4	9	5	4	12.0	8.0	32.8
Bronchitis (500-502)	2	2	0	3	3	0	4.0	4.8	-
Other bronchopul. diseases (525-527)	4	4	0	5	5	0	6.7	8.0	-
Stomach & duodenal ulcers (540, 541)	2	2	0	2	2	0	2.7	3.2	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	1.3	1.6	-
#Gastritis, etc. (543, 571, 572)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (581)	2	2	0	3	3	0	4.0	4.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	4.0	-	24.6
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	5.7	-	23.9
Congenital malformations (750-759)	3	3	0	9	8	1	12.0	12.7	8.2
Diseases of early infancy (760-776)	23	13	10	30	18	12	17.0	13.4	28.6
Accidents (800-962)	52	41	11	60	50	10	80.0	79.6	81.9
Motor vehicle accidents (810-835)	25	22	3	35	31	4	46.7	49.4	32.8
Suicide (963, 970-979)	3	3	0	4	4	0	5.3	6.4	-
Homicide (964, 980-985)	5	4	1	10	5	5	13.3	8.0	41.0
Ill-defined (780-795)	32	20	12	28	17	11	37.3	27.1	90.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
JASPER COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	198	127	71	330	121	209	20.4	14.5	26.7
Fetal deaths	9	4	5	13	5	8	39.4	41.3	38.3
Total deaths	165	86	79	202	99	103	12.5	11.9	13.2
Deaths under 1 year	14	3	11	18	2	16	54.5	16.5	76.6
Deaths under 28 days	8	2	6	11	1	10	33.3	8.3	47.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	-	6.2	-	12.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	16	12	4	15	9	6	92.8	108.0	76.6
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	6.2	12.0	-
Asthma (241)	0	0	0	1	0	1	6.2	-	12.8
Diabetes mellitus (260)	5	3	2	10	5	5	61.9	60.0	63.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	24	11	13	25	13	12	154.7	156.1	153.2
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	71	39	32	79	45	34	488.9	540.2	434.2
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	3	2	1	18.6	24.0	12.8
Other circulatory diseases (451-468)	4	3	1	4	2	2	24.8	24.0	25.5
#Influenza & pneumonia (480-493)	6	1	5	9	3	6	55.7	36.0	76.6
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	4	3	1	4	3	1	24.8	36.0	12.8
Cirrhosis of liver (581)	0	0	0	1	1	0	6.2	12.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	1	2	3	1	2	18.6	12.0	25.5
Infections of kidney (600)	2	0	2	3	1	2	18.6	12.0	25.5
Hyperplasia of prostate (610)	0	0	0	2	0	2	12.4	-	25.5
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	6	1	5	9	1	8	27.3	8.3	38.3
Accidents (800-962)	8	4	4	14	8	6	86.6	96.0	76.6
Motor vehicle accidents (810-835)	3	3	0	6	5	1	37.1	60.0	12.8
Suicide (963, 970-979)	1	1	0	1	1	0	6.2	12.0	-
Homicide (964, 980-985)	2	1	1	5	1	4	30.9	12.0	51.1
Ill-defined (780-795)	5	1	4	6	1	5	37.1	12.0	63.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
JEFFERSON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	69	12	57	194	26	168	16.8	9.0	19.5
Fetal deaths	2	0	2	5	1	4	25.8	38.5	23.8
Total deaths	82	15	67	131	37	94	11.4	12.8	10.9
Deaths under 1 year	2	0	2	7	0	7	36.1	-	41.7
Deaths under 28 days	1	0	1	3	0	3	15.5	-	17.8
<b>SELECTED CAUSES OF DEATH</b>									
AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	13	5	8	22	8	14	190.8	275.9	162.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	0	2	3	0	3	26.0	-	34.8
Anemias (290-293)	0	0	0	1	0	1	8.7	-	11.6
Intracranial lesions (330-334)	11	0	11	13	1	12	112.7	34.5	139.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	33	6	27	49	16	33	425.0	551.7	382.4
Other hypertensive disease (444-447)	0	0	0	1	1	0	8.7	34.5	-
General arteriosclerosis (450)	1	0	1	4	2	2	34.7	69.0	23.2
Other circulatory diseases (451-468)	2	1	1	2	1	1	17.3	34.5	11.6
#Influenza & pneumonia (480-493)	1	0	1	4	1	3	34.7	34.5	34.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	1	0	1	8.7	-	11.6
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	0	2	17.3	-	23.2
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	17.3	-	23.2
Infections of kidney (600)	2	0	2	2	0	2	17.3	-	23.2
Hyperplasia of prostate (610)	1	0	1	1	0	1	8.7	-	11.6
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	1	0	1	3	0	3	15.5	-	17.8
Accidents (800-962)	4	2	2	9	4	5	78.0	137.9	57.9
Motor vehicle accidents (810-835)	2	1	1	5	2	3	43.4	69.0	34.8
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	2	0	2	17.3	-	23.2
Ill-defined (780-795)	6	0	6	6	0	6	52.0	-	69.5

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
JEFF DAVIS COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	198	105	93	263	103	160	19.0	16.1	21.5
Fetal deaths	5	0	5	8	0	8	30.4	-	50.0
Total deaths	135	69	66	143	72	71	10.3	11.2	9.5
Deaths under 1 year	11	2	9	16	4	12	60.8	38.8	75.0
Deaths under 28 days	8	2	6	12	4	8	45.6	38.8	50.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	23	14	9	27	15	12	194.9	234.0	161.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	2	1	1	14.4	15.6	13.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	20	8	12	20	9	11	144.4	140.4	147.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	45	27	18	38	20	18	274.4	312.0	241.9
Other hypertensive disease (444-447)	1	1	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	3	2	1	4	3	1	28.9	46.8	13.4
#Influenza & pneumonia (480-493)	6	2	4	6	2	4	43.3	31.2	53.8
Bronchitis (500-502)	1	1	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	3	3	0	21.7	46.8	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	7.2	15.6	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	0	1	1	0	1	7.2	-	13.4
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	2	0	2	2	0	2	76.0	-	125.0
Congenital malformations (750-759)	2	0	2	1	0	1	7.2	-	13.4
Diseases of early infancy (760-776)	6	2	4	11	4	7	41.8	38.8	43.8
Accidents (800-962)	5	3	2	9	6	3	65.0	93.6	40.3
Motor vehicle accidents (810-835)	5	3	2	7	5	2	50.5	78.0	26.9
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	2	0	2	2	0	2	14.4	-	26.9
Ill-defined (780-795)	9	1	8	9	2	7	65.0	31.2	94.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
JONES COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,779	754	1,025	943	609	334	15.8	13.8	21.6
Fetal deaths	84	21	63	40	16	24	42.4	26.3	71.8
Total deaths	716	466	250	601	439	162	10.1	9.9	10.5
Deaths under 1 year	67	21	46	28	17	11	29.7	27.9	32.9
Deaths under 28 days	49	15	34	14	10	4	14.8	16.4	12.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	96	68	28	92	71	21	154.0	160.3	135.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	0	0	0	-	-	-
Diabetes mellitus (260)	13	3	10	9	3	6	15.1	6.8	38.8
Anemias (290-293)	2	1	1	1	0	1	1.7	-	6.5
Intracranial lesions (330-334)	80	59	21	71	59	12	118.9	133.2	77.7
Nonmeningococcal meningitis (340)	5	2	3	4	2	2	6.7	4.5	12.9
Heart diseases (400-443)	211	159	52	193	145	48	323.1	327.5	310.7
Other hypertensive disease (444-447)	4	0	4	4	1	3	6.7	2.2	19.4
General arteriosclerosis (450)	7	5	2	7	5	2	11.7	11.3	12.9
Other circulatory diseases (451-468)	4	2	2	5	4	1	8.4	9.0	6.5
#Influenza & pneumonia (480-493)	25	17	8	18	14	4	30.1	31.6	25.9
Bronchitis (500-502)	1	1	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	5	4	1	4	4	0	6.7	9.0	-
Stomach & duodenal ulcers (540, 541)	5	5	0	5	5	0	8.4	11.3	-
Hernia & intestinal obstruction (560, 561, 570)	5	3	2	5	3	2	8.4	6.8	12.9
#Gastritis, etc. (543, 571, 572)	7	3	4	6	3	3	10.0	6.8	19.4
Cirrhosis of liver (541)	5	2	3	3	1	2	5.0	2.2	12.9
Acute nephritis (590, 591)	3	0	3	3	0	3	5.0	-	19.4
Chronic nephritis, etc. (592-594)	13	7	6	10	6	4	16.7	13.6	25.9
Infections of kidney (600)	9	6	3	5	3	2	8.4	6.8	12.9
Hyperplasia of prostate (610)	3	2	1	2	2	0	3.3	4.5	-
Maternal causes (640-689)	1	0	1	0	0	0	-	-	-
Congenital malformations (750-759)	6	6	0	7	7	0	11.7	15.8	-
Diseases of early infancy (760-776)	45	13	32	10	6	4	10.6	9.8	12.0
Accidents (800-962)	62	46	16	48	42	6	80.4	94.8	38.8
Motor vehicle accidents (810-835)	38	27	11	31	27	4	51.9	61.0	25.9
Suicide (963, 970-979)	6	6	0	6	6	0	10.0	13.6	-
Homicide (964, 980-985)	13	5	8	8	3	5	13.4	6.8	32.4
Ill-defined (780-795)	53	24	29	52	26	26	87.0	58.7	168.3

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
KEMPER COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	109	15	94	191	32	159	16.6	6.0	25.7
Fetal deaths	7	0	7	11	1	10	57.6	31.2	62.9
Total deaths	78	35	43	114	58	56	9.9	10.9	9.0
Deaths under 1 year	0	0	0	2	0	2	10.5	-	12.6
Deaths under 28 days	0	0	0	2	0	2	10.5	-	12.6
<b>SELECTED CAUSES OF DEATH</b>									
AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	5	4	14	7	7	121.7	131.8	113.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	1	0	2	2	0	17.4	37.7	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	6	1	5	12	3	9	104.3	56.5	145.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	40	18	22	46	25	21	400.0	470.8	339.2
Other hypertensive disease (444-447)	0	0	0	1	0	1	8.7	-	16.2
General arteriosclerosis (450)	0	0	0	3	2	1	26.1	37.7	16.2
Other circulatory diseases (451-468)	1	0	1	2	0	2	17.4	-	32.3
#Influenza & pneumonia (480-493)	3	2	1	3	2	1	26.1	37.7	16.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	3	3	0	26.1	56.5	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	8.7	18.8	-
Cirrhosis of liver (581)	0	0	0	1	1	0	8.7	18.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	1	0	1	1	0	8.7	18.8	-
Hyperplasia of prostate (610)	0	0	0	2	2	0	17.4	37.7	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	1	0	1	5.2	-	6.3
Accidents (800-962)	2	1	1	6	5	1	52.2	94.2	16.2
Motor vehicle accidents (810-835)	1	0	1	3	2	1	26.1	37.7	16.2
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	2	0	2	17.4	-	32.3
Ill-defined (780-795)	12	4	8	14	4	10	121.7	75.3	161.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LAFAYETTE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	533	320	213	467	272	195	18.7	15.7	25.5
Fetal deaths	14	7	7	10	4	6	21.4	14.7	30.8
Total deaths	198	139	59	196	129	67	7.9	7.5	8.8
Deaths under 1 year	16	2	14	18	4	14	38.5	14.7	71.8
Deaths under 28 days	11	1	10	13	3	10	27.8	11.0	51.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	1	0	1	4.0	-	13.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	40	31	9	41	29	12	164.5	167.7	157.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	4	0	2	2	0	8.0	11.6	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	13	9	20	12	8	80.2	69.4	104.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	68	57	11	62	50	12	248.7	289.2	157.1
Other hypertensive disease (444-447)	1	0	1	2	1	1	8.0	5.8	13.1
General arteriosclerosis (450)	2	2	0	4	3	1	16.0	17.4	13.1
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	1	0	1	4	2	2	16.0	11.6	26.2
Bronchitis (500-502)	1	1	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	5	5	0	5	5	0	20.0	28.9	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	2	2	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	4.0	-	13.1
Cirrhosis of liver (581)	0	0	0	1	1	0	4.0	5.8	-
Acute nephritis (590, 591)	1	0	1	1	0	1	4.0	-	13.1
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	4.0	-	13.1
Infections of kidney (600)	1	1	0	1	1	0	4.0	-	13.1
Hyperplasia of prostate (610)	3	2	1	3	2	1	12.0	11.6	13.1
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	1	1	2	1	1	8.0	5.8	13.1
Diseases of early infancy (760-776)	8	0	8	11	3	8	23.6	11.0	41.0
Accidents (800-962)	17	10	7	15	7	8	60.2	40.5	104.7
Motor vehicle accidents (810-835)	7	5	2	5	2	3	20.0	11.6	39.3
Suicide (963, 970-979)	3	3	0	4	4	0	16.0	23.1	-
Homicide (964, 980-985)	1	0	1	1	0	1	4.0	-	13.1
Ill-defined (780-795)	6	3	3	5	2	3	20.0	11.6	39.3

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LAMAR COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	79	56	23	257	196	61	17.1	15.3	27.0
Fetal deaths	1	1	0	8	3	5	31.1	15.3	82.0
Total deaths	77	54	23	136	109	27	9.0	8.5	11.9
Deaths under 1 year	2	0	2	6	3	3	23.3	15.3	49.2
Deaths under 28 days	0	0	0	4	3	1	15.6	15.3	16.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	8	3	20	17	3	132.8	132.8	132.7
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	6.6	7.8	-
Asthma (241)	1	1	0	1	1	0	6.6	7.8	-
Diabetes mellitus (260)	1	1	0	1	1	0	6.6	7.8	-
Anemias (290-293)	0	0	0	1	1	0	6.6	7.8	-
Intracranial lesions (330-334)	15	9	6	27	20	7	179.3	156.2	309.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	25	19	6	39	32	7	259.0	250.0	309.7
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	7	4	3	7	4	3	46.5	31.2	132.7
Other circulatory diseases (451-468)	1	0	1	3	2	1	19.9	15.6	44.2
#Influenza & pneumonia (480-493)	4	2	2	8	6	2	53.1	46.9	88.5
Bronchitis (500-502)	0	0	0	1	1	0	6.6	7.8	-
Other bronchopul. diseases (525-527)	1	0	1	2	1	1	13.3	7.8	44.2
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	6.6	7.8	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	1	0	6.6	7.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	0	0	0	-	-	-
Accidents (800-962)	3	3	0	4	3	1	15.6	15.3	16.4
Motor vehicle accidents (810-835)	3	3	0	12	11	1	79.7	85.9	44.2
Suicide (963, 970-979)	1	1	0	7	7	0	46.5	54.7	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	1	1	0	0	0	0	6.6	7.8	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LAUDERDALE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,951	999	952	1,415	748	667	18.4	14.3	27.2
Fetal deaths	108	72	36	81	52	29	57.2	69.5	43.5
Total deaths	986	627	359	770	459	311	10.0	8.8	12.7
Deaths under 1 year	66	22	44	50	16	34	35.3	21.4	51.0
Deaths under 28 days	49	18	31	32	13	19	22.6	17.4	28.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	2	1	1	2.6	1.9	4.1
Syphilis (020-029)	5	4	1	2	1	1	2.6	1.9	4.1
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	125	84	41	108	67	41	140.8	128.3	167.4
Benign & unspec. neoplasms (210-239)	3	2	1	2	1	1	2.6	1.9	4.1
Asthma (241)	7	6	1	5	4	1	6.5	7.7	4.1
Diabetes mellitus (260)	10	2	8	8	1	7	10.4	1.9	28.6
Anemias (290-293)	2	1	1	1	1	0	1.3	1.9	-
Intracranial lesions (330-334)	148	102	46	123	85	38	160.4	162.8	155.2
Nonmeningococcal meningitis (340)	5	1	4	3	1	2	3.9	1.9	8.2
Heart diseases (400-443)	275	200	75	207	135	72	269.9	258.6	294.0
Other hypertensive disease (444-447)	6	2	4	4	2	2	5.2	3.8	8.2
General arteriosclerosis (450)	12	9	3	7	4	3	9.1	7.7	12.2
Other circulatory diseases (451-468)	13	11	2	10	8	2	13.0	15.3	8.2
#Influenza & pneumonia (480-493)	44	22	22	31	16	15	40.4	30.6	61.2
Bronchitis (500-502)	0	0	0	1	1	0	1.3	1.9	-
Other bronchopul. diseases (525-527)	19	15	4	18	14	4	23.5	26.8	16.3
Stomach & duodenal ulcers (540, 541)	3	3	0	2	2	0	2.6	3.8	-
Hernia & intestinal obstruction (560, 561, 570)	5	4	1	4	3	1	5.2	5.7	4.1
#Gastritis, etc. (543, 571, 572)	8	7	1	1	1	0	1.3	1.9	-
Cirrhosis of liver (581)	6	4	2	3	1	2	3.9	1.9	8.2
Acute nephritis (590, 591)	1	1	0	1	1	0	1.3	1.9	-
Chronic nephritis, etc. (592-594)	1	1	0	1	1	0	1.3	1.9	-
Infections of kidney (600)	16	11	5	6	3	3	7.8	5.7	12.2
Hyperplasia of prostate (610)	4	3	1	2	1	1	2.6	1.9	4.1
Maternal causes (640-689)	1	1	0	1	1	0	7.1	13.4	-
Congenital malformations (750-759)	6	5	1	7	5	2	9.1	9.6	8.2
Diseases of early infancy (760-776)	38	13	25	23	8	15	16.2	10.7	22.5
Accidents (800-962)	72	48	24	57	37	20	74.3	70.9	81.7
Motor vehicle accidents (810-835)	26	21	5	21	18	3	27.4	34.5	12.2
Suicide (963, 970-979)	5	4	1	5	5	0	6.5	9.6	-
Homicide (964, 980-985)	8	2	6	8	2	6	10.4	3.8	24.5
Ill-defined (780-795)	96	30	66	90	27	63	117.3	51.7	257.2

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LAWRENCE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	102	42	60	184	97	87	16.2	12.4	24.5
Fetal deaths	2	2	0	6	3	3	32.6	30.9	34.5
Total deaths	89	56	33	142	99	43	12.5	12.6	12.1
Deaths under 1 year	2	0	2	3	0	3	16.3	-	34.5
Deaths under 28 days	2	0	2	2	0	2	10.9	-	23.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	8.8	12.8	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	6	5	15	10	5	131.7	127.6	140.8
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	8.8	12.8	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	5	4	1	43.9	51.0	28.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	19	10	9	21	11	10	184.4	140.3	281.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	32	23	9	49	40	9	430.2	510.2	253.5
Other hypertensive disease (444-447)	0	0	0	3	3	0	26.3	38.3	-
General arteriosclerosis (450)	2	1	1	2	1	1	17.6	12.8	28.2
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	1	1	0	3	2	1	26.3	25.5	28.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	3	0	4	4	0	35.1	51.0	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	8.8	12.8	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	8.8	-	28.2
Cirrhosis of liver (581)	0	0	0	1	1	0	8.8	12.8	-
Acute nephritis (590, 591)	1	0	1	1	0	1	8.8	-	28.2
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	1	0	1	1	0	1	5.4	-	11.5
Accidents (800-962)	5	5	0	17	13	4	149.2	165.8	112.7
Motor vehicle accidents (810-835)	2	2	0	14	10	4	122.9	127.6	112.7
Suicide (963, 970-979)	1	1	0	1	1	0	8.8	12.8	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
III-defined (780-795)	8	4	4	11	5	6	96.6	63.8	169.0

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LEAKE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	215	108	107	320	138	182	15.6	10.5	24.8
Fetal deaths	3	1	2	14	5	9	43.8	36.2	49.4
Total deaths	162	109	53	197	135	62	9.6	10.2	8.4
Deaths under 1 year	8	1	7	12	4	8	37.5	29.0	44.0
Deaths under 28 days	1	0	1	2	1	1	6.2	7.2	5.5
<b>SELECTED CAUSES OF DEATH</b>									
AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	4.9	-	13.6
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	26	17	9	35	25	10	170.7	190.0	136.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	4.9	-	13.6
Diabetes mellitus (260)	2	1	1	4	1	3	19.5	7.6	40.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	28	18	10	34	24	10	165.8	182.4	136.2
Nonmeningococcal meningitis (340)	0	0	0	1	1	0	4.9	7.6	-
Heart diseases (400-443)	57	45	12	63	49	14	307.3	372.3	190.7
Other hypertensive disease (444-447)	1	0	1	2	0	2	9.8	-	27.2
General arteriosclerosis (450)	4	3	1	7	6	1	34.1	45.6	13.6
Other circulatory diseases (451-468)	0	0	0	2	2	0	9.8	15.2	-
#Influenza & pneumonia (480-493)	8	1	7	8	1	7	39.0	7.6	95.4
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	3	0	3	3	0	14.6	22.8	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	4.9	7.6	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	4.9	7.6	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	4.9	-	13.6
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	4	4	0	3	3	0	14.6	22.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	3	1	2	14.6	7.6	27.2
Diseases of early infancy (760-776)	2	1	1	2	1	1	6.2	7.2	5.5
Accidents (800-962)	14	10	4	12	8	4	58.5	60.8	54.5
Motor vehicle accidents (810-835)	9	7	2	5	3	2	24.4	22.8	27.2
Suicide (963, 970-979)	4	4	0	3	3	0	14.6	22.8	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	4	0	4	7	3	4	34.1	22.8	54.5

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LEE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,252	988	264	968	709	259	19.7	18.4	24.2
Fetal deaths	25	9	16	33	14	19	34.1	19.7	73.4
Total deaths	496	382	114	452	339	113	9.2	8.8	10.5
Deaths under 1 year	38	28	10	37	27	10	38.2	38.1	38.6
Deaths under 28 days	27	22	5	24	20	4	24.8	28.2	15.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	1	0	2.0	2.6	-
Syphilis (020-029)	1	1	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	69	55	14	65	52	13	132.2	135.3	121.3
Benign & unspec. neoplasms (210-239)	2	2	0	2	2	0	4.1	5.2	-
Asthma (241)	2	2	0	3	3	0	6.1	7.8	-
Diabetes mellitus (260)	4	3	1	3	2	1	6.1	5.2	9.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	61	56	5	60	52	8	122.0	135.3	74.6
Nonmeningococcal meningitis (340)	1	0	1	2	0	2	4.1	-	18.6
Heart diseases (400-443)	196	142	54	171	121	50	347.8	314.8	466.4
Other hypertensive disease (444-447)	2	0	2	2	0	2	4.1	-	18.6
General arteriosclerosis (450)	5	5	0	4	3	1	8.1	7.8	9.3
Other circulatory diseases (451-468)	7	6	1	9	7	2	18.3	18.2	18.6
#Influenza & pneumonia (480-493)	8	6	2	6	4	2	12.2	10.4	18.6
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	3	0	3	3	0	6.1	7.8	-
Stomach & duodenal ulcers (540, 541)	4	3	1	2	1	1	4.1	2.6	9.3
Hernia & intestinal obstruction (560, 561, 570)	5	5	0	2	2	0	4.1	5.2	-
#Gastritis, etc. (543, 571, 572)	4	2	2	6	4	2	12.2	10.4	18.6
Cirrhosis of liver (581)	4	3	1	3	2	1	6.1	5.2	9.3
Acute nephritis (590, 591)	4	2	2	2	1	1	4.1	2.6	9.3
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	2.0	-	9.3
Infections of kidney (600)	3	3	0	5	5	0	10.2	13.0	-
Hyperplasia of prostate (610)	1	1	0	1	1	0	2.0	2.6	-
Maternal causes (640-689)	1	1	0	0	0	0	-	-	-
Congenital malformations (750-759)	8	7	1	10	8	2	20.3	20.8	18.6
Diseases of early infancy (760-776)	19	15	4	17	14	3	17.6	19.7	11.6
Accidents (800-962)	45	37	8	38	27	11	77.3	70.2	102.6
Motor vehicle accidents (810-835)	22	17	5	21	14	7	42.7	36.4	65.3
Suicide (963, 970-979)	3	3	0	2	2	0	4.1	5.2	-
Homicide (964, 980-985)	6	0	6	7	2	5	14.2	5.2	46.6
Ill-defined (780-795)	4	3	1	7	6	1	14.2	15.6	9.3

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LEFLORE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,195	413	782	1,109	348	761	22.3	18.2	24.9
Fetal deaths	57	18	39	59	18	41	53.2	51.7	53.9
Total deaths	493	197	296	452	160	292	9.1	8.4	9.5
Deaths under 1 year	50	5	45	46	4	42	41.5	11.5	55.2
Deaths under 28 days	30	4	26	25	3	22	22.5	8.6	28.9
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	1	0	1	1	0	1	2.0	-	3.3
Malignant neoplasms (140-205)	61	28	33	58	25	33	116.6	130.6	107.9
Benign & unspec. neoplasms (210-239)	2	0	2	2	0	2	4.0	-	6.5
Asthma (241)	1	1	0	0	0	0	-	-	-
Diabetes mellitus (260)	7	3	4	7	2	5	14.1	10.4	16.3
Anemias (290-293)	2	0	2	2	0	2	4.0	-	6.5
Intracranial lesions (330-334)	71	34	37	65	33	32	130.7	172.4	104.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	139	63	76	130	49	81	261.4	256.0	264.8
Other hypertensive disease (444-447)	6	1	5	6	1	5	12.1	5.2	16.3
General arteriosclerosis (450)	8	2	6	7	3	4	14.1	15.7	13.1
Other circulatory diseases (451-468)	7	5	2	5	3	2	10.0	15.7	6.5
#Influenza & pneumonia (480-493)	17	2	15	18	2	16	36.2	10.4	52.3
Bronchitis (500-502)	1	1	0	1	1	0	2.0	5.2	-
Other bronchopul. diseases (525-527)	8	6	2	8	6	2	16.1	31.3	6.5
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	2.0	-	3.3
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	8	3	5	5	0	5	10.0	-	16.3
Cirrhosis of liver (581)	1	1	0	1	1	0	2.0	5.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	2	0	2	4.0	-	6.5
Infections of kidney (600)	10	3	7	6	1	5	12.1	5.2	16.3
Hyperplasia of prostate (610)	2	1	1	1	0	1	2.0	-	3.3
Maternal causes (640-689)	1	0	1	1	0	1	9.0	-	13.1
Congenital malformations (750-759)	5	2	3	4	3	1	8.0	15.7	3.3
Diseases of early infancy (760-776)	20	2	18	17	1	16	15.3	2.9	21.0
Accidents (800-962)	38	18	20	27	8	19	54.3	41.8	62.1
Motor vehicle accidents (810-835)	20	11	9	11	2	9	22.1	10.4	29.4
Suicide (963, 970-979)	4	3	1	6	5	1	12.1	26.1	3.3
Homicide (964, 980-985)	6	1	5	8	1	7	16.1	5.2	22.9
Ill-defined (780-795)	49	8	41	50	9	41	100.5	47.0	134.0

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LINCOLN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	517	292	225	472	256	216	18.1	14.4	26.0
Fetal deaths	19	10	9	29	13	16	61.4	50.8	74.1
Total deaths	297	200	97	312	211	101	11.9	11.8	12.1
Deaths under 1 year	14	5	9	17	5	12	36.0	19.5	55.6
Deaths under 28 days	8	4	4	11	4	7	23.3	15.6	32.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	0	1	1	0	1	3.8	-	12.0
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	37	27	10	40	30	10	153.1	168.5	120.2
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	3.8	5.6	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	2	0	2	7.6	-	24.0
Anemias (290-293)	2	0	2	2	0	2	7.6	-	24.0
Intracranial lesions (330-334)	34	19	15	38	19	19	145.5	106.7	228.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	124	95	29	118	88	30	451.8	494.4	360.6
Other hypertensive disease (444-447)	1	1	0	0	0	0	-	-	-
General arteriosclerosis (450)	3	2	1	4	3	1	15.3	16.8	12.0
Other circulatory diseases (451-468)	1	1	0	2	2	0	7.6	11.2	-
#Influenza & pneumonia (480-493)	14	5	9	14	6	8	53.6	33.7	96.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	6	5	1	7	6	1	26.8	33.7	12.0
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	3.8	-	12.0
Cirrhosis of liver (581)	2	2	0	2	2	0	7.6	11.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	5	0	6	6	0	23.0	33.7	-
Infections of kidney (600)	3	2	1	3	2	1	11.5	11.2	12.0
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	6	3	3	9	3	6	19.1	11.7	27.8
Accidents (800-962)	29	20	9	32	27	5	122.5	151.7	60.1
Motor vehicle accidents (810-835)	21	14	7	15	14	1	57.4	78.6	12.0
Suicide (963, 970-979)	1	1	0	1	1	0	3.8	5.6	-
Homicide (964, 980-985)	1	0	1	2	1	1	7.6	5.6	12.0
Ill-defined (780-795)	22	9	13	23	10	13	88.0	56.2	156.2

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
LOWNDES COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,175	660	515	1,123	599	524	20.3	16.9	26.2
Fetal deaths	34	14	20	34	14	20	30.3	23.4	38.2
Total deaths	417	202	215	467	242	225	8.4	6.8	11.2
Deaths under 1 year	34	13	21	33	13	20	29.4	21.7	38.2
Deaths under 28 days	20	10	10	19	9	10	16.9	15.0	19.1
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	3	2	1	5.4	5.6	5.0
Syphilis (020-029)	1	0	1	1	0	1	1.8	-	5.0
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	1	1	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	48	33	15	56	37	19	101.1	104.6	94.9
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	1.8	2.8	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	0	4	5	1	4	9.0	2.8	20.0
Anemias (290-293)	2	1	1	2	1	1	3.6	2.8	5.0
Intracranial lesions (330-334)	64	30	34	66	34	32	119.2	96.2	159.8
Nonmeningococcal meningitis (340)	2	0	2	2	0	2	3.6	-	10.0
Heart diseases (400-443)	134	75	59	146	83	63	263.6	234.7	314.7
Other hypertensive disease (444-447)	7	4	3	8	5	3	14.4	14.1	15.0
General arteriosclerosis (450)	6	3	3	10	5	5	18.0	14.1	25.0
Other circulatory diseases (451-468)	5	3	2	7	5	2	12.6	14.1	10.0
#Influenza & pneumonia (480-493)	9	3	6	13	5	8	23.5	14.1	40.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	1	1	3	2	1	5.4	5.6	5.0
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	5	2	3	4	3	1	7.2	8.5	5.0
#Gastritis, etc. (543, 571, 572)	2	1	1	3	1	2	5.4	2.8	10.0
Cirrhosis of liver (581)	3	1	2	2	0	2	3.6	-	10.0
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	3	2	1	5.4	5.6	5.0
Infections of kidney (600)	4	2	2	6	3	3	10.8	8.5	15.0
Hyperplasia of prostate (610)	0	0	0	1	1	0	1.8	2.8	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	1	2	4	2	2	7.2	5.6	10.0
Diseases of early infancy (760-776)	20	10	10	19	9	10	16.9	15.0	19.1
Accidents (800-962)	35	15	20	42	22	20	75.8	62.2	99.9
Motor vehicle accidents (810-835)	17	10	7	17	10	7	30.7	28.3	35.0
Suicide (963, 970-979)	1	1	0	2	2	0	3.6	5.6	-
Homicide (964, 980-985)	5	3	2	4	2	2	7.2	5.6	10.0
Ill-defined (780-795)	41	5	36	43	7	36	77.6	19.8	179.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 MADISON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	599	98	501	792	174	618	24.7	16.8	28.4
Fetal deaths	37	10	27	52	17	35	65.6	97.7	56.6
Total deaths	300	103	197	334	102	232	10.4	9.9	10.7
Deaths under 1 year	31	6	25	38	6	32	48.0	34.5	51.8
Deaths under 28 days	17	5	12	21	4	17	26.5	23.0	27.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	35	22	13	36	22	14	112.2	213.0	64.4
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	3.1	-	4.6
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	6	3	3	5	2	3	15.6	19.4	13.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	35	10	25	41	14	27	127.8	135.5	124.2
Nonmeningococcal meningitis (340)	0	0	0	3	0	3	9.4	-	13.8
Heart diseases (400-443)	76	30	46	83	29	54	258.8	280.7	248.4
Other hypertensive disease (444-447)	2	0	2	6	1	5	18.7	9.7	23.0
General arteriosclerosis (450)	4	4	0	3	3	0	9.4	29.0	-
Other circulatory diseases (451-468)	3	1	2	3	1	2	9.4	9.7	9.2
#Influenza & pneumonia (480-493)	8	0	8	8	0	8	24.9	-	36.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	1	0	1	1	0	3.1	9.7	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	3.1	9.7	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	0	2	6.2	-	9.2
#Gastritis, etc. (543, 571, 572)	3	0	3	3	0	3	9.4	-	13.8
Cirrhosis of liver (581)	2	0	2	2	0	2	6.2	-	9.2
Acute nephritis (590, 591)	1	0	1	1	0	1	3.1	-	4.6
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	3.1	-	4.6
Infections of kidney (600)	0	0	0	1	1	0	3.1	9.7	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	1	0	1	12.6	-	16.2
Congenital malformations (750-759)	4	2	2	6	2	4	18.7	19.4	18.4
Diseases of early infancy (760-776)	15	5	10	19	4	15	24.0	23.0	24.3
Accidents (800-962)	35	16	19	34	9	25	106.0	87.1	115.0
Motor vehicle accidents (810-835)	19	11	8	20	7	13	62.4	67.8	59.8
Suicide (963, 970-979)	1	0	1	0	0	0	-	-	-
Homicide (964, 980-985)	4	1	3	5	2	3	15.6	19.4	13.8
Ill-defined (780-795)	57	4	53	58	4	54	180.8	38.7	248.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
MARION COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	364	212	152	457	252	205	19.4	16.1	26.0
Fetal deaths	9	3	6	14	6	8	30.6	23.8	39.0
Total deaths	225	149	76	248	155	93	10.5	9.9	11.8
Deaths under 1 year	10	4	6	14	6	8	30.6	23.8	39.0
Deaths under 28 days	6	3	3	9	4	5	19.7	15.9	24.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	4	4	0	17.0	25.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	29	19	10	32	23	9	135.9	146.8	114.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	2	0	5	2	3	21.2	12.8	38.1
Anemias (290-293)	1	0	1	1	0	1	4.2	-	12.7
Intracranial lesions (330-334)	46	34	12	43	24	19	182.7	153.2	241.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	61	41	20	66	44	22	280.4	280.8	279.5
Other hypertensive disease (444-447)	2	0	2	1	0	1	4.2	-	12.7
General arteriosclerosis (450)	4	4	0	2	2	0	8.5	12.8	-
Other circulatory diseases (451-468)	0	0	0	2	1	1	8.5	6.4	12.7
#Influenza & pneumonia (480-493)	8	6	2	9	7	2	38.2	44.7	25.4
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	2	1	1	8.5	6.4	12.7
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	4.2	6.4	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	4.2	-	12.7
Cirrhosis of liver (581)	1	1	0	1	1	0	4.2	6.4	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	4.2	-	12.7
Infections of kidney (600)	0	0	0	1	1	0	4.2	6.4	-
Hyperplasia of prostate (610)	1	1	0	1	1	0	4.2	6.4	-
Maternal causes (640-689)	2	1	1	2	1	1	43.8	39.7	48.8
Congenital malformations (750-759)	1	1	0	2	1	1	8.5	6.4	12.7
Diseases of early infancy (760-776)	5	2	3	7	3	4	15.3	11.9	19.5
Accidents (800-962)	17	11	6	26	17	9	110.4	108.5	114.4
Motor vehicle accidents (810-835)	10	7	3	12	9	3	51.0	57.4	38.1
Suicide (963, 970-979)	1	1	0	1	1	0	4.2	6.4	-
Homicide (964, 980-985)	2	0	2	2	0	2	8.5	-	25.4
Ill-defined (780-795)	29	13	16	28	13	15	118.9	83.0	190.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
MARSHALL COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	342	22	320	516	129	387	20.9	15.3	23.8
Fetal deaths	18	2	16	28	5	23	54.3	38.8	59.4
Total deaths	188	70	118	263	101	162	10.7	12.0	10.0
Deaths under 1 year	10	0	10	19	4	15	36.8	31.0	38.8
Deaths under 28 days	6	0	6	11	2	9	21.3	15.5	23.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	0	1	4.0	-	6.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	0	1	4.0	-	6.2
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	5	6	31	14	17	125.6	166.3	104.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	1	0	1	4.0	-	6.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	10	17	41	13	28	166.2	154.4	172.3
Nonmeningococcal meningitis (340)	0	0	0	1	1	0	4.0	11.9	-
Heart diseases (400-443)	71	30	41	90	42	48	364.8	498.8	295.4
Other hypertensive disease (444-447)	1	0	1	3	1	2	12.2	11.9	12.3
General arteriosclerosis (450)	1	0	1	1	0	1	4.0	-	6.2
Other circulatory diseases (451-468)	1	1	0	3	0	3	12.2	-	18.5
#Influenza & pneumonia (480-493)	3	0	3	5	0	5	20.3	-	30.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	2	1	4	2	2	16.2	23.8	12.3
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	3	1	2	12.2	11.9	12.3
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	4.0	-	6.2
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	5	3	2	5	3	2	20.3	35.6	12.3
Infections of kidney (600)	1	0	1	1	0	1	4.0	-	6.2
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	6	0	6	9	1	8	17.4	7.8	20.7
Accidents (800-962)	25	9	16	25	11	14	101.3	130.6	86.2
Motor vehicle accidents (810-835)	12	7	5	12	8	4	48.6	95.0	24.6
Suicide (963, 970-979)	1	1	0	1	1	0	4.0	11.9	-
Homicide (964, 980-985)	4	1	3	2	0	2	8.1	-	12.3
Ill-defined (780-795)	21	7	14	23	7	16	93.2	83.1	98.5

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
MONROE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	747	466	281	674	381	293	20.6	17.4	27.1
Fetal deaths	20	7	13	22	5	17	32.6	13.1	58.0
Total deaths	330	230	100	357	239	118	10.9	10.9	10.9
Deaths under 1 year	20	14	6	21	12	9	31.2	31.5	30.7
Deaths under 28 days	15	10	5	15	9	6	22.2	23.6	20.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	43	31	12	46	35	11	140.5	159.6	101.8
Benign & unspec. neoplasms (210-239)	2	0	2	3	1	2	9.2	4.6	18.5
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	5	2	3	5	3	2	15.3	13.7	18.5
Anemias (290-293)	0	0	0	1	1	0	3.0	4.6	-
Intracranial lesions (330-334)	42	21	21	47	22	25	143.6	100.3	231.5
Nonmeningococcal meningitis (340)	1	1	0	1	1	0	3.0	4.6	-
Heart diseases (400-443)	135	111	24	138	108	30	421.6	492.5	277.8
Other hypertensive disease (444-447)	5	2	3	5	2	3	15.3	9.1	27.8
General arteriosclerosis (450)	4	2	2	6	3	3	18.3	13.7	27.8
Other circulatory diseases (451-468)	2	1	1	3	2	1	9.2	9.1	9.2
#Influenza & pneumonia (480-493)	6	5	1	9	7	2	27.5	31.9	18.5
Bronchitis (500-502)	1	1	0	1	1	0	3.0	4.6	-
Other bronchopul. diseases (525-527)	4	4	0	4	4	0	12.2	18.2	-
Stomach & duodenal ulcers (540, 541)	2	1	1	2	1	1	6.1	4.6	9.2
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	3.0	-	9.2
#Gastritis, etc. (543, 571, 572)	3	3	0	2	2	0	6.1	9.1	-
Cirrhosis of liver (581)	1	1	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	1	0	3.0	4.6	-
Chronic nephritis, etc. (592-594)	14	4	10	15	4	11	45.8	18.2	101.8
Infections of kidney (600)	1	1	0	1	1	0	3.0	4.6	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	2	2	0	29.7	52.5	-
Congenital malformations (750-759)	0	0	0	2	1	1	6.1	4.6	9.2
Diseases of early infancy (760-776)	15	10	5	15	9	6	22.2	23.6	20.5
Accidents (800-962)	22	19	3	25	21	4	76.4	95.8	37.0
Motor vehicle accidents (810-835)	12	11	1	10	8	2	30.6	36.5	18.5
Suicide (963, 970-979)	1	1	0	1	1	0	3.0	4.6	-
Homicide (964, 980-985)	3	0	3	3	0	3	9.2	-	27.8
Ill-defined (780-795)	6	2	4	5	0	5	15.3	-	46.3

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
MONTGOMERY COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	310	124	186	256	93	163	17.2	10.9	25.5
Fetal deaths	9	3	6	7	2	5	27.3	21.5	30.7
Total deaths	154	90	64	142	86	56	9.5	10.1	8.8
Deaths under 1 year	7	1	6	6	1	5	23.4	10.8	30.7
Deaths under 28 days	5	0	5	4	0	4	15.6	-	24.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	11	7	18	11	7	120.6	128.8	109.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	6.7	11.7	-
Diabetes mellitus (260)	1	0	1	2	1	1	13.4	11.7	15.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	31	16	15	24	16	8	160.8	187.4	125.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	61	42	19	55	39	16	368.6	456.7	250.8
Other hypertensive disease (444-447)	2	0	2	3	1	2	20.1	11.7	31.3
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	3	3	0	1	0	1	6.7	-	15.7
#Influenza & pneumonia (480-493)	1	1	0	0	0	0	-	-	-
Bronchitis (500-502)	0	0	0	1	1	0	6.7	11.7	-
Other bronchopul. diseases (525-527)	1	1	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	6.7	-	15.7
Hernia & intestinal obstruction (560, 561, 570)	2	1	1	2	1	1	13.4	11.7	15.7
#Gastritis, etc. (543, 571, 572)	1	1	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	1	0	1	6.7	-	15.7
Acute nephritis (590, 591)	1	0	1	1	0	1	6.7	-	15.7
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	1	0	1	1	0	6.7	11.7	-
Hyperplasia of prostate (610)	1	1	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	1	1	2	1	1	13.4	11.7	15.7
Diseases of early infancy (760-776)	5	0	5	4	0	4	15.6	-	24.5
Accidents (800-962)	8	4	4	10	5	5	67.0	58.5	78.4
Motor vehicle accidents (810-835)	6	2	4	5	2	3	33.5	23.4	47.0
Suicide (963, 970-979)	1	1	0	2	1	1	13.4	11.7	15.7
Homicide (964, 980-985)	1	0	1	0	0	0	-	-	-
Ill-defined (780-795)	7	3	4	8	4	4	53.6	46.8	62.7

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
NESHOBIA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	365	209	156	365	225	140	18.2	15.1	26.8
Fetal deaths	18	6	12	19	5	14	52.0	22.2	100.0
Total deaths	205	144	61	243	181	62	12.1	12.2	11.8
Deaths under 1 year	10	7	3	16	10	6	43.8	44.4	42.8
Deaths under 28 days	6	5	1	11	8	3	30.1	35.6	21.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	5.0	-	19.1
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	1	0	1	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	1	1	0	1	1	0	5.0	6.7	-
Malignant neoplasms (140-205)	34	28	6	39	34	5	194.0	228.6	95.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	3	1	5	3	2	24.9	20.2	38.2
Anemias (290-293)	0	0	0	1	1	0	5.0	6.7	-
Intracranial lesions (330-334)	43	30	13	45	34	11	223.9	228.6	210.3
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	67	51	16	85	68	17	422.9	457.3	325.0
Other hypertensive disease (444-447)	1	0	1	1	0	1	5.0	-	19.1
General arteriosclerosis (450)	3	3	0	3	3	0	14.9	20.2	-
Other circulatory diseases (451-468)	1	0	1	1	0	1	5.0	-	19.1
#Influenza & pneumonia (480-493)	3	1	2	4	1	3	19.9	6.7	57.4
Bronchitis (500-502)	1	1	0	1	1	0	5.0	6.7	-
Other bronchopul. diseases (525-527)	2	1	1	2	2	0	10.0	13.4	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	1	1	2	1	1	10.0	6.7	19.1
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	1	1	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	1	1	0	5.0	6.7	-
Infections of kidney (600)	0	0	0	4	3	1	19.9	20.2	19.1
Hyperplasia of prostate (610)	1	1	0	1	1	0	5.0	6.7	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	2	1	3	2	1	14.9	13.4	19.1
Diseases of early infancy (760-776)	6	4	2	11	7	4	30.1	31.1	28.6
Accidents (800-962)	15	7	8	16	7	9	79.6	47.1	172.1
Motor vehicle accidents (810-835)	7	3	4	7	3	4	34.8	20.2	76.5
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	2	1	1	10.0	6.7	19.1
Ill-defined (780-795)	4	1	3	4	2	2	19.9	13.4	38.2

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 NEWTON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	278	157	121	304	137	167	15.0	9.5	28.7
Fetal deaths	27	15	12	27	15	12	88.8	109.5	71.8
Total deaths	192	138	54	230	161	69	11.4	11.2	11.8
Deaths under 1 year	6	2	4	8	3	5	26.3	21.9	30.0
Deaths under 28 days	5	2	3	7	3	4	23.0	21.9	24.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	27	18	9	31	19	12	153.3	131.9	206.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	3	1	2	14.8	6.9	34.4
Anemias (290-293)	1	0	1	2	0	2	9.9	-	34.4
Intracranial lesions (330-334)	38	30	8	39	32	7	192.9	222.2	120.3
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	4.9	-	17.2
Heart diseases (400-443)	67	52	15	73	58	15	361.0	402.8	257.7
Other hypertensive disease (444-447)	3	3	0	3	2	1	14.8	13.9	17.2
General arteriosclerosis (450)	1	0	1	2	1	1	9.9	6.9	17.2
Other circulatory diseases (451-468)	2	1	1	4	3	1	19.8	20.8	17.2
#Influenza & pneumonia (480-493)	3	2	1	4	3	1	19.8	20.8	17.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	9.9	13.9	-
Stomach & duodenal ulcers (540, 541)	1	1	0	2	1	1	9.9	6.9	17.2
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	3	3	0	14.8	20.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	4.9	-	17.2
Infections of kidney (600)	0	0	0	1	0	1	4.9	-	17.2
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	1	2	3	1	2	14.8	6.9	34.4
Diseases of early infancy (760-776)	4	2	2	6	3	3	19.7	21.9	18.0
Accidents (800-962)	16	11	5	25	16	9	123.6	111.1	154.6
Motor vehicle accidents (810-835)	8	6	2	11	8	3	54.4	55.6	51.5
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	2	0	2	2	0	2	9.9	-	34.4
Ill-defined (780-795)	9	6	3	10	6	4	49.4	41.7	68.7

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
NOXUBEE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	367	67	300	360	66	294	23.3	14.6	26.9
Fetal deaths	11	1	10	10	0	10	27.8	-	34.0
Total deaths	140	42	98	168	57	111	10.9	12.6	10.2
Deaths under 1 year	18	0	18	19	0	19	52.8	-	64.6
Deaths under 28 days	11	0	11	12	0	12	33.3	-	40.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	0	1	6.5	-	9.2
Dysentery (045-048)	1	0	1	1	0	1	6.5	-	9.2
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	10	4	6	15	6	9	97.2	132.7	82.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	3	1	2	19.4	22.1	18.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	16	3	13	22	8	14	142.5	177.0	128.2
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	35	23	12	38	24	14	246.1	531.0	128.2
Other hypertensive disease (444-447)	3	0	3	4	0	4	25.9	-	36.6
General arteriosclerosis (450)	0	0	0	1	1	0	6.5	22.1	-
Other circulatory diseases (451-468)	2	0	2	3	0	3	19.4	-	27.5
#Influenza & pneumonia (480-493)	3	0	3	7	1	6	45.3	22.1	54.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	0	1	1	0	1	6.5	-	9.2
Stomach & duodenal ulcers (540, 541)	2	0	2	3	1	2	19.4	22.1	18.3
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	6.5	-	9.2
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	0	1	1	0	1	6.5	-	9.2
Hyperplasia of prostate (610)	0	0	0	1	1	0	6.5	22.1	-
Maternal causes (640-689)	1	0	1	1	0	1	27.8	-	34.0
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	10	0	10	11	0	11	30.6	-	37.4
Accidents (800-962)	12	6	6	10	4	6	64.8	88.5	54.9
Motor vehicle accidents (810-835)	6	3	3	5	2	3	32.4	44.2	27.5
Suicide (963, 970-979)	1	0	1	2	1	1	13.0	22.1	9.2
Homicide (964, 980-985)	3	0	3	1	0	1	6.5	-	9.2
Ill-defined (780-795)	28	3	25	32	5	27	207.2	110.6	247.2

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
OKTIBBEHA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	589	276	313	542	274	268	17.7	14.7	22.2
Fetal deaths	14	2	12	15	4	11	27.7	14.6	41.0
Total deaths	208	81	127	226	97	129	7.4	5.2	10.7
Deaths under 1 year	24	5	19	26	7	19	48.0	25.5	70.9
Deaths under 28 days	16	5	11	18	7	11	33.2	25.5	41.0
<b>SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	1	1	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	10	11	27	13	14	88.1	70.0	116.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	1	0	1	3.3	-	8.3
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	9	21	34	14	20	111.0	75.3	165.9
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	65	26	39	69	31	38	225.2	166.8	315.2
Other hypertensive (444-447)	4	0	4	3	0	3	9.8	-	24.9
General arteriosclerosis (450)	4	2	2	4	2	2	13.0	10.8	16.6
Other circulatory diseases (451-468)	1	1	0	1	1	0	3.3	5.4	-
#Influenza & pneumonia (480-493)	3	0	3	4	1	3	13.0	5.4	24.9
Bronchitis (500-502)	0	0	0	1	1	0	3.3	5.4	-
Other bronchopul. diseases (525-527)	1	1	0	1	0	1	3.3	-	8.3
Stomach & duodenal ulcers (540, 541)	2	2	0	2	2	0	6.5	10.8	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	2	1	1	6.5	5.4	8.3
#Gastritis, etc. (543, 571, 572)	6	0	6	6	0	6	19.6	-	49.8
Cirrhosis of liver (581)	2	1	1	2	1	1	6.5	5.4	8.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	3.3	5.4	-
Infections of kidney (600)	1	1	0	1	1	0	3.3	5.4	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.3	-	8.3
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	2	2	0	6.5	10.8	-
Diseases of early infancy (760-776)	16	5	11	16	5	11	29.5	18.2	41.0
Accidents (800-962)	24	13	11	24	12	12	78.3	64.6	99.5
Motor vehicle accidents (810-835)	10	5	5	10	5	5	32.6	26.9	41.5
Suicide (963, 970-979)	2	2	0	1	1	0	3.3	5.4	-
Homicide (964, 980-985)	2	1	1	2	1	1	6.5	5.4	8.3
Ill-defined (780-795)	10	0	10	12	1	11	39.2	5.4	91.2

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
PANOLA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	710	223	487	604	185	419	19.7	14.1	23.9
Fetal deaths	21	0	21	22	1	21	36.4	5.4	50.1
Total deaths	249	97	152	320	132	188	10.4	10.0	10.7
Deaths under 1 year	20	2	18	23	2	21	38.1	10.8	50.1
Deaths under 28 days	15	2	13	15	1	14	24.8	5.4	33.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	3.2	-	5.7
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	26	11	15	35	18	17	114.0	136.8	96.9
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	3.2	-	5.7
Asthma (241)	0	0	0	1	1	0	3.2	7.6	-
Diabetes mellitus (260)	3	1	2	5	3	2	16.3	22.8	11.4
Anemias (290-293)	1	0	1	1	0	1	3.2	-	5.7
Intracranial lesions (330-334)	31	10	21	40	16	24	130.3	121.6	136.8
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	3.2	-	5.7
Heart diseases (400-443)	97	38	59	125	55	70	407.2	418.1	399.1
Other hypertensive disease (444-447)	3	1	2	3	1	2	9.8	7.6	11.4
General arteriosclerosis (450)	5	4	1	6	4	2	19.5	30.4	11.4
Other circulatory diseases (451-468)	2	2	0	4	2	2	13.0	15.2	11.4
#Influenza & pneumonia (480-493)	5	1	4	10	4	6	32.6	30.4	34.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	2	1	4	2	2	13.0	15.2	11.4
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	3.2	-	5.7
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	3	2	1	9.8	15.2	5.7
#Gastritis, etc. (543, 571, 572)	2	1	1	3	1	2	9.8	7.6	11.4
Cirrhosis of liver (581)	1	0	1	1	0	1	3.2	-	5.7
Acute nephritis (590, 591)	0	0	0	1	0	1	3.2	-	5.7
Chronic nephritis, etc. (592-594)	2	0	2	3	0	3	9.8	-	17.1
Infections of kidney (600)	0	0	0	2	1	1	6.5	7.6	5.7
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.2	-	5.7
Maternal causes (640-689)	0	0	0	1	1	0	16.6	54.0	-
Congenital malformations (750-759)	2	0	2	3	1	2	9.8	7.6	11.4
Diseases of early infancy (760-776)	11	2	9	11	1	10	18.2	5.4	23.9
Accidents (800-962)	32	20	12	26	12	14	84.7	91.2	79.8
Motor vehicle accidents (810-835)	16	11	5	13	7	6	42.4	53.2	34.2
Suicide (963, 970-979)	3	3	0	2	2	0	6.5	15.2	-
Homicide (964, 980-985)	1	0	1	0	0	0	-	-	-
Ill-defined (780-795)	13	0	13	15	2	13	48.9	15.2	74.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
PEARL RIVER COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	601	483	118	540	405	135	18.6	16.7	28.0
Fetal deaths	7	6	1	10	8	2	18.5	19.8	14.8
Total deaths	212	168	44	246	191	55	8.5	7.9	11.4
Deaths under 1 year	14	7	7	22	10	12	40.7	24.7	88.9
Deaths under 28 days	10	7	3	17	9	8	31.5	22.2	59.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	19	17	2	27	22	5	92.9	90.8	103.5
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	3.4	4.1	-
Asthma (241)	1	1	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	1	0	1	3.4	-	20.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	18	5	28	23	5	96.3	94.9	103.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	97	77	20	98	77	21	337.1	317.6	434.8
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	7	6	1	7	6	1	24.1	24.8	20.7
Other circulatory diseases (451-468)	2	2	0	4	4	0	13.8	16.5	-
#Influenza & pneumonia (480-493)	10	9	1	13	12	1	44.7	49.5	20.7
Bronchitis (500-502)	0	0	0	1	1	0	3.4	4.1	-
Other bronchopul. diseases (525-527)	1	1	0	1	1	0	3.4	4.1	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	3.4	4.1	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	3.4	4.1	-
#Gastritis, etc. (543, 571, 572)	1	1	0	2	2	0	6.9	8.2	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	3	3	0	1	1	0	3.4	4.1	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	18.5	-	74.1
Congenital malformations (750-759)	0	0	0	2	1	1	6.9	4.1	20.7
Diseases of early infancy (760-776)	10	7	3	16	9	7	29.6	22.2	51.8
Accidents (800-962)	14	10	4	19	15	4	65.4	61.9	82.8
Motor vehicle accidents (810-835)	9	7	2	11	9	2	37.8	37.1	41.4
Suicide (963, 970-979)	3	2	1	3	2	1	10.3	8.2	20.7
Homicide (964, 980-985)	2	2	0	3	2	1	10.3	8.2	20.7
Ill-defined (780-795)	10	6	4	7	3	4	24.1	12.4	82.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
PERRY COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	99	59	40	184	115	69	19.9	17.8	24.6
Fetal deaths	3	3	0	8	4	4	43.5	34.8	58.0
Total deaths	78	54	24	94	68	26	10.2	10.5	9.3
Deaths under 1 year	5	2	3	5	3	2	27.2	26.1	29.0
Deaths under 28 days	2	2	0	3	3	0	16.3	26.1	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	9	4	5	25	18	7	270.0	278.6	250.0
Benign & unspec. neoplasms (210-239)	1	1	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	0	0	0	-	-	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	15	10	5	13	9	4	140.4	139.3	142.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	31	25	6	24	18	6	259.2	278.6	214.3
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	2	2	0	2	2	0	21.6	31.0	-
Other circulatory diseases (451-468)	0	0	0	1	1	0	10.8	15.5	-
#Influenza & pneumonia (480-493)	3	1	2	3	2	1	32.4	31.0	35.7
Bronchitis (500-502)	1	1	0	2	2	0	21.6	31.0	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	10.8	15.5	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	2	1	1	21.6	15.5	35.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	2	2	0	3	3	0	16.3	26.1	-
Accidents (800-962)	9	7	2	11	9	2	118.8	139.3	71.4
Motor vehicle accidents (810-835)	7	7	0	5	5	0	54.0	77.4	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	1	0	1	1	0	1	10.8	-	35.7
Ill-defined (780-795)	4	1	3	5	2	3	54.0	31.0	107.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
PIKE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	612	317	295	553	230	323	17.1	12.5	23.2
Fetal deaths	32	10	22	31	8	23	56.0	34.8	71.2
Total deaths	350	218	132	362	216	146	11.2	11.7	10.5
Deaths under 1 year	22	5	17	24	6	18	43.4	26.1	55.7
Deaths under 28 days	16	4	12	17	4	13	30.7	17.4	40.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	3	0	3	9.3	-	21.6
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	54	41	13	58	46	12	179.7	250.1	86.4
Benign & unspec. neoplasms (210-239)	1	0	1	1	0	1	3.1	-	7.2
Asthma (241)	0	0	0	1	0	1	3.1	-	7.2
Diabetes mellitus (260)	1	1	0	1	0	1	3.1	-	7.2
Anemias (290-293)	2	0	2	2	0	2	6.2	-	14.4
Intracranial lesions (330-334)	62	41	21	60	37	23	185.9	201.2	165.6
Nonmeningococcal meningitis (340)	1	1	0	2	1	1	6.2	5.4	7.2
Heart diseases (400-443)	99	67	32	99	64	35	306.7	348.0	252.0
Other hypertensive disease (444-447)	4	3	1	4	3	1	12.4	16.3	7.2
General arteriosclerosis (450)	9	7	2	11	8	3	34.1	43.5	21.6
Other circulatory diseases (451-468)	3	3	0	7	6	1	21.7	32.6	7.2
#Influenza & pneumonia (480-493)	8	3	5	10	4	6	31.0	21.8	43.2
Bronchitis (500-502)	1	0	1	1	0	1	3.1	-	7.2
Other bronchopul. diseases (525-527)	4	4	0	4	4	0	12.4	21.8	-
Stomach & duodenal ulcers (540, 541)	1	0	1	2	1	1	6.2	5.4	7.2
Hernia & intestinal obstruction (560, 561, 570)	2	0	2	3	0	3	9.3	-	21.6
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	3.1	5.4	-
Cirrhosis of liver (581)	2	2	0	3	3	0	9.3	16.3	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	0	0	0	-	-	-
Infections of kidney (600)	1	0	1	4	1	3	12.4	5.4	21.6
Hyperplasia of prostate (610)	1	0	1	1	0	1	3.1	-	7.2
Maternal causes (640-689)	1	1	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	2	0	2	2	0	6.2	10.9	-
Diseases of early infancy (760-776)	12	2	10	12	2	10	21.7	8.7	31.0
Accidents (800-962)	23	17	6	17	12	5	52.7	65.2	36.0
Motor vehicle accidents (810-835)	16	13	3	11	10	1	34.1	54.4	7.2
Suicide (963, 970-979)	4	4	0	4	4	0	12.4	21.8	-
Homicide (964, 980-985)	5	1	4	5	1	4	15.5	5.4	28.8
Ill-defined (780-795)	33	10	23	33	9	24	102.2	48.9	172.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 PONTOTOC COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	318	236	82	278	209	69	15.2	13.9	21.5
Fetal deaths	4	3	1	2	1	1	7.2	4.8	14.5
Total deaths	181	149	32	215	176	39	11.8	11.7	12.1
Deaths under 1 year	9	2	7	11	4	7	39.6	19.1	101.4
Deaths under 28 days	9	2	7	10	3	7	36.0	14.4	101.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	22	21	1	27	25	2	148.0	166.3	62.3
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	5.5	6.7	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	3	2	1	16.4	13.3	31.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	21	4	31	28	3	170.0	186.3	93.4
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	73	66	7	78	67	11	427.6	445.8	342.7
Other hypertensive disease (444-447)	1	0	1	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	1	1	0	5.5	6.6	-
Other circulatory diseases (451-468)	2	2	0	3	2	1	16.4	13.3	31.2
#Influenza & pneumonia (480-493)	4	4	0	5	5	0	27.4	33.3	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	5	5	0	6	6	0	32.9	40.0	-
Stomach & duodenal ulcers (540, 541)	1	0	1	3	2	1	16.4	13.3	31.2
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.5	6.6	-
#Gastritis, etc. (543, 571, 572)	2	2	0	2	2	0	11.0	13.3	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	0	0	0	-	-	-
Infections of kidney (600)	4	2	2	5	3	2	27.4	20.0	62.3
Hyperplasia of prostate (610)	0	0	0	1	0	1	5.5	-	31.2
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	9	2	7	9	2	7	32.4	9.6	101.4
Accidents (800-962)	14	12	2	21	17	4	115.1	113.1	124.6
Motor vehicle accidents (810-835)	10	8	2	12	10	2	65.8	66.5	62.3
Suicide (963, 970-979)	2	2	0	3	3	0	16.4	20.0	-
Homicide (964, 980-985)	1	0	1	2	0	2	11.0	-	62.3
Ill-defined (780-795)	4	3	1	5	4	1	27.4	26.6	31.2

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
PRENTISS COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	383	331	52	336	284	52	16.2	15.8	21.7
Fetal deaths	17	7	10	11	5	6	32.7	17.6	115.4
Total deaths	195	172	23	198	177	21	9.7	9.8	8.8
Deaths under 1 year	6	5	1	4	4	0	11.9	14.1	-
Deaths under 28 days	3	3	0	2	2	0	6.0	7.0	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	1	1	0	4.9	5.6	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	34	30	4	36	32	4	176.7	178.1	166.7
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	4.9	5.6	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	17	8	22	16	6	108.0	89.0	250.0
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	76	68	8	74	65	9	363.3	361.7	375.0
Other hypertensive disease (444-447)	2	2	0	2	2	0	9.8	11.1	-
General arteriosclerosis (450)	7	7	0	7	7	0	34.4	39.0	-
Other circulatory diseases (451-468)	3	2	1	4	3	1	19.6	16.7	41.7
#Influenza & pneumonia (480-493)	5	5	0	5	5	0	24.5	27.8	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	3	3	0	3	3	0	14.7	16.7	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	1	0	4.9	5.6	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	1	0	1	1	0	4.9	5.6	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	1	1	0	4.9	5.6	-
Hyperplasia of prostate (610)	1	1	0	1	1	0	4.9	5.6	-
Maternal causes (640-689)	0	0	0	1	1	0	29.8	35.2	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	3	3	0	2	2	0	6.0	7.0	-
Accidents (800-962)	11	10	1	12	12	0	58.9	66.8	-
Motor vehicle accidents (810-835)	8	7	1	7	7	0	34.4	39.0	-
Suicide (963, 970-979)	3	3	0	2	2	0	9.8	11.1	-
Homicide (964, 980-985)	2	1	1	2	1	1	9.8	5.6	41.7
Ill-defined (780-795)	9	9	0	11	11	0	54.0	61.2	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
QUITMAN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	492	132	360	474	133	341	24.0	17.1	28.3
Fetal deaths	8	1	7	12	1	11	25.3	7.5	32.2
Total deaths	136	34	102	203	72	131	10.2	9.3	10.9
Deaths under 1 year	25	5	20	26	4	22	54.8	30.1	64.5
Deaths under 28 days	15	5	10	16	4	12	33.8	30.1	35.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	5.0	12.9	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	8	2	6	25	12	13	126.3	154.6	108.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	5.0	12.9	-
Diabetes mellitus (260)	3	0	3	4	1	3	20.2	12.9	24.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	3	6	20	8	12	101.1	103.1	99.8
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	51	16	35	59	23	36	298.1	296.4	299.2
Other hypertensive disease (444-447)	1	0	1	2	0	2	10.1	-	16.6
General arteriosclerosis (450)	0	0	0	1	1	0	5.0	12.9	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	9	1	8	13	4	9	65.7	51.5	74.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	1	1	3	2	1	15.2	25.8	8.3
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	3	0	3	15.2	-	24.9
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	1	0	5.0	12.9	-
Chronic nephritis, etc. (592-594)	2	0	2	4	0	4	20.2	-	33.2
Infections of kidney (600)	2	2	0	2	2	0	10.1	25.8	-
Hyperplasia of prostate (610)	0	0	0	1	0	1	5.0	-	8.3
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	3	0	3	4	2	2	20.2	25.8	16.6
Diseases of early infancy (760-776)	12	5	7	13	3	10	27.4	22.6	29.3
Accidents (800-962)	12	4	8	20	10	10	101.1	128.9	83.1
Motor vehicle accidents (810-835)	3	1	2	10	7	3	50.5	90.2	24.9
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	3	0	3	4	0	4	20.2	-	33.2
Ill-defined (780-795)	12	0	12	14	0	14	70.7	-	116.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
RANKIN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	85	43	42	709	476	233	18.1	18.8	16.8
Fetal deaths	1	1	0	52	35	17	73.3	73.5	73.0
Total deaths	360	210	150	247	158	89	6.3	6.2	6.4
Deaths under 1 year	2	1	1	10	8	2	14.1	16.8	8.6
Deaths under 28 days	2	1	1	9	7	2	12.7	14.7	8.6
<b>SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	4	2	2	0	0	0	-	-	-
Syphilis (020-029)	2	0	2	1	0	1	2.6	-	7.2
Dysentery (045-048)	3	1	2	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	43	30	13	29	22	7	74.1	87.1	50.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	2	0	2	5.1	-	14.4
Diabetes mellitus (260)	8	4	4	9	6	3	23.0	23.8	21.6
Anemias (290-293)	0	0	0	1	0	1	2.6	-	7.2
Intracranial lesions (330-334)	32	16	16	21	9	12	53.7	35.6	86.5
Nonmeningococcal meningitis (340)	1	1	0	0	0	0	-	-	-
Heart diseases (400-443)	90	51	39	92	65	27	235.2	257.4	194.7
Other hypertensive disease (444-447)	2	0	2	2	0	2	5.1	-	14.4
General arteriosclerosis (450)	15	11	4	12	10	2	30.7	39.6	14.4
Other circulatory diseases (451-468)	14	4	10	4	1	3	10.2	4.0	21.6
#Influenza & pneumonia (480-493)	24	14	10	3	2	1	7.7	7.9	7.2
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	7	7	0	5	5	0	12.8	19.8	-
Stomach & duodenal ulcers (540, 541)	5	3	2	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	3	1	2	2	2	0	5.1	7.9	-
#Gastritis, etc. (543, 571, 572)	1	1	0	1	1	0	2.6	4.0	-
Cirrhosis of liver (581)	1	0	1	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	1	0	1	2.6	-	7.2
Chronic nephritis, etc. (592-594)	3	0	3	3	0	3	7.7	-	21.6
Infections of kidney (600)	4	1	3	4	1	3	10.2	4.0	21.6
Hyperplasia of prostate (610)	1	0	1	1	0	1	2.6	-	7.2
Maternal causes (640-689)	0	0	0	1	0	1	14.1	-	42.9
Congenital malformations (750-759)	2	0	2	3	1	2	7.7	4.0	14.4
Diseases of early infancy (760-776)	2	1	1	8	7	1	11.3	14.7	4.3
Accidents (800-962)	32	23	9	17	13	4	43.4	51.5	28.8
Motor vehicle accidents (810-835)	5	4	1	9	7	2	23.0	27.7	14.4
Suicide (963, 970-979)	5	5	0	4	4	0	10.2	15.8	-
Homicide (964, 980-985)	2	1	1	4	2	2	10.2	7.9	14.4
Ill-defined (780-795)	16	11	5	7	3	4	17.9	11.9	28.8

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
SCOTT COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	416	267	149	465	253	212	19.7	16.7	25.0
Fetal deaths	6	2	4	12	5	7	25.8	19.8	33.0
Total deaths	203	132	71	241	150	91	10.2	9.9	10.8
Deaths under 1 year	13	8	5	14	8	6	30.1	31.6	28.3
Deaths under 28 days	9	6	3	11	6	5	23.6	23.7	23.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	17	12	5	25	15	10	105.8	98.9	118.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	1	0	1	4.2	-	11.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	23	14	9	27	17	10	114.3	112.1	118.2
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	100	73	27	104	74	30	440.1	487.8	354.6
Other hypertensive disease (444-447)	1	1	0	1	1	0	4.2	6.6	-
General arteriosclerosis (450)	6	3	3	6	3	3	25.4	19.8	35.5
Other circulatory diseases (451-468)	1	0	1	1	0	1	4.2	-	11.8
#Influenza & pneumonia (480-493)	2	1	1	5	2	3	21.2	13.2	35.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	8.5	13.2	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	4.2	6.6	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	4	2	2	16.9	13.2	23.6
#Gastritis, etc. (543, 571, 572)	1	1	0	3	3	0	12.7	19.8	-
Cirrhosis of liver (581)	1	1	0	2	1	1	8.5	6.6	11.8
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	2	0	2	2	0	8.5	13.2	-
Infections of kidney (600)	0	0	0	1	1	0	4.2	6.6	-
Hyperplasia of prostate (610)	1	0	1	2	0	2	8.5	-	23.6
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	0	1	3	1	2	12.7	6.6	23.6
Diseases of early infancy (760-776)	8	5	3	10	6	4	21.5	23.7	18.9
Accidents (800-962)	21	12	9	23	13	10	97.3	85.7	118.2
Motor vehicle accidents (810-835)	13	8	5	11	6	5	46.6	39.6	59.1
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	3	1	2	2	0	2	8.5	-	23.6
Ill-defined (780-795)	8	2	6	6	1	5	25.4	6.6	59.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
SHARKEY COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	97	11	86	255	54	201	25.1	16.7	29.1
Fetal deaths	1	0	1	8	4	4	31.4	74.1	19.9
Total deaths	91	26	65	111	32	79	10.9	9.9	11.4
Deaths under 1 year	9	0	9	12	1	11	47.0	18.5	54.7
Deaths under 28 days	5	0	5	8	1	7	31.4	18.5	34.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	9.9	-	14.5
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	7	3	4	9	6	3	88.8	185.8	43.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	5	1	4	49.3	31.0	57.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	11	2	9	12	1	11	118.3	31.0	159.2
Nonmeningococcal meningitis (340)	1	0	1	2	1	1	19.7	31.0	14.5
Heart diseases (400-443)	39	11	28	42	10	32	414.2	309.6	463.1
Other hypertensive disease (444-447)	1	0	1	1	0	1	9.9	-	14.5
General arteriosclerosis (450)	0	0	0	2	1	1	19.7	31.0	14.5
Other circulatory diseases (451-468)	1	0	1	3	1	2	29.6	31.0	28.9
#Influenza & pneumonia (480-493)	2	0	2	3	1	2	29.6	31.0	28.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	3	0	3	1	0	1	9.9	-	14.5
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	9.9	-	14.5
Infections of kidney (600)	0	0	0	0	0	1	9.9	-	14.5
Hyperplasia of prostate (610)	0	0	0	2	0	0	-	-	-
Maternal causes (640-689)	0	0	0	2	0	2	19.7	-	28.9
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	2	0	2	1	1	0	9.9	31.0	-
Accidents (800-962)	18	8	10	17	7	10	15.7	-	19.9
Motor vehicle accidents (810-835)	6	5	1	6	5	1	59.2	154.8	14.5
Suicide (963, 970-979)	1	1	0	1	1	0	9.9	31.0	-
Homicide (964, 980-985)	1	0	1	1	0	1	9.9	-	14.5
Ill-defined (780-795)	2	0	2	2	0	2	19.7	-	28.9

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
SIMPSON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	320	196	124	390	218	172	19.6	16.0	27.2
Fetal deaths	11	4	7	18	9	9	46.2	41.3	52.3
Total deaths	230	161	69	236	172	64	11.8	12.6	10.1
Deaths under 1 year	12	2	10	16	6	10	41.0	27.5	58.1
Deaths under 28 days	3	0	3	7	3	4	17.9	13.8	23.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	24	11	13	1	0	1	5.0	-	15.8
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	5.0	7.4	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	24	13	11	24	13	11	120.5	95.6	174.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	4	2	2	3	1	2	15.1	7.4	31.6
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	40	30	10	44	35	9	220.9	257.4	142.4
Nonmeningococcal meningitis (340)	0	0	0	1	1	0	5.0	7.4	-
Heart diseases (400-443)	81	68	13	88	72	16	441.8	529.4	253.2
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	4	3	1	4	3	1	20.1	22.0	15.8
Other circulatory diseases (451-468)	4	3	1	6	6	0	30.1	44.1	-
#Influenza & pneumonia (480-493)	7	5	2	7	4	3	35.1	29.4	47.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	1	0	2	2	0	10.0	14.7	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	0	1	5.0	-	15.8
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	2	1	1	10.0	7.4	15.8
Cirrhosis of liver (581)	2	2	0	2	2	0	10.0	14.7	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	0	1	5.0	-	15.8
Infections of kidney (600)	3	2	1	1	0	1	5.0	-	15.8
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	2	0	2	2	0	2	10.0	-	31.6
Diseases of early infancy (760-776)	2	0	2	6	3	3	15.4	13.8	17.4
Accidents (800-962)	15	15	0	17	14	3	85.3	102.9	47.5
Motor vehicle accidents (810-835)	13	13	0	12	11	1	60.2	80.9	15.8
Suicide (963, 970-979)	1	0	1	2	1	1	10.0	7.4	15.8
Homicide (964, 980-985)	1	0	1	2	0	2	10.0	-	31.6
Ill-defined (780-795)	9	3	6	8	3	5	40.2	22.0	79.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
SMITH COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	16	0	16	260	134	126	19.4	13.9	33.7
Fetal deaths	0	0	0	8	3	5	30.8	22.4	39.7
Total deaths	54	40	14	133	103	30	9.9	10.7	8.0
Deaths under 1 year	3	1	2	13	2	11	50.0	14.9	87.3
Deaths under 28 days	2	1	1	8	2	6	30.8	14.9	47.6
<b>SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	7.5	10.4	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	5	4	1	19	16	3	141.9	165.8	80.2
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	2	2	0	14.9	20.7	-
Anemias (290-293)	1	1	0	1	1	0	7.5	10.4	-
Intracranial lesions (330-334)	4	4	0	18	13	5	134.4	134.7	133.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	23	18	5	51	46	5	380.9	476.7	133.7
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	1	0	1	1	0	7.5	10.4	-
Other circulatory diseases (451-468)	0	0	0	2	2	0	14.9	20.7	-
#Influenza & pneumonia (480-493)	0	0	0	1	0	1	-	-	26.7
Bronchitis (500-502)	1	1	0	1	1	0	7.5	-	10.4
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	1	0	2	1	1	14.9	10.4	26.7
Cirrhosis of liver (581)	1	0	1	2	1	1	14.9	10.4	26.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	2	2	0	14.9	20.7	-
Infections of kidney (600)	0	0	0	2	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	0	0	0	-	-	-
Accidents (800-962)	2	1	1	8	2	6	30.8	14.9	47.6
Motor vehicle accidents (810-835)	11	7	4	11	8	3	82.2	82.9	80.2
Suicide (963, 970-979)	6	5	1	6	6	0	44.8	62.2	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	2	1	1	2	1	1	14.9	10.4	26.7
	1	0	1	3	1	2	22.4	10.4	53.5

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
STONE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	120	79	41	130	78	52	16.2	12.7	27.5
Fetal deaths	2	1	1	2	1	1	15.4	12.8	19.2
Total deaths	66	53	13	77	61	16	9.6	9.9	8.5
Deaths under 1 year	2	2	0	6	4	2	46.2	51.3	38.5
Deaths under 28 days	1	1	0	6	4	2	46.2	51.3	38.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	13	8	5	17	12	5	211.4	195.1	264.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	12.4	16.3	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	10	6	4	7	3	4	87.1	48.8	211.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	30	27	3	28	25	3	348.2	406.5	158.7
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	2	2	0	1	1	0	12.4	16.3	-
#Influenza & pneumonia (480-493)	2	2	0	2	2	0	24.9	32.5	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	3	3	0	37.3	48.8	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	2	2	0	24.9	32.5	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	1	1	0	12.4	16.3	-
Infections of kidney (600)	1	0	1	1	0	1	12.4	-	52.9
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	1	1	0	4	2	2	30.8	51.3	38.5
Accidents (800-962)	3	3	0	8	7	1	99.5	113.8	52.9
Motor vehicle accidents (810-835)	2	2	0	6	6	0	74.6	97.6	-
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	0	0	0	1	1	0	12.4	16.3	-
Ill-defined (780-795)	0	0	0	0	0	0	-	-	-

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 SUNFLOWER COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	801	132	669	951	178	773	21.9	13.0	26.0
Fetal deaths	38	4	34	44	5	39	46.3	28.1	50.4
Total deaths	369	120	249	416	142	274	9.6	10.4	9.2
Deaths under 1 year	34	3	31	37	4	33	38.9	22.5	42.7
Deaths under 28 days	18	3	15	19	4	15	20.0	22.5	19.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	3	1	2	6.9	7.3	6.7
Syphilis (020-029)	1	0	1	1	0	1	2.3	-	3.4
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	54	18	36	63	23	40	145.2	168.4	134.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	3	1	2	3	1	2	6.9	7.3	6.7
Diabetes mellitus (260)	8	3	5	7	3	4	16.1	22.0	13.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	42	14	28	53	21	32	122.1	153.7	107.6
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	2.3	-	3.4
Heart diseases (400-443)	114	39	75	122	47	75	281.1	344.1	252.2
Other hypertensive disease (444-447)	4	0	4	4	0	4	9.2	-	13.4
General arteriosclerosis (450)	4	1	3	6	2	4	13.8	14.6	13.4
Other circulatory diseases (451-468)	4	3	1	7	3	4	16.1	22.0	13.4
#Influenza & pneumonia (480-493)	12	4	8	16	4	12	36.9	29.3	40.3
Bronchitis (500-502)	1	1	0	1	1	0	2.3	7.3	-
Other bronchopul. diseases (525-527)	3	2	1	3	1	2	6.9	7.3	6.7
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	2.3	7.3	-
Hernia & intestinal obstruction (560, 561, 570)	2	1	1	1	0	1	2.3	-	3.4
#Gastritis, etc. (543, 571, 572)	5	1	4	5	1	4	11.5	7.3	13.4
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	1	2	6.9	7.3	6.7
Infections of kidney (600)	2	1	1	4	3	1	9.2	22.0	3.4
Hyperplasia of prostate (610)	1	0	1	2	1	1	4.6	7.3	3.4
Maternal causes (640-689)	0	0	0	1	0	1	10.5	-	12.9
Congenital malformations (750-759)	3	0	3	3	0	3	6.9	-	10.1
Diseases of early infancy (760-776)	21	3	18	22	4	18	23.1	22.5	23.3
Accidents (800-962)	37	21	16	33	14	19	76.0	102.5	63.9
Motor vehicle accidents (810-835)	19	15	4	13	10	3	30.0	73.2	10.1
Suicide (963, 970-979)	2	1	1	1	0	1	2.3	-	3.4
Homicide (964, 980-985)	3	1	2	5	2	3	11.5	14.6	10.1
Ill-defined (780-795)	31	3	28	31	5	26	71.4	36.6	87.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
TALLAHATCHIE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	491	88	403	529	109	420	23.0	13.6	28.1
Fetal deaths	18	2	16	19	1	18	35.9	9.2	42.8
Total deaths	208	69	139	264	107	157	11.5	13.4	10.5
Deaths under 1 year	27	3	24	30	3	27	56.7	27.5	64.3
Deaths under 28 days	17	2	15	19	2	17	35.9	18.3	40.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	1	1	0	4.4	12.5	-
Dysentery (045-048)	2	0	2	2	0	2	8.7	-	13.4
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	8	10	31	16	15	135.1	200.2	100.3
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	2	1	3	2	1	13.1	25.0	6.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	7	10	25	14	11	109.0	175.2	73.6
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	79	29	50	95	43	52	414.0	538.2	347.7
Other hypertensive disease (444-447)	4	0	4	2	0	2	8.7	-	13.4
General arteriosclerosis (450)	3	2	1	4	2	2	17.4	25.0	13.4
Other circulatory diseases (451-468)	1	0	1	3	0	3	13.1	-	20.1
#Influenza & pneumonia (480-493)	8	1	7	12	4	8	52.3	50.1	53.5
Bronchitis (500-502)	1	0	1	2	1	1	8.7	12.5	6.7
Other bronchopul. diseases (525-527)	4	3	1	7	6	1	30.5	75.1	6.7
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	4.4	-	6.7
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	4.4	-	6.7
Cirrhosis of liver (581)	1	0	1	2	1	1	8.7	12.5	6.7
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	2	1	1	8.7	12.5	6.7
Infections of kidney (600)	0	0	0	1	0	1	4.4	-	6.7
Hyperplasia of prostate (610)	0	0	0	1	0	1	4.4	-	6.7
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	0	1	4.4	-	6.7
Diseases of early infancy (760-776)	17	2	15	18	2	16	34.0	18.3	38.1
Accidents (800-962)	11	4	7	16	4	12	69.7	50.1	80.2
Motor vehicle accidents (810-835)	4	3	1	5	2	3	21.8	25.0	20.1
Suicide (963, 970-979)	4	4	0	4	4	0	17.4	50.1	-
Homicide (964, 980-985)	5	2	3	4	1	3	17.4	12.5	20.1
Ill-defined (780-795)	20	3	17	19	2	17	82.8	25.0	113.7

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
TATE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	184	0	184	421	147	274	21.2	16.0	25.8
Fetal deaths	4	0	4	15	3	12	35.6	20.4	43.8
Total deaths	86	24	62	155	73	82	7.8	7.9	7.7
Deaths under 1 year	5	0	5	12	2	10	28.5	13.6	36.5
Deaths under 28 days	3	0	3	10	2	8	23.8	13.6	29.2
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	1	0	2	2	0	10.1	21.7	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	17	6	11	26	15	11	131.1	162.9	103.6
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	2	1	1	10.1	10.8	9.4
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	3	6	20	11	9	100.8	119.4	84.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	33	8	25	54	26	28	272.3	282.3	263.6
Other hypertensive disease (444-447)	0	0	0	0	0	0	-	-	-
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	1	1	0	5.0	10.8	-
#Influenza & pneumonia (480-493)	1	1	0	1	1	0	5.0	10.8	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	1	1	0	5.0	10.8	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	5.0	10.8	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	2	1	1	10.1	10.8	9.4
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	5.0	-	9.4
Cirrhosis of liver (581)	0	0	0	1	1	0	5.0	10.8	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	3	1	2	15.1	10.8	18.8
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	0	1	1	0	1	5.0	-	9.4
Diseases of early infancy (760-776)	2	0	2	9	2	7	21.4	13.6	25.5
Accidents (800-962)	10	2	8	17	5	12	85.7	54.3	113.0
Motor vehicle accidents (810-835)	2	2	0	5	2	3	25.2	21.7	28.2
Suicide (963, 970-979)	1	1	0	1	1	0	5.0	10.8	-
Homicide (964, 980-985)	1	0	1	1	0	1	5.0	-	9.4
Ill-defined (780-795)	6	0	6	5	0	5	25.2	-	47.1

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
TIPPAH COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	362	273	89	300	230	70	16.9	15.7	22.3
Fetal deaths	36	31	5	32	25	7	106.7	108.7	100.0
Total deaths	136	99	37	147	114	33	8.3	7.8	10.5
Deaths under 1 year	8	5	3	7	4	3	23.3	17.4	42.8
Deaths under 28 days	6	4	2	5	3	2	16.7	13.0	28.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	19	12	7	21	18	3	118.3	123.2	95.5
Benign & unspec. neoplasms (210-239)	1	1	0	1	1	0	5.6	6.8	-
Asthma (241)	1	1	0	1	1	0	5.6	6.8	-
Diabetes mellitus (260)	1	0	1	1	0	1	5.6	-	31.8
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	17	8	27	21	6	152.1	143.7	191.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	46	34	12	43	31	12	242.2	212.2	382.2
Other hypertensive disease (444-447)	1	0	1	2	1	1	11.3	6.8	31.8
General arteriosclerosis (450)	3	2	1	2	2	0	11.3	13.7	-
Other circulatory diseases (451-468)	2	2	0	2	2	0	11.3	13.7	-
#Influenza & pneumonia (480-493)	1	1	0	3	2	1	16.9	13.7	31.8
Bronchitis (500-502)	1	0	1	1	0	1	5.6	-	31.8
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	2	2	0	11.3	13.7	-
#Gastritis, etc. (543, 571, 572)	0	0	0	2	2	0	11.3	13.7	-
Cirrhosis of liver (581)	3	3	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	1	0	1	1	0	1	5.6	-	31.8
Infections of kidney (600)	1	1	0	1	1	0	5.6	6.8	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	1	1	0	33.3	43.5	-
Congenital malformations (750-759)	2	2	0	2	2	0	11.3	13.7	-
Diseases of early infancy (760-776)	3	2	1	2	1	1	6.7	4.3	14.3
Accidents (800-962)	7	6	1	15	12	3	84.5	82.1	95.5
Motor vehicle accidents (810-835)	3	3	0	8	7	1	45.1	47.9	31.8
Suicide (963, 970-979)	4	4	0	5	5	0	28.2	34.2	-
Homicide (964, 980-985)	3	3	0	2	2	0	11.3	13.7	-
Ill-defined (780-795)	7	4	3	8	5	3	45.1	34.2	95.5

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
TISHOMINGO COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	170	158	12	233	215	18	12.9	12.3	31.0
Fetal deaths	5	5	0	5	5	0	21.4	23.2	-
Total deaths	114	109	5	153	144	9	8.5	8.2	15.5
Deaths under 1 year	1	0	1	3	2	1	12.9	9.3	55.6
Deaths under 28 days	0	0	0	1	1	0	4.3	4.6	-
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	2	2	0	11.1	11.5	-
Syphilis (020-029)	0	0	0	1	1	0	5.5	5.7	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	21	21	0	30	30	0	166.5	172.0	-
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	1	0	1	1	0	5.5	5.7	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	17	17	0	21	21	0	116.5	120.4	-
Nonmeningococcal meningitis (340)	0	0	0	1	1	0	5.5	5.7	-
Heart diseases (400-443)	44	43	1	51	49	2	283.0	281.0	344.8
Other hypertensive disease (444-447)	1	1	0	1	1	0	5.5	5.7	-
General arteriosclerosis (450)	5	3	2	5	3	2	27.7	17.2	344.8
Other circulatory diseases (451-468)	1	1	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	0	0	0	3	2	1	16.6	11.5	172.4
Bronchitis (500-502)	0	0	0	1	1	0	5.5	5.7	-
Other bronchopul. diseases (525-527)	1	1	0	2	2	0	11.1	11.5	-
Stomach & duodenal ulcers (540, 541)	1	0	1	1	0	1	5.5	-	172.4
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	1	1	0	5.5	5.7	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	5.5	5.7	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	0	0	0	1	1	0	4.3	4.6	-
Accidents (800-962)	10	10	0	16	15	1	88.8	86.0	172.4
Motor vehicle accidents (810-835)	4	4	0	9	8	1	49.9	45.9	172.4
Suicide (963, 970-979)	0	0	0	2	2	0	11.1	11.5	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	5	4	1	5	4	1	27.7	22.9	172.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
TUNICA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	367	45	322	387	61	326	24.8	15.2	28.1
Fetal deaths	12	0	12	11	0	11	28.4	-	33.7
Total deaths	135	36	99	158	37	121	10.1	9.2	10.4
Deaths under 1 year	16	0	16	20	1	19	51.7	16.4	58.3
Deaths under 28 days	8	0	8	8	1	7	20.7	16.4	21.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	2	1	1	3	1	2	19.2	24.9	17.2
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	12	5	7	20	7	13	128.0	174.1	112.1
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	0	2	4	2	2	25.6	49.8	17.2
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	5	21	29	7	22	185.6	174.1	189.6
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	6.4	-	8.6
Heart diseases (400-443)	39	12	27	40	10	30	256.1	248.8	258.6
Other hypertensive disease (444-447)	1	0	1	1	0	1	6.4	-	8.6
General arteriosclerosis (450)	0	0	0	0	0	0	-	-	-
Other circulatory diseases (451-468)	0	0	0	2	0	2	12.8	-	17.2
#Influenza & pneumonia (480-493)	6	0	6	8	0	8	51.2	-	69.0
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	12.8	49.8	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	0	1	6.4	-	8.6
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	2	0	2	12.8	-	17.2
Cirrhosis of liver (581)	2	1	1	2	1	1	12.8	24.9	8.6
Acute nephritis (590, 591)	1	0	1	1	0	1	6.4	-	8.6
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	0	1	1	0	1	6.4	-	8.6
Congenital malformations (750-759)	0	0	0	1	0	1	6.4	-	8.6
Diseases of early infancy (760-776)	6	0	6	6	1	5	15.5	16.4	15.3
Accidents (800-962)	18	9	9	12	4	8	76.8	99.5	69.0
Motor vehicle accidents (810-835)	13	7	6	5	1	4	32.0	24.9	34.5
Suicide (963, 970-979)	1	1	0	1	1	0	6.4	24.9	-
Homicide (964, 980-985)	1	0	1	1	0	1	6.4	-	8.6
Ill-defined (780-795)	13	0	13	16	0	16	102.4	-	137.9

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
UNION COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	345	247	98	345	248	97	17.9	15.0	35.4
Fetal deaths	13	8	5	8	6	2	23.2	24.2	20.6
Total deaths	191	159	32	201	170	31	10.4	10.3	11.3
Deaths under 1 year	10	8	2	8	6	2	23.2	24.2	20.6
Deaths under 28 days	8	7	1	6	5	1	17.4	20.2	10.3
<b>SELECTED CAUSES OF DEATH</b>									
<b>AND INTERNATIONAL LIST NO.</b>									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	1	1	0	5.2	6.0	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	24	22	2	29	27	2	150.6	163.5	73.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	1	1	0	5.2	6.0	-
Diabetes mellitus (260)	2	2	0	1	1	0	5.2	6.0	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	27	23	4	28	23	5	145.4	139.3	182.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	73	65	8	76	69	7	394.8	417.9	255.5
Other hypertensive disease (444-447)	1	1	0	0	0	0	-	-	-
General arteriosclerosis (450)	1	0	1	1	0	1	5.2	-	36.5
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	4	3	1	3	2	1	15.6	12.1	36.5
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	1	1	0	1	1	0	5.2	6.0	-
Stomach & duodenal ulcers (540, 541)	5	4	1	5	4	1	26.0	24.2	36.5
Hernia & intestinal obstruction (560, 561, 570)	1	1	0	2	2	0	10.4	12.1	-
#Gastritis, etc. (543, 571, 572)	2	2	0	2	2	0	10.4	12.1	-
Cirrhosis of liver (581)	0	0	0	1	1	0	5.2	6.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	3	2	1	3	2	1	15.6	12.1	36.5
Infections of kidney (600)	1	1	0	1	1	0	5.2	6.0	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	5.2	6.0	-
Diseases of early infancy (760-776)	7	6	1	4	3	1	11.6	12.1	10.3
Accidents (800-962)	12	10	2	14	13	1	72.7	78.7	36.5
Motor vehicle accidents (810-835)	7	6	1	9	8	1	46.8	48.4	36.5
Suicide (963, 970-979)	1	1	0	1	1	0	5.2	6.0	-
Homicide (964, 980-985)	0	0	0	0	0	0	-	-	-
Ill-defined (780-795)	22	12	10	22	12	10	114.3	72.7	365.0

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
WALTHALL COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence			Rate*		
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	328	160	168	268	99	169	18.9	12.9	26.0
Fetal deaths	16	8	8	13	7	6	48.5	70.7	35.5
Total deaths	123	67	56	144	84	60	10.1	10.9	9.2
Deaths under 1 year	15	3	12	15	5	10	56.0	50.5	59.2
Deaths under 28 days	8	2	6	10	5	5	37.3	50.5	29.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	2	1	1	14.1	13.0	15.4
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	12	6	6	22	14	8	155.0	182.3	122.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	7.0	-	15.4
Diabetes mellitus (260)	5	3	2	4	2	2	28.2	26.0	30.7
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	13	8	5	17	10	7	119.8	130.2	107.5
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	36	24	12	36	23	13	253.7	299.5	199.7
Other hypertensive disease (444-447)	2	0	2	2	0	2	14.1	-	30.7
General arteriosclerosis (450)	6	4	2	4	2	2	28.2	26.0	30.7
Other circulatory diseases (451-468)	1	0	1	3	3	0	21.1	39.1	-
#Influenza & pneumonia (480-493)	4	0	4	6	2	4	42.3	26.0	61.4
Bronchitis (500-502)	0	0	0	1	0	1	7.0	-	15.4
Other bronchopul. diseases (525-527)	4	2	2	5	3	2	35.2	39.1	30.7
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	1	0	1	0	0	0	-	-	-
Cirrhosis of liver (581)	2	2	0	2	2	0	14.1	26.0	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	1	1	0	1	1	0	37.3	101.0	-
Congenital malformations (750-759)	3	1	2	2	0	2	14.1	-	30.7
Diseases of early infancy (760-776)	5	1	4	8	4	4	29.8	40.4	23.7
Accidents (800-962)	13	8	5	11	8	3	77.5	104.2	46.1
Motor vehicle accidents (810-835)	7	5	2	6	4	2	42.3	52.1	30.7
Suicide (963, 970-979)	3	3	0	3	3	0	21.1	39.1	-
Homicide (964, 980-985)	0	0	0	1	1	0	7.0	13.0	-
Ill-defined (780-795)	6	2	4	6	2	4	42.3	26.0	61.4

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
WARREN COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,372	571	801	909	453	456	20.0	17.4	23.6
Fetal deaths	122	55	67	76	45	31	83.6	99.3	68.0
Total deaths	650	277	373	532	220	312	11.7	8.4	16.1
Deaths under 1 year	70	16	54	42	9	33	46.2	19.9	72.4
Deaths under 28 days	55	15	40	31	7	24	34.1	15.4	52.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	2.2	-	5.2
Syphilis (020-029)	1	0	1	1	0	1	2.2	-	5.2
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	1	0	1	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	93	45	48	80	34	46	176.3	130.6	237.9
Benign & unspec. neoplasms (210-239)	1	0	1	0	0	0	-	-	-
Asthma (241)	1	0	1	1	0	1	2.2	-	5.2
Diabetes mellitus (260)	15	3	12	11	3	8	24.2	11.5	41.4
Anemias (290-293)	2	0	2	1	0	1	2.2	-	5.2
Intracranial lesions (330-334)	69	30	39	68	29	39	149.9	111.4	201.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	178	87	91	158	72	86	348.2	276.5	444.8
Other hypertensive disease (444-447)	7	3	4	3	1	2	6.6	3.8	10.3
General arteriosclerosis (450)	16	10	6	10	6	4	22.0	23.0	20.7
Other circulatory diseases (451-468)	10	6	4	6	4	2	13.2	15.4	10.3
#Influenza & pneumonia (480-493)	17	10	7	15	10	5	33.0	38.4	25.8
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	7	6	1	8	6	2	17.6	23.0	10.3
Stomach & duodenal ulcers (540, 541)	2	2	0	3	3	0	6.6	11.5	-
Hernia & intestinal obstruction (560, 561, 570)	6	1	5	3	1	2	6.6	3.8	10.3
#Gastritis, etc. (543, 571, 572)	7	5	2	5	4	1	11.0	15.4	5.2
Cirrhosis of liver (581)	6	4	2	6	4	2	13.2	15.4	10.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	1	5	5	1	4	11.0	3.8	20.7
Infections of kidney (600)	3	1	2	2	1	1	4.4	3.8	5.2
Hyperplasia of prostate (610)	6	1	5	4	1	3	8.8	3.8	15.5
Maternal causes (640-689)	3	0	3	2	0	2	22.0	-	43.8
Congenital malformations (750-759)	6	3	3	3	1	2	6.6	3.8	10.3
Diseases of early infancy (760-776)	50	11	39	28	5	23	30.8	11.0	50.4
Accidents (800-962)	47	22	25	28	11	17	61.7	42.2	87.9
Motor vehicle accidents (810-835)	23	15	8	9	5	4	19.8	19.2	20.7
Suicide (963, 970-979)	3	2	1	3	2	1	6.6	7.7	5.2
Homicide (964, 980-985)	3	0	3	1	0	1	2.2	-	5.2
Ill-defined (780-795)	55	8	47	49	10	39	108.0	38.4	201.7

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
WASHINGTON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	1,962	658	1,304	1,851	586	1,265	23.7	16.5	29.7
Fetal deaths	126	67	59	125	57	68	67.5	97.3	53.8
Total deaths	770	292	478	799	276	523	10.2	7.8	12.3
Deaths under 1 year	68	10	58	74	12	62	40.0	20.5	49.0
Deaths under 28 days	41	6	35	47	8	39	25.4	13.6	30.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	5	3	2	5	2	3	6.4	5.6	7.0
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	91	51	40	100	48	52	128.0	135.1	122.0
Benign & unspec. neoplasms (210-239)	2	1	1	2	1	1	2.6	2.8	2.3
Asthma (241)	1	0	1	1	0	1	1.3	-	2.3
Diabetes mellitus (260)	14	1	13	16	1	15	20.5	2.8	35.2
Anemias (290-293)	1	0	1	2	1	1	2.6	2.8	2.3
Intracranial lesions (330-334)	63	32	31	61	27	34	78.1	76.0	79.8
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	1.3	-	2.3
Heart diseases (400-443)	228	102	126	224	92	132	286.7	258.9	309.8
Other hypertensive disease (444-447)	2	0	2	4	1	3	5.1	2.8	7.0
General arteriosclerosis (450)	6	3	3	4	1	3	5.1	2.8	7.0
Other circulatory diseases (451-468)	5	5	0	6	5	1	7.7	14.1	2.3
#Influenza & pneumonia (480-493)	28	13	15	26	11	15	33.3	31.0	35.2
Bronchitis (500-502)	1	1	0	1	1	0	1.3	2.8	-
Other bronchopul. diseases (525-527)	7	5	2	9	6	3	11.5	16.9	7.0
Stomach & duodenal ulcers (540, 541)	1	1	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	4	0	4	4	0	4	5.1	-	9.4
#Gastritis, etc. (543, 571, 572)	3	0	3	5	0	5	6.4	-	11.7
Cirrhosis of liver (581)	5	4	1	5	4	1	6.4	11.2	2.3
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	6	1	5	8	1	7	10.2	2.8	16.4
Infections of kidney (600)	2	0	2	2	0	2	2.6	-	4.7
Hyperplasia of prostate (610)	3	0	3	3	0	3	3.8	-	7.0
Maternal causes (640-689)	1	0	1	1	0	1	5.4	-	7.9
Congenital malformations (750-759)	8	3	5	11	4	7	14.1	11.2	16.4
Diseases of early infancy (760-776)	34	4	30	39	6	33	21.1	10.2	26.1
Accidents (800-962)	59	30	29	73	36	37	93.4	101.3	86.8
Motor vehicle accidents (810-835)	21	10	11	26	12	14	33.3	33.8	32.8
Suicide (963, 970-979)	6	6	0	5	5	0	6.4	14.1	-
Homicide (964, 980-985)	12	3	9	14	3	11	17.9	8.4	25.8
Ill-defined (780-795)	164	19	145	160	16	144	204.8	45.0	337.9

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
 AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
 WAYNE COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	371	260	111	327	184	143	19.7	17.5	23.5
Fetal deaths	15	10	5	14	8	6	42.8	43.5	42.0
Total deaths	156	92	64	185	113	72	11.2	10.8	11.8
Deaths under 1 year	8	4	4	8	4	4	24.5	21.7	28.0
Deaths under 28 days	5	2	3	5	2	3	15.3	10.9	21.0
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	1	0	1	1	0	1	6.0	-	16.4
Syphilis (020-029)	0	0	0	2	2	0	12.0	19.0	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	25	14	11	26	15	11	156.7	142.7	180.9
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	3	1	2	3	1	2	18.1	9.5	32.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	25	14	11	27	16	11	162.7	152.2	180.9
Nonmeningococcal meningitis (340)	0	0	0	1	0	1	6.0	-	16.4
Heart diseases (400-443)	61	41	20	62	40	22	373.7	380.6	361.8
Other hypertensive disease (444-447)	1	0	1	1	0	1	6.0	-	16.4
General arteriosclerosis (450)	4	2	2	5	3	2	30.1	28.5	32.9
Other circulatory diseases (451-468)	2	2	0	2	2	0	12.0	19.0	-
#Influenza & pneumonia (480-493)	0	0	0	4	4	0	24.1	38.0	-
Bronchitis (500-502)	1	1	0	1	1	0	6.0	9.5	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	12.0	19.0	-
Stomach & duodenal ulcers (540, 541)	2	1	1	1	0	1	6.0	-	16.4
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	0	0	0	-	-	-
Cirrhosis of liver (581)	1	0	1	1	0	1	6.0	-	16.4
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	1	1	3	1	2	18.1	9.5	32.9
Infections of kidney (600)	1	0	1	2	0	2	12.0	-	32.9
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	0	0	0	-	-	-
Diseases of early infancy (760-776)	5	2	3	5	2	3	15.3	10.9	21.0
Accidents (800-962)	10	7	3	20	15	5	120.6	142.7	82.2
Motor vehicle accidents (810-835)	1	0	1	7	5	2	42.2	47.6	32.9
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	3	1	2	4	1	3	24.1	9.5	49.3
III-defined (780-795)	5	2	3	5	2	3	30.1	19.0	49.3

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
WEBSTER COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	120	75	45	162	100	62	15.4	12.5	24.5
Fetal deaths	2	1	1	4	2	2	24.7	20.0	32.2
Total deaths	112	95	17	123	104	19	11.7	13.0	7.5
Deaths under 1 year	4	1	3	8	2	6	49.4	20.0	96.8
Deaths under 28 days	4	1	3	5	2	3	30.9	20.0	48.4
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	9.5	12.5	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	1	0	1	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	1	1	0	9.5	12.5	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	11	10	1	12	10	2	114.2	125.3	79.0
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	0	0	0	1	1	0	9.5	12.5	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	22	18	4	28	24	4	266.4	300.8	158.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	42	38	4	40	35	5	380.6	438.6	197.6
Other hypertensive disease (444-447)	0	0	0	1	1	0	9.5	12.5	-
General arteriosclerosis (450)	5	4	1	3	2	1	28.5	25.1	39.5
Other circulatory diseases (451-468)	0	0	0	2	2	0	19.0	25.1	-
#Influenza & pneumonia (480-493)	1	1	0	4	4	0	38.0	50.1	-
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	2	0	2	2	0	19.0	25.1	-
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	1	0	1	9.5	-	39.5
#Gastritis, etc. (543, 571, 572)	0	0	0	1	0	1	9.5	-	39.5
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	4	4	0	3	3	0	28.5	37.6	-
Hyperplasia of prostate (610)	1	0	1	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	1	1	0	9.5	12.5	-
Diseases of early infancy (760-776)	3	1	2	4	2	2	24.7	20.0	32.2
Accidents (800-962)	8	7	1	6	5	1	57.1	62.6	39.5
Motor vehicle accidents (810-835)	8	7	1	2	1	1	19.0	12.5	39.5
Suicide (963, 970-979)	4	4	0	4	4	0	38.0	50.1	-
Homicide (964, 980-985)	1	1	0	2	2	0	19.0	25.1	-
Ill-defined (780-795)	3	1	2	3	1	2	28.5	12.5	79.0

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
WILKINSON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	230	87	143	231	58	173	20.0	16.9	21.3
Fetal deaths	6	1	5	9	2	7	39.0	34.5	40.5
Total deaths	109	50	59	135	54	81	11.7	15.7	10.0
Deaths under 1 year	5	0	5	8	2	6	34.6	34.5	34.7
Deaths under 28 days	1	0	1	4	1	3	17.3	17.2	17.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	17	11	6	18	10	8	155.7	291.5	98.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	0	1	3	0	3	26.0	-	36.9
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	9	2	7	16	5	11	138.4	145.8	135.3
Nonmeningococcal meningitis (340)	1	0	1	1	0	1	8.6	-	12.3
Heart diseases (400-443)	32	19	13	33	19	14	285.5	553.9	172.2
Other hypertensive disease (444-447)	1	0	1	2	0	2	17.3	-	24.6
General arteriosclerosis (450)	2	1	1	4	0	4	34.6	-	49.2
Other circulatory diseases (451-468)	1	1	0	2	1	1	17.3	29.2	12.3
#Influenza & pneumonia (480-493)	3	1	2	4	1	3	34.6	29.2	36.9
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	2	1	1	4	1	3	34.6	29.2	36.9
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	1	0	1	1	0	1	8.6	-	12.3
#Gastritis, etc. (543, 571, 572)	1	0	1	1	0	1	8.6	-	12.3
Cirrhosis of liver (581)	0	0	0	1	1	0	8.6	29.2	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	0	1	1	0	1	8.6	-	12.3
Hyperplasia of prostate (610)	0	0	0	1	0	1	8.6	-	12.3
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	3	2	1	26.0	58.3	12.3
Diseases of early infancy (760-776)	0	0	0	1	0	1	4.3	-	5.8
Accidents (800-962)	17	11	6	17	10	7	147.0	291.5	86.1
Motor vehicle accidents (810-835)	9	7	2	10	7	3	86.5	204.1	36.9
Suicide (963, 970-979)	0	0	0	0	0	0	-	-	-
Homicide (964, 980-985)	2	0	2	3	0	3	26.0	-	36.9
III-defined (780-795)	15	2	13	14	2	12	121.1	58.3	147.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
WINSTON COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	332	142	190	378	153	225	22.1	16.1	29.5
Fetal deaths	9	4	5	11	4	7	29.1	26.1	31.1
Total deaths	156	110	46	187	126	61	10.9	13.3	8.0
Deaths under 1 year	13	4	9	12	4	8	31.7	26.1	35.6
Deaths under 28 days	6	3	3	6	3	3	15.9	19.6	13.3
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	0	0	0	-	-	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	15	10	5	21	16	5	122.7	168.6	65.5
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	2	1	1	2	1	1	11.7	10.5	13.1
Anemias (290-293)	1	1	0	0	0	0	-	-	-
Intracranial lesions (330-334)	33	27	6	40	30	10	233.6	316.1	131.1
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	58	44	14	62	45	17	362.1	474.2	222.8
Other hypertensive disease (444-447)	1	1	0	1	1	0	5.8	10.5	-
General arteriosclerosis (450)	5	3	2	9	4	5	52.6	42.1	65.5
Other circulatory diseases (451-468)	2	2	0	2	2	0	11.7	21.1	-
#Influenza & pneumonia (480-493)	4	2	2	5	2	3	29.2	21.1	39.3
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	2	1	1	11.7	10.5	13.1
Stomach & duodenal ulcers (540, 541)	0	0	0	0	0	0	-	-	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	2	0	2	2	0	2	11.7	-	26.2
Cirrhosis of liver (581)	0	0	0	1	0	1	5.8	-	13.1
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	2	0	2	4	1	3	23.4	10.5	39.3
Infections of kidney (600)	0	0	0	0	0	0	-	-	-
Hyperplasia of prostate (610)	1	1	0	1	1	0	5.8	10.5	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	1	1	0	1	1	0	5.8	10.5	-
Diseases of early infancy (760-776)	6	2	4	5	2	3	13.2	13.1	13.3
Accidents (800-962)	14	10	4	19	13	6	111.0	137.0	78.6
Motor vehicle accidents (810-835)	6	3	3	10	5	5	58.4	52.7	65.5
Suicide (963, 970-979)	2	2	0	3	3	0	17.5	31.6	-
Homicide (964, 980-985)	1	1	0	0	0	0	-	-	-
Ill-defined (780-795)	4	0	4	3	0	3	17.5	-	39.3

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
YALOBUSHA COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	210	74	136	250	102	148	20.5	15.2	26.9
Fetal deaths	6	1	5	9	3	6	36.0	29.4	40.5
Total deaths	123	76	47	164	107	57	13.4	16.0	10.3
Deaths under 1 year	7	1	6	11	4	7	44.0	39.2	47.3
Deaths under 28 days	2	0	2	5	3	2	20.0	29.4	13.5
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	1	0	8.2	14.9	-
Syphilis (020-029)	0	0	0	0	0	0	-	-	-
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	18	14	4	21	16	5	172.1	239.2	90.7
Benign & unspec. neoplasms (210-239)	0	0	0	1	1	0	8.2	14.9	-
Asthma (241)	0	0	0	0	0	0	-	-	-
Diabetes mellitus (260)	1	1	0	2	2	0	16.4	29.9	-
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	26	11	15	36	18	18	295.1	269.0	326.7
Nonmeningococcal meningitis (340)	0	0	0	0	0	0	-	-	-
Heart diseases (400-443)	39	29	10	47	35	12	385.2	523.2	217.8
Other hypertensive disease (444-447)	0	0	0	1	0	1	8.2	-	18.1
General arteriosclerosis (450)	0	0	0	1	1	0	8.2	14.9	-
Other circulatory diseases (451-468)	0	0	0	0	0	0	-	-	-
#Influenza & pneumonia (480-493)	8	6	2	10	8	2	82.0	119.6	36.3
Bronchitis (500-502)	1	1	0	2	2	0	16.4	29.9	-
Other bronchopul. diseases (525-527)	1	1	0	2	2	0	16.4	29.9	-
Stomach & duodenal ulcers (540, 541)	1	1	0	1	1	0	8.2	14.9	-
Hernia & intestinal obstruction (560, 561, 570)	0	0	0	0	0	0	-	-	-
#Gastritis, etc. (543, 571, 572)	0	0	0	1	1	0	8.2	14.9	-
Cirrhosis of liver (581)	0	0	0	0	0	0	-	-	-
Acute nephritis (590, 591)	0	0	0	0	0	0	-	-	-
Chronic nephritis, etc. (592-594)	0	0	0	0	0	0	-	-	-
Infections of kidney (600)	1	1	0	1	1	0	8.2	14.9	-
Hyperplasia of prostate (610)	0	0	0	0	0	0	-	-	-
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	1	1	0	8.2	14.9	-
Diseases of early infancy (760-776)	2	0	2	3	1	2	12.0	9.8	13.5
Accidents (800-962)	10	7	3	13	9	4	106.6	134.5	72.6
Motor vehicle accidents (810-835)	6	3	3	9	5	4	73.8	74.7	72.6
Suicide (963, 970-979)	1	1	0	1	1	0	8.2	14.9	-
Homicide (964, 980-985)	3	0	3	3	0	3	24.6	-	54.4
III-defined (780-795)	7	1	6	10	3	7	82.0	44.8	127.0

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS,  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
YAZOO COUNTY, MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	589	192	397	699	221	478	20.4	15.4	24.0
Fetal deaths	28	4	24	41	7	34	58.6	31.7	71.1
Total deaths	308	133	175	381	159	222	11.1	11.1	11.1
Deaths under 1 year	19	5	14	31	5	26	44.3	22.6	54.4
Deaths under 28 days	15	4	11	22	4	18	31.5	18.1	37.6
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	0	0	0	1	0	1	2.9	-	5.0
Syphilis (020-029)	1	0	1	1	0	1	2.9	-	5.0
Dysentery (045-048)	0	0	0	0	0	0	-	-	-
Scarlet fever & streptococcal sore throat (050, 051)	0	0	0	0	0	0	-	-	-
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	0	0	0	0	0	0	-	-	-
Meningococcal infections (057)	0	0	0	0	0	0	-	-	-
Acute poliomyelitis (080)	0	0	0	0	0	0	-	-	-
Measles (085)	0	0	0	0	0	0	-	-	-
Malignant neoplasms (140-205)	43	23	20	56	27	29	163.5	188.8	145.4
Benign & unspec. neoplasms (210-239)	0	0	0	0	0	0	-	-	-
Asthma (241)	1	1	0	1	1	0	2.9	7.0	-
Diabetes mellitus (260)	2	0	2	2	1	1	5.8	7.0	5.0
Anemias (290-293)	0	0	0	0	0	0	-	-	-
Intracranial lesions (330-334)	30	16	14	38	19	19	110.9	132.9	95.2
Nonmeningocephal meningitis (340)	1	0	1	1	0	1	2.9	-	5.0
Heart diseases (400-443)	93	48	45	109	56	53	318.2	391.6	265.7
Other hypertensive disease (444-447)	2	1	1	3	1	2	8.8	7.0	10.0
General arteriosclerosis (450)	4	3	1	6	5	1	17.5	35.0	5.0
Other circulatory diseases (451-468)	1	1	0	4	2	2	11.7	14.0	10.0
#Influenza & pneumonia (480-493)	9	1	8	11	2	9	32.1	14.0	45.1
Bronchitis (500-502)	0	0	0	0	0	0	-	-	-
Other bronchopul. diseases (525-527)	0	0	0	0	0	0	-	-	-
Stomach & duodenal ulcers (540, 541)	0	0	0	1	1	0	2.9	7.0	-
Hernia & intestinal obstruction (560, 561, 570)	2	1	1	3	1	2	8.8	7.0	10.0
#Gastritis, etc. (543, 571, 572)	1	0	1	2	1	1	5.8	7.0	5.0
Cirrhosis of liver (581)	1	1	0	1	1	0	2.9	7.0	-
Acute nephritis (590, 591)	1	0	1	1	0	1	2.9	-	5.0
Chronic nephritis, etc. (592-594)	4	0	4	6	0	6	17.5	-	30.1
Infections of kidney (600)	1	0	1	1	1	0	2.9	7.0	-
Hyperplasia of prostate (610)	1	0	1	1	0	1	2.9	-	5.0
Maternal causes (640-689)	0	0	0	0	0	0	-	-	-
Congenital malformations (750-759)	0	0	0	2	1	1	5.8	7.0	5.0
Diseases of early infancy (760-776)	12	4	8	18	4	14	25.8	18.1	29.3
Accidents (800-962)	30	14	16	36	15	21	105.1	104.9	105.3
Motor vehicle accidents (810-835)	16	8	8	20	10	10	58.4	69.9	50.1
Suicide (963, 970-979)	2	2	0	2	2	0	5.8	14.0	-
Homicide (964, 980-985)	2	0	2	3	0	3	8.8	-	15.0
Ill-defined (780-795)	58	11	47	59	11	48	172.3	76.9	240.6

See page 55 for footnotes.

BIRTHS, FETAL DEATHS, TOTAL DEATHS, DEATHS UNDER 1 YEAR, DEATHS UNDER 28 DAYS  
AND DEATHS FROM SELECTED CAUSES USED FOR RANKING LEADING CAUSES OF DEATH  
STATE OF MISSISSIPPI, 1967

By Place of Occurrence, and by Place of Residence, by Race, with Rates

	Occurrence			Residence					
	Total	White	Non-white	Number			Rate*		
				Total	White	Non-white	Total	White	Non-white
Live births	46714	22765	23949	46594	22675	23919	19.7	15.9	25.6
Fetal deaths	2321	1065	1256	2291	1031	1260	49.2	45.5	52.7
Total deaths	22121	12538	9583	22609	12816	9793	9.6	9.0	10.5
Deaths under 1 year	1614	497	1117	1645	512	1133	35.3	22.6	47.4
Deaths under 28 days	1040	383	657	1053	388	665	22.6	17.1	27.8
SELECTED CAUSES OF DEATH AND INTERNATIONAL LIST NO.									
Tuberculosis (001-019)	78	36	42	77	36	41	3.3	2.5	4.4
Syphilis (020-029)	18	8	10	22	11	11	0.9	0.8	1.2
Dysentery (045-048)	7	1	6	7	1	6	0.3	0.1	0.6
Scarlet fever & streptococcal sore throat (050, 051)	2	0	2	2	0	2	0.1	-	0.2
Diphtheria (055)	0	0	0	0	0	0	-	-	-
Whooping cough (056)	3	1	2	3	1	2	0.1	0.1	0.2
Meningococcal infections (057)	5	3	2	6	3	3	0.2	0.2	0.3
Acute poliomyelitis (080)	1	0	1	1	0	1	**	-	0.1
Measles (085)	3	1	2	3	1	2	0.1	0.1	0.2
Malignant neoplasms (140-205)	3000	1966	1034	3136	2061	1075	132.8	144.4	115.1
Benign & unspec. neoplasms (210-239)	36	17	19	39	20	19	1.6	1.4	2.0
Asthma (241)	51	30	21	50	29	21	2.1	2.0	2.2
Diabetes mellitus (260)	326	141	185	332	146	186	14.1	10.2	19.9
Anemias (290-293)	48	12	36	48	12	36	2.0	0.8	3.8
Intracranial lesions (330-334)	2880	1567	1313	2981	1632	1349	126.2	114.4	144.4
Nonmeningococcal meningitis (340)	44	12	32	47	14	33	2.0	1.0	3.5
Heart diseases (400-443)	7223	4607	2616	7300	4640	2660	309.2	325.2	284.7
Other hypertensive disease (444-447)	196	62	134	196	65	131	8.3	4.6	14.0
General arteriosclerosis (450)	339	218	121	348	221	127	14.7	15.5	13.6
Other circulatory diseases (451-468)	264	180	84	288	191	97	12.2	13.4	10.4
#Influenza & pneumonia (480-493)	649	312	337	657	310	347	27.8	21.7	37.1
Bronchitis (500-502)	51	44	7	50	43	7	2.1	3.0	0.7
Other bronchopul. diseases (525-527)	268	209	59	275	214	61	11.6	15.0	6.5
Stomach & duodenal ulcers (540, 541)	85	56	29	88	58	30	3.7	4.1	3.2
Hernia & intestinal obstruction (560, 561, 570)	104	59	45	111	64	47	4.7	4.5	5.0
#Gastritis, etc. (543, 571, 572)	158	68	90	164	74	90	6.9	5.2	9.6
Cirrhosis of liver (581)	128	87	41	132	90	42	5.6	6.3	4.5
Acute nephritis (590, 591)	25	7	18	25	7	18	1.0	0.5	1.9
Chronic nephritis, etc. (592-594)	180	53	127	185	57	128	7.8	4.0	13.7
Infections of kidney (600)	150	90	60	156	93	63	6.6	6.5	6.7
Hyperplasia of prostate (610)	52	18	34	56	19	37	2.4	1.3	4.0
Maternal causes (640-689)	32	8	24	36	10	26	7.7	4.4	10.9
Congenital malformations (750-759)	195	103	92	227	123	104	9.6	8.6	11.1
Diseases of early infancy (760-776)	872	307	565	875	308	567	18.8	13.6	23.7
Accidents (800-962)	1861	1151	710	1890	1166	724	80.0	81.7	77.5
Motor vehicle accidents (810-835)	944	662	282	931	649	282	39.4	45.5	30.2
Suicide (963, 970-979)	159	145	14	164	149	15	6.9	10.4	1.6
Homicide (964, 980-985)	261	66	195	271	66	205	11.5	4.6	21.9
Ill-defined (780-795)	1627	455	1172	1601	436	1165	67.8	30.6	124.7

See page 55 for footnotes.